

Growth Factor-Mediated Cell–Cell Interactions in the Ovary

MICHAEL K. SKINNER & JEFF A. PARROTT

I. Introduction

The local production and integrated actions of various growth factors are required for the growth and development of all tissues. Growth factors mediate critical cell–cell interactions that control cell proliferation and organ development and the number of specific growth factors identified has increased dramatically. Growth factors often exist in families composed of unique gene products that have similarities in structure and function, but often differ in the way gene expression is controlled. The existence of multiple members in a growth factor family allows for unique sites of expression and more precise developmental and hormonal regulation of growth factor production. Table 1 contains a partial list of several of the major types of growth factors, including nomenclature and major functions attributed to them.

Ovarian development also requires growth factor-mediated cell–cell interactions as a general mechanism for controlling cellular proliferation. Most of the information available on growth factors and the ovary deals with follicular development and adult ovarian function. These are the primary stages of development discussed here. However, it is likely that many of the same cell–cell interactions and growth factors are also important for other stages of ovarian development (e.g., embryonic or prepubertal stages). Since research has been focused primarily on identifying the sites of production and actions of specific growth factors, this chapter emphasizes the growth factors potentially involved in ovarian cell–cell interac-

TABLE 1 Properties and Nomenclature of Several Common Growth Factors

Growth Factor		Approx. size (kDA)	Examples of physiological action
Insulin-like growth factor-I	IGF-I	7.5	Skeletal growth
Insulin-like growth factor-II	IGF-II	7.5	Fetal development
Epidermal growth factor	EGF	6	Tissue growth
Transforming growth factor α	TGF α	5	Tissue growth
Transforming growth factor β	TGF β	25/dimer	Growth inhibition/tissue repair
Fibroblast growth factor	FGF	17	Angiogenesis/tissue growth
Vascular endothelial growth factor	VEGF	25-50/dimer	Angiogenesis/tissue growth
Nerve growth factor	NGF	13	Neuronal development
Interleukin-1	IL-1	17	Immune response/inflammation
Platelet-derived growth factor	PDGF	30/dimer	Tissue growth
Stem cell factor (c-kit ligand)	SCF	30	Tissue growth/fetal development

tions. The specific functions of these growth factors *in vivo* in large part remain to be elucidated.

II. Growth Factors in the Ovary

Ovarian physiology requires rapid and continuous regulation of the growth associated with folliculogenesis. Growth factor-mediated interactions among theca cells, granulosa cells, and the oocyte are needed to maintain ovarian function and oogenesis. Granulosa cells provide the cyto-architectural support for the developing oocyte and also help to form the follicle and antrum. Theca cells surround and form the exterior wall of the follicle. The interactions between theca cells and granulosa cells provide an example of a mesenchymal (stromal)-epithelial cell interaction. The effects of this cellular interaction on oocyte development and the influence of the endocrine system on this cellular interaction are essential to female reproductive biology. The theca cells and granulosa cells of the preantral and small antral follicles must undergo extensive proliferation and functional differentiation prior to ovulation (Hsueh *et al.*, 1984). In most large animals, follicle size increases from millimeters to centimeters. The primary expansion in cell number takes place in both the granulosa and theca cell populations. In addition to the cell proliferation required during follicle development, follicles at various stages of development become atretic and cell growth is arrested. Therefore, the regulation of cell proliferation in the follicle requires both stimulatory and inhibitory growth factors.

TABLE 2 Growth Factors in the Ovary

Growth factor	Proposed site of synthesis	Proposed site of action	Proposed function ^a
IGF-1	Granulosa	Granulosa Theca	+Growth/+differentiation +Growth/+differentiation
FGF	Granulosa	Granulosa Endothelium	+Growth Angiogenesis
TGF β	Theca Granulosa	Granulosa Theca	-Growth/+differentiation -Growth/+differentiation
TGF α	Theca	Granulosa Theca	+Growth/-differentiation +growth
VEGF	Granulosa	Endothelium	Angiogenesis/+growth
NGF	Ovary	Neurons	Innervation
SCF	Granulosa	Oocyte	Oocyte maturation

^aA plus sign denotes an increase and a minus sign indicates a decrease.

Follicle-stimulating hormone (FSH) and estrogen have been shown to stimulate proliferation of follicle cells *in vivo* (Goldenberg *et al.*, 1972; Louvet and Vaitukaitis, 1976; Richards, 1979). These hormones, however, have negligible effects on cell growth *in vitro*. This implies that *in vivo* hormone actions are most likely indirectly mediated by the local production of growth factors. The mechanisms and specific growth factors involved in the control of ovarian cell proliferation remain to be fully elucidated (Carson *et al.*, 1989). However, several major growth factors have been identified as being produced and/or acting in the ovary. These are summarized in Table 2.

A. Insulin-like Growth Factor

The insulin-like growth factors (IGF) derive their name from their structural similarity to insulin (Froesch *et al.*, 1985). IGF-I (previously termed somatomedin C) is considered essential for cellular replication and is a progression factor for cell growth and DNA synthesis. Production and secretion of IGF-I by the liver accounts for the high levels of IGF-I in serum and interstitial fluid (Daughaday and Rotwein, 1989). IGF-II is another member of this family and may act as a growth factor during fetal development.

IGF-I is produced by granulosa cells under the control of growth hormone (Davoren and Hsueh 1986), FSH, and estradiol (Hammond *et al.*, 1985; Hsu and Hammond, 1987). The gene expression of IGF-1 has also been localized in granulosa cells (Hernandez *et al.*, 1989; Oliver *et al.*, 1989) but not theca cells. The receptors for IGF-1 have been localized to granulosa cells (Baranao and Hammond, 1984; Davoren *et al.*, 1986; Adashi *et al.*,

1988b) and are affected by the actions of FSH (Adashi *et al.*, 1986, 1988c,d). The regulation of IGF receptor gene expression has also been examined (Hernandez *et al.*, 1991, 1992). IGF-1 stimulates granulosa cell oxytocin production (Schams *et al.*, 1988), the P450 side chain cleavage enzyme (Veldhuis *et al.*, 1986), the aromatase gene expression (Steinkampf *et al.*, 1988), lipoprotein metabolism (Veldhuis *et al.*, 1987), adenylate cyclase activity (Adashi *et al.*, 1986), plasminogen activator production (Tilly and Johnson, 1990), and LH receptor induction (Adashi *et al.*, 1985b).

In addition to affecting cellular function, IGF stimulates the proliferation of bovine (Savion *et al.*, 1981) and porcine (Baranao and Hammond, 1984) granulosa cells, but not rat granulosa cells *in vitro* (Adashi *et al.*, 1984). Although theca cells do not appear to produce IGF-1 (Oliver *et al.*, 1989), IGF-II gene expression has been localized to theca cells and not granulosa cells (Hernandez *et al.*, 1990a). Theca cells also contain IGF receptors and respond to IGF through an alteration in steroidogenesis (Hernandez *et al.*, 1988) and LH receptor binding (Cara *et al.*, 1990). Potential interactions between granulosa cells and theca cells through the local production and action of IGF have been suggested (Adashi *et al.*, 1985a; Geisthovel *et al.*, 1990). The localization of IGF-1 expression to granulosa and not theca cells implies a potential IGF-1-mediated paracrine interaction between granulosa and theca cells. IGF-1 can also play a role as an autocrine factor for granulosa cells. A physiological parameter to consider, however, is the high circulatory levels of liver-derived IGF-1 (>100 ng/ml) available to both cell types. This is an additional source of IGF-1 that needs to be considered in understanding the importance of IGF-mediated cell-cell interactions.

IGF binding proteins (IGFBP) are produced by ovarian cell types and are present in the follicle. These binding proteins can reduce the effective concentration and modulate the actions of IGF. Several forms of IGFBP have been identified and are present in the ovary. Both granulosa cells and theca cells produce IGFBP(2) (Samares *et al.*, 1992; Ricciarelli *et al.*, 1991; Nakatani *et al.*, 1991), IGFBP(3) (Ricciarelli *et al.*, 1992; Samares *et al.*, 1992; Mondschein *et al.*, 1990), and IGFBP(4), which appears to be species-specific for cellular localization. IGFBP(4) and IGFBP(5) appear to be expressed primarily by granulosa cells (Erickson *et al.*, 1992a,b; Nakatani *et al.*, 1991). All the forms are present in follicular fluid at various stages of development. Although the specific function(s) of these IGFBPs remains to be elucidated, it has been postulated that they may inhibit or control the actions of IGF (Ui *et al.*, 1989).

B. Transforming Growth Factor- α /Epidermal Growth Factor

Transforming growth factor- α (TGF α) is one of the structurally related peptides belonging to the epidermal growth factor (EGF) family (Derynck, 1988; Carpenter and Cohen, 1990). Because they have a similar protein

structure, these factors act at the same receptor to stimulate cell growth (Carpenter, 1987). TGF α is synthesized as a transmembrane precursor, which may activate EGF receptors on neighboring cells or be proteolytically cleaved to release mature peptide. TGF α appears to be produced by non-transformed cells, and may play an important role as a growth regulator in normal tissues.

Although EGF was not found to be produced in the ovary, an EGF-like substance was found in theca cells (Skinner *et al.*, 1987b) and was identified as TGF α (Skinner and Coffey, 1988; Kudlow *et al.*, 1987). Granulosa cells do not express TGF α (Skinner and Coffey, 1988; Lobb *et al.*, 1989) but have been shown to contain the EGF receptor (Mondschein and Schomberg, 1981; Chabot *et al.*, 1986; Feng *et al.*, 1986). EGF generally is inhibitory for adenylate cyclase (Dodson and Schomberg, 1987), LH receptor activity (Mondschein and Schomberg, 1981; Knecht and Catt, 1983a; May *et al.*, 1987), and FSH-induced aromatase activity (Hsueh *et al.*, 1981; May *et al.*, 1982). TGF α has inhibitory effects on granulosa cells (Adashi *et al.*, 1987) that are similar to those of EGF. The effects of TGF α , however, can vary among species (Gangrade *et al.*, 1991). Theca cells also contain the EGF receptor (Skinner and Coffey, 1988), and EGF/TGF α influences theca cell steroidogenesis (Erickson and Case, 1983). One of the initial observations on growth regulation in the ovary was the ability of EGF to stimulate granulosa cell proliferation (Gospodarowicz *et al.*, 1977). The growth of theca cells can also be stimulated by TGF α /EGF (Skinner and Coffey, 1988). The actions of TGF α /EGF and the potential presence of EGF in the ovaries of various species can vary and remain to be fully elucidated.

Circulatory levels of EGF/TGF α are negligible, therefore, the factors must be produced locally within a specific organ. The ability of the theca cell to produce TGF α that can stimulate the growth of both granulosa and theca cells implies that TGF α may have an important role in promoting cell proliferation during follicle development. TGF α has been localized in developing ovarian follicles (Lobb *et al.*, 1989; Chegini and Williams, 1992). An interesting observation is that this appears to be a mesenchymal/stromal-controlled growth process (Skinner, 1990). Therefore, TGF α is postulated to mediate a paracrine interaction between theca and granulosa cells and an autocrine interaction between theca cells. The ability of hormones to influence TGF α production remains to be elucidated and may provide a mechanism through which hormones can regulate ovarian follicle cell growth.

C. Transforming Growth Factor- β

Transforming growth factor- β (TGF β) is a multifunctional regulatory molecule that can stimulate or inhibit aspects of cellular growth and differentiation (Roberts and Sporn, 1988). TGF β acts as a growth inhibitor by

inhibiting the actions of growth factors such as EGF/TGF α . TGF β can also promote cellular differentiation, extracellular matrix production, and chemotaxis. Different subtypes of TGF β are produced as latent secreted precursors. Most cell types contain receptors for this ubiquitous factor.

Skinner *et al.* (1987a) and Gangrade and May (1990) demonstrated that ovarian theca cells express and produce TGF β *in vivo*, immunocytochemical localization of TGF β is primarily confined to the theca cell layer (Thompson *et al.*, 1989). Several recent studies have confirmed the immunocytochemical localization of TGF β isoforms in follicle cells (Chegini and Williams, 1992; Chegini and Flanders, 1992); this may vary with follicle development and hormone treatment (Roy *et al.*, 1992). Although freshly isolated bovine granulosa cells do not appear to express TGF β (Skinner *et al.*, 1987a), cultured rat granulosa cells produce TGF β which can be suppressed by FSH (Kim and Schomberg, 1989; Mulheron and Schomberg, 1990). Therefore, theca cells appear to be a predominant source of ovarian TGF β , but granulosa cells also have the capacity to express TGF β . The specific types of TGF β expressed and their hormonal and developmental regulation remain to be fully elucidated (Mulheron *et al.*, 1991, 1992).

Local production of TGF β allows it to act on various ovarian cell types. TGF β stimulates a number of granulosa cell functions, including FSH-induced LH receptors (Knecht *et al.*, 1986; Dodson and Schomberg, 1987), EGF actions (Feng *et al.*, 1986), FSH-induced aromatase activity (Ying *et al.*, 1986; Hutchinson *et al.*, 1987), IGF-1 production (Mondschein *et al.*, 1988), and inhibin production (Zhiwen *et al.*, 1988). TGF β can also influence theca cell function and steroidogenesis (Magoffin *et al.*, 1989; Caubo *et al.*, 1989; Hernandez *et al.*, 1990b), and oocyte maturation (Feng *et al.*, 1988; Tsafiriri *et al.*, 1989). However, it is not known if TGF β acts directly or indirectly on oocytes. In addition to effects on cellular differentiation, TGF β can also influence ovarian cell growth. TGF β has been shown to inhibit TGF α /EGF-induced bovine and porcine granulosa cell growth (Skinner *et al.*, 1987a; Mondschein *et al.*, 1988). Conflicting data were found with rat granulosa cells (Dorrington *et al.*, 1988). TGF β can also inhibit TGF α /EGF-induced theca cell growth (Roberts and Skinner, 1991). Observations suggest that TGF β may have an important role as a growth inhibitor in the ovary. The ability of TGF β to inhibit cell growth allows for a more differentiated state of the cell that is reflected in the generally stimulatory effects of TGF β on cell function. Therefore, the influence of TGF β on cell function may be indirectly mediated through the inhibition of cellular proliferation. Growth inhibition may be important in preventing premature cell growth of the preantral follicle, arresting cell growth during atresia, and controlling cell growth during follicle cell expansion.

The local production and action of TGF β within the developing ovarian follicle implies that TGF β is an important paracrine and autocrine factor for ovarian cell-cell interactions. The hormonal regulation of TGF β production

(Bendell and Dorrington, 1991) may also have a role in the endocrine regulation of ovary growth. The physiological significance of TGF β in the ovarian follicle remains to be elucidated.

D. Fibroblast Growth Factor

Fibroblast growth factor (FGF) can influence aspects of both cellular growth and differentiation (Gospodarowicz *et al.*, 1987). Aside from growth stimulation, recent studies indicate that FGF may play a critical role in angiogenesis and tissue repair. FGF has many cellular targets and is important in many organ systems, including the ovary (Gospodarowicz and Ferrara, 1989).

Basic FGF is produced by granulosa cells in the developing embryonic gonad (Gonzalez *et al.*, 1990) and in the adult ovary (Neufeld *et al.*, 1987; Koos and Olson, 1989; Guthridge *et al.*, 1992). The angiogenic factor in the ovary and corpus luteum has been identified as FGF (Gospodarowicz *et al.*, 1985). FGF can act on granulosa cells to alter the steroidogenic capacity of the cell (Baird and Hsueh, 1986; Adashi *et al.*, 1988a), gonadotropin receptors (Mondschein and Schomberg, 1981), and plasminogen activator expression (LaPolt *et al.*, 1990; Tilly and Johnson, 1990). In addition to these effects on cell function, FGF can stimulate granulosa cell proliferation (Gospodarowicz *et al.*, 1977; Gospodarowicz and Bialecki, 1979), and may indirectly cause many of the effects observed on cell function. An additional role for FGF production in the ovary is to act as an angiogenic factor and promote vascularization of the developing follicle and corpus luteum.

E. Platelet-Derived Growth Factor and Vascular Endothelial Growth Factor

Platelet-derived growth factor (PDGF) is a common growth factor that allows cells to become competent to enter the growth cycle. PDGF acts on granulosa cells to enhance FSH-induced progesterone secretion, adenylate cyclase activity (Knecht and Catt, 1983b), plasminogen activator production (Tilly and Johnson, 1990), and LH receptor induction (Knecht and Catt, 1983b; Mondschein and Schomberg, 1984). The effects of PDGF on ovarian cell growth remain to be elucidated, but action as a potential competence factor for cell proliferation is a plausible activity. The local production of PDGF in the ovary remains to be examined. A factor that is structurally related to PDGF has been identified as vascular endothelial growth factor (VEGF) (Leung *et al.*, 1989; Conn *et al.*, 1990). This growth factor is expressed in the ovary, particularly in luteal tissue, and is postulated to have a role in angiogenesis of the follicle (Phillips *et al.*, 1990; Ravindranath *et al.*, 1992).

F. Nerve Growth Factor

Nerve growth factor (NGF) is another mitogen that may mediate intercellular interactions involving growth (Yanker and Shooter, 1982). NGF is important for the development and maintenance of sympathetic neurons in the peripheral nervous system and cholinergic neurons in the central nervous system. Its expression typically correlates with the amount of sympathetic innervation.

NGF is also expressed in the ovary (Lara *et al.*, 1990a) and is affected by ovarian innervation (Lara *et al.*, 1990a,b). NGF antibodies inhibit ovarian sympathetic innervation (Lara *et al.*, 1990c). The low-affinity NGF receptor is expressed in the ovary and is regulated upon ovulation (Dissen *et al.*, 1991). NGF production in the ovary is therefore likely to have actions on ovarian function through sympathetic innervation.

G. Additional Growth Factors

Several additional types of growth factors act and/or are produced by ovarian cells. One such factor is stem cell factor (SCF)/c-kit ligand. Zsebo *et al.* (1990a) characterized stem cell factor and found that it influences stem cell growth and development. SCF acts at the c-kit tyrosine kinase receptor (Zsebo *et al.*, 1990b) and therefore is also referred to as the c-kit ligand. The c-kit tyrosine kinase receptor is expressed in the ovary by oocytes at various stages of development (Manova *et al.*, 1990; Horie *et al.*, 1991). The c-kit receptor expression appears to decline with the onset of meiotic maturation, suggesting a role for SCF in meiotic arrest (Horie *et al.*, 1991). Expression of SCF by follicular cells suggests a role for SCF to mediate cell-cell interaction with the oocyte. It is anticipated that many additional growth factors will be identified upon further investigation. For example, keratinocyte growth factor and hepatocyte growth factor are present in the ovary and appear to mediate theca-granulosa all interactions (unpublished observation).

III. Summary

It is apparent that a large number of growth factors are produced and act in the ovary. Most of the research to date has focused on how specific growth factors affect differentiated functions of gonadal cell types. Factors that promote cell growth generally have suppressive effects on differentiation and attenuate hormone responsiveness. Factors that inhibit growth generally enhance differentiation and increase hormone responsiveness. When considering the function and physiology of locally produced growth factors, a distinction needs to be made between growth and differentiation. A factor that promotes cell proliferation and the cell cycle will indirectly

reduce the differentiated state of the cell, while the inverse is true of a factor that arrests cell proliferation and inhibits the cell cycle. Although specific growth factors may have a role in regulating differentiated functions, the possibility exists that many of the actions observed may be the indirect result of effects on cell growth. Therefore, the physiological importance of growth factor regulation of differentiated function remains to be elucidated. The control of cell growth, however, is a major function of specific growth factors. The integrated actions of various factors such as TGF α and TGF β could provide an efficient mechanism for regulating the cell proliferation required in gonadal development. Further investigation of the developmental regulation of the expression, production, and action of individual growth factors will provide insight into the potential physiological roles for the various growth factors. Evidence obtained to date implies that growth factors will be critical regulatory agents involved in ovarian cell-cell communication.

The endocrine regulation of ovarian cell growth has been well documented *in vivo*. The actions of gonadotropins and reproductive steroids, however, are distinct from the pharmacology of most growth factors. In addition, these hormones often have negligible effects on cell proliferation *in vitro*. The possibility that the actions of reproductive hormones on gonadal cell growth are indirectly mediated through alterations in the expression of locally produced growth factors needs to be seriously considered. Current work suggests that hormones may regulate growth factor production. Further investigation of the hormonal regulation of the production and actions of growth factors will help elucidate the mechanisms involved in the endocrine regulation of gonadal development.

References

- Adashi, E. Y., Resnick, C. E., Svoboda, M. E., and Van Wyk, J. J. (1984). A novel role for somatomedin-C in the cytodifferentiation of ovarian granulosa cells. *Endocrinology* 115, 1227-1229.
- Adashi, E. Y., Resnick, C. E., D'Ercole, A. J., Svoboda, M. E., and Van Wyk, J. J. (1985a). Insulin-like growth factors as intraovarian regulators of granulosa cell growth and function. *Endocr. Rev.* 6, 400-420.
- Adashi, E. Y., Resnick, C. E., Svoboda, M. E., and Van Wyk, J. J. (1985b). Somatomedin-C enhances induction of luteinizing hormone receptors by follicle-stimulating hormone in cultured rat granulosa cells. *Endocrinology* 116, 2369-2375.
- Adashi, E. Y., Resnick, C. E., Svoboda, M. E., and Van Wyk, J. J. (1986). Somatomedin-C as an amplifier of follicle-stimulating hormone action: enhanced accumulation of adenosine 3',5'-monophosphate. *Endocrinology* 118, 149-155.
- Adashi, E. Y., Resnick, C. E., and Twardzik, D. R. (1987). Transforming growth factor- α attenuates the acquisition of aromatase activity by cultured rat granulosa cells. *J. Cell. Biochem.* 33, 1-13.
- Adashi, E. Y., Resnick, C. E., Croft, C. S., May, J. V., and Gospodarowicz, D. (1988a). Basic fibroblast growth factor as a regulator of ovarian granulosa cell differentiation: a novel non-mitogenic role. *Mol. Cell. Endocrinol.* 55, 7-14.

- Adashi, E. Y., Resnick, C. E., Hernandez, E. R., Svoboda, M. E., and Van Wyk, J. J. (1988b). Characterization and regulation of a specific cell membrane receptor for somatomedin-C/insulin-like growth factor I in cultured rat granulosa cells. *Endocrinology* 122, 194–201.
- Adashi, E. Y., Resnick, C. E., Hernandez, E. R., Svoboda, M. E., and Van Wyk, J. J. (1988c). In vivo regulation of granulosa cell somatomedin-C/insulin-like growth factor I receptors. *Endocrinology* 122, 1383–1389.
- Adashi, E. Y., Resnick, C. E., Hernandez, E. R., May, J. V., Knecht, M., Svoboda, M. E., and Van Wyk, J. J. (1988d). Insulin-like growth factor-1 as an amplifier of follicle-stimulating hormone action: studies on mechanism(s) and site(s) of action in cultured rat granulosa cells. *Endocrinology* 122, 1583–1591.
- Baird, A., and Hsueh, A. J. W. (1986). Fibroblast growth factor as an intraovarian hormone: differential regulation of steroidogenesis by an angiogenic factor. *Regul. Pept.* 16, 243–250.
- Baranao, J. L., and Hammond, J. M. (1984). Comparative effects of insulin and insulin-like growth factors on DNA synthesis and differentiation of porcine granulosa cells. *Biochem. Biophys. Res. Commun.* 124, 484–490.
- Bendell, J. J., and Dorrington, J. (1991). Estradiol-17 beta stimulates DNA syntheses in rat granulosa cells: action mediated by transforming growth factor-beta. *Endocrinology* 128, 2663–2665.
- Cara, J. F., Fan, J., Azzarello, J., and Rosenfield, R. L. (1990). Insulin-like-growth factor-I enhances luteinizing hormone binding to rat ovarian theca-interstitial cells. *J. Clin. Invest.* 86, 560–565.
- Carpenter, G. (1987). Receptors for epidermal growth factor and other polypeptide mitogens. *Annu. Rev. Biochem.* 56, 881–914.
- Carpenter, G., and Cohen, S. (1990). Epidermal growth factor. *J. Biol. Chem.* 265, 7709–7712.
- Carson, R. S., Zhang, Z., Hutchinson, L. A., Herington, A. C., and Findlay, J. K. (1989). Growth factors in ovarian function. *J. Reprod. Fertil.* 85, 735–746.
- Caubo, B., DeVinna, R. S., and Tonetta, S. A. (1989). Regulation of steroidogenesis in cultured porcine theca cells. *Endocrinology* 125, 321–326.
- Chabot, J. G., St.-Arnaud, R., Walker, P., and Pelletier, G. (1986). Distribution of epidermal growth factor receptors in the rat ovary. *Mol. Cell. Endocrinol.* 44, 99–108.
- Chegini, N., and Flanders, K. C. (1992). Presence of transforming growth factor-beta and their selective cellular localization in human ovarian tissue of various reproductive stages. *Endocrinology* 130, 1707–1715.
- Chegini, N., and Williams, R. S. (1992). Immunocytochemical localization of transforming growth factors (TGFs) TGF-alpha and TGF-beta in human ovarian tissues. *J. Clin. Endocrinol. Metab.* 74, 973–980.
- Conn, G., Bayne, M. L., Soderman, D. D., Kwok, P. W., Sullivan, K. A., Palisi, T. M., Hope, D. A., and Thomas, K. S. (1990). Amino acid and cDNA sequences of a vascular endothelial cell mitogen that is homologous to platelet-derived growth factor. *Proc. Natl. Acad. Sci. USA* 87, 2628–2632.
- Daughaday, W. H., and Rotwein, P. (1989). Insulin-like growth factors I and II: peptide, messenger RNA and gene structure, serum, and tissue concentrations. *Endocr. Rev.* 10, 68–91.
- Davoren, J. B., and Hsueh, A. J. W. (1986). Growth hormone increases ovarian levels of immunoreactive somatomedin C/insulin-like growth factor I in vivo. *Endocrinology* 119, 2155–2162.
- Davoren, J. B., Kasson, B. G., Li, C. H., and Hsueh, A. J. W. (1986). Specific insulin-like growth factor (IGF) I- and II-binding sites on rat granulosa cells: relation to IGF action. *Endocrinology* 119, 2155–2162.
- Derynck, R. (1988). Transforming growth factor alpha. *Cell* 54, 593–595.
- Dissen, G. A., Hill, D. F., Costa, M. E., Ma, Y. J., and Ojeda, S. R. (1991). Nerve growth factor receptors in the peripubertal rat ovary. *Mol. Endocrinol.* 5, 1642–1650.

- Dodson, W. C., and Schomberg, D. W. (1987). The effect of transforming growth factor-beta on follicle-stimulating hormone-induced differentiation of cultured rat granulosa. *Endocrinology* 120, 512-516.
- Dorrington, J. H., Chuma, A. V., and Bendall, J. J. (1988). Transforming growth factor- β and follicle-stimulating hormone promote rat granulosa cell proliferation. *Endocrinology* 123, 352-359.
- Erickson, G. F., and Case, E. (1983). Epidermal growth factor antagonizes ovarian theca-interstitial cytodifferentiation. *Mol. Cell. Endocrinol.* 31, 71-76.
- Erickson, G. F., Nakatani, A., Ling, N., and Shimasaki, S. (1992a). Localization of insulin-like growth factor-binding protein-5 messenger ribonucleic acid in rat ovaries during the estrous cycle. *Endocrinology* 130, 1867-1878.
- Erickson, G. F., Nakatani, A., Ling, N., and Shimasaki, S. (1992b). Cyclic changes in insulin-like growth factor-binding protein-4 messenger ribonucleic acid in the rat ovary. *Endocrinology* 130, 625-636.
- Feng, P., Catt, K. J., and Knecht, M. (1986). Transforming growth factor beta regulates the inhibitory actions of epidermal growth factor during granulosa cell differentiation. *J. Biol. Chem.* 261, 14167-14170.
- Feng, P., Catt, K. J., and Knecht, M. (1988). Transforming growth factor- β stimulated meiotic maturation of the rat oocyte. *Endocrinology* 122, 181-188.
- Froesch, E. R., Schmid, C., Schwander, J., and Zapf, J. (1985). Actions of insulin-like growth factors. *Annu. Rev. Physiol.* 47, 443-467.
- Gangrade, B. K., and May, J. V. (1990). The production of transforming growth factor-beta in the porcine ovary and its secretion *in vitro*. *Endocrinology* 127, 2372-2380.
- Gangrade, B. K., Davis, J. S., and May, J. V. (1991). A novel mechanism for the induction of aromatase in ovarian cells *in vitro*: role of transforming growth factor alpha-induced protein tyrosine kinase. *Endocrinology* 129, 2790-2792.
- Geisthovel, F., Moretti-Rojas, I., Rojas, F. J., and Asch, R. H. (1990). Insulin-like growth factors and thecal-granulosa-cell function. *Hum. Reprod.* 5, 785-799.
- Goldenberg, R. L., Vaitukaitis, J. L., and Ross, G. T. (1972). Estrogen and follicle-stimulating hormone interactions on follicle growth in rats. *Endocrinology* 90, 1492-1497.
- Gonzales, A. M., Buscaglia, M., Ong, M., and Baird, A. (1990). Distribution of basic fibroblast growth factor in the 18-day rat fetus: localization in the basement membranes of diverse tissues. *J. Cell. Biol.* 110, 753-765.
- Gospodarowicz, D., III, and Birdwell, C. R. (1977). Effects of fibroblast and epidermal growth factors on ovarian cell proliferation *in vitro*. 1. Characterization of the response of granulosa cells to FGF and EGF. *Endocrinology* 100, 1108-1120.
- Gospodarowicz, D., and Bialecki, H. (1979). Fibroblast and epidermal growth factors are mitogenic agents for cultured granulosa cells of rodent, porcine, and human origin. *Endocrinology* 104, 757-764.
- Gospodarowicz, D., and Ferrara, N. (1989). Fibroblast growth factor and the control of pituitary and gonad development and function. *J. Steroid Biochem.* 32, 183-191.
- Gospodarowicz, D., Cheng, J., Lui, G.-M., and Bohlen, P. (1985). Corpus luteum angiogenic factor is related to fibroblast growth factor. *Endocrinology* 117, 201-213.
- Gospodarowicz, D., Ferrara, N., Schweigerer, L., and Neufeld, G. (1987). Structural characterization and biological functions of fibroblast growth factor. *Endocr. Rev.* 8, 95-114.
- Guthridge, M., Schmitt, J., Bertolini, J., Cowling, J., Runting, A., Katsahambas, S., Drummond, A. E., and Hearn, M. T. (1992). Studies on basic fibroblast growth factor (FGF-beta) gene expression in the rat and pig ovary using *in situ* hybridization and quantitative reverse transcriptase-polymerase chain reaction techniques. *Exs. Angiogenesis* 61, 219-229.
- Hammond, J. M., Baranoa, J. L. S., Skaleris, D., Knight, A. B., Ronanus, J. A., and Rechler, M. M. (1985). Production of insulin like growth factors by ovarian granulosa cells. *Endocrinology* 117, 2553-2555.

- Hernandez, E. R., Resnick, C. E., Svoboda, M. E., Van Wyk, J. J., Payne, D. W., and Adashi, E. Y. (1988). Somatomedin-C/insulin-like growth factor I as an enhancer of androgen biosynthesis by cultured rat ovarian cells. *Endocrinology* 122, 1603–1612.
- Hernandez, E. R., Roberts, C. T. Jr., LeRoith, D., and Adashi, E. Y. (1989). Rat ovarian insulin-like growth factor I (IGF-I) gene expression is granulosa cell-selective: 5'-untranslated mRNA variant representation and hormonal regulation. *Endocrinology* 125, 572–574.
- Hernandez, E. R., Roberts, C. T. Jr., Hurwitz, A., LeRoith, D., and Adashi, E. Y. (1990a). Rat ovarian insulin-like growth factor II gene expression is theca-interstitial cell-exclusive: Hormonal regulation and receptor distribution. *Endocrinology* 127, 3249–3251.
- Hernandez, E. R., Hurwitz, A., Payne, D. W., Dharmarajan, A. M., Purchio, A. F., and Adashi, E. Y. (1990b). Transforming growth factor- β 1 inhibits ovarian androgen production: gene expression, cellular localization, mechanism(s), and site(s) of action. *Endocrinology* 127, 2804–2811.
- Hernandez, E. R., Hurwitz, A., Botero, L., Ricciarelli, E., Werner, H., Roberts, C. T. Jr., LeRoith, D., and Adashi, E. Y. (1991). Insulin-like growth factor receptor gene expression in the rat ovary: divergent regulation of distinct receptor species. *Mol. Endocrinol.* 5, 1799–1805.
- Hernandez, E. R., Hurwitz, A., Vera, A., Pellicer, A., Adashi, E. Y., LeRoith, D., and Roberts, C. R. Jr. (1992). Expression of the genes encoding the insulin-like growth factors and their receptors in the human ovary. *J. Clin. Endocrinol. Metab.* 74, 419–425.
- Horie, K., Takakura, K., Taii, S., Narimoto, K., Noda, Y., Nishikawa, S., Nakayama, H., Fujita, J., and Mori, T. (1991). The expression of c-kit protein during oogenesis and early embryonic development. *Biol. Reprod.* 45, 547–552.
- Hsu, C., and Hammond, J. M. (1987). Gonadotropins and estradiol stimulate immunoreactive insulin-like growth factor-I production by porcine cells *in vitro*. *Endocrinology* 120, 198–207.
- Hsueh, A. J. W., Welsh, T. H., and Jones, P. B. C. (1981). Inhibition of ovarian and testicular steroidogenesis by epidermal growth factor. *Endocrinology* 108, 2002–2004.
- Hsueh, A. J. W., Adashi, E. Y., Jones, P. B. C., and Welsh, T. H. Jr. (1984). Hormonal regulation of the differentiation of cultured ovarian granulosa cells. *Endocr. Rev.* 5, 76–127.
- Hutchinson, L. A., Findlay, J. K., de Vos, F. L., and Robertson, D. M. (1987). Effects of bovine inhibin, transforming growth factor-beta and bovine Activin-A on granulosa cell differentiation. *Biochem. Biophys. Res. Commun.* 146, 1405–1412.
- Kim, I.-C., and Schomberg, D. W. (1989). The production of transforming growth factor- β activity by the rat granulosa cells. *Endocrinology* 124, 1345–1351.
- Knecht, M., and Catt, K. (1983a). Epidermal growth factor and gonadotropin-releasing hormone inhibit cyclic AMP-dependent luteinizing hormone receptor formation in ovarian granulosa cells. *J. Cell Biol.* 21, 209–217.
- Knecht, M., and Catt, K. J. (1983b). Modulation of cAMP-mediated differentiation in ovarian granulosa cells by epidermal growth factor and platelet-derived growth factor. *J. Biol. Chem.* 258, 2789–2794.
- Knecht, M., Feng, P., and Catt, K. J. (1986). Transforming growth factor-beta regulates the expression of luteinizing hormone receptors in ovarian granulosa cells. *Biochem. Biophys. Res. Commun.* 139, 800–807.
- Koos, R. D., and Olson, C. E. (1989). Expression of basic fibroblast growth factor in the rat ovary: detection of mRNA using reverse transcription-polymerase chain reaction amplification. *Mol. Endocrinol.* 3, 2041–2048.
- Kudlow, J. E., Kobrin, M. S., Purchio, A. F., Twardzik, D. R., Hernandez, E. R., Asa, S. L., and Adashi, E. Y. (1987). Ovarian transforming growth factor-a gene expression: immunohistochemical localization to theca-interstitial cells. *Endocrinology* 121, 1577–1579.
- LaPolt, P. S., Yamoto, M., Veljkovic, M., Sincich, C., Tor, N. Y., Tsafiriri, A., and Hsueh,

- A. J. W. (1990). Basic fibroblast growth factor induction of granulosa cell tissue-type plasminogen activator expression and oocyte maturation: potential role as a paracrine ovarian hormone. *Endocrinology* 127, 2357-2363.
- Lara, H. E., Hill, D. F., Katz, K. H., and Ojeda, S. R. (1990a). The gene encoding nerve growth factor is expressed in the immature rat ovary: effect of denervation and hormonal treatment. *Endocrinology* 126, 357-363.
- Lara, H. E., McDonald, J. K., Ahmed, C. E., and Ojeda, S. R. (1990b). Guanethidine-mediated destruction of ovarian sympathetic nerves disrupts ovarian development and function in rats. *Endocrinology* 127, 2199-2209.
- Lara, H. E., McDonald, J. K., and Ojeda, S. R. (1990c). Involvement of nerve growth factor in female sexual development. *Endocrinology* 126, 364-375.
- Leung, D. W., Cachianes, G., Kuang, W. J., Goeddel, D. V., and Ferrara, N. (1989). Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 246, 1306-1309.
- Lobb, D. K., Kobrin, M. S., Kudlow, J. E., and Dorrington, J. H. (1989). Transforming growth factor-alpha in the adult bovine ovary: identification in growing ovarian follicles. *Biol. Reprod.* 40, 1087-1093.
- Louvet, J.-P., and Vaitukaitis, J. L. (1976). Induction of follicle-stimulating hormone (FSH) receptors in rat ovaries by estrogen priming. *Endocrinology* 99, 758-764.
- Magoffin, D. A., Gancedo, B., and Erickson, G. F. (1989). Transforming growth factor- β promotes differentiation of ovarian thecal-interstitial cells but inhibits androgen production. *Endocrinology* 125, 1951-1958.
- Manova, K., Nocka, K., Besmer, P., and Bachvarova, R. F. (1990). Gonadal expression of c-kit encoded at the W locus of the mouse. *Development* 110, 1057-1069.
- May, J. V., Gilliam, F. R., Rein, M. S., Mondschein, J. S., and Schomberg, D. W. (1982). Growth and differentiated functions of porcine and rat granulosa cells following cryopreservation. *Biol. Reprod.* 27, 641-651.
- May, J. V., Buck, P. A., and Schomberg, D. W. (1987). Epidermal growth factor enhances [¹²⁵I]iodo-follicle-stimulating hormone binding by cultured porcine granulosa cells. *Endocrinology* 120, 2413-2420.
- Mondschein, J. S., and Schomberg, D. W. (1981). Growth factors modulate gonadotropin receptor induction in granulosa cell cultures. *Science* 211, 1179-1180.
- Mondschein, J. S., and Schomberg, D. W. (1984). Effects of partially and highly purified platelet-derived growth factor preparations in luteinizing hormone receptor induction in granulosa cell cultures. *Biol. Reprod.* 30, 603-608.
- Mondschein, J. S., Canning, S. F., and Hammond, J. M. (1988). Effects of transforming growth factor- β on the production of immunoreactive insulin-like growth factor I and progesterone and on [³H]thymidine incorporation in porcine granulosa cell cultures. *Endocrinology* 123, 1970-1976.
- Mondschein, J. S., Smith, S. A., and Hammond, J. M. (1990). Production of insulin-like growth factor binding proteins (IGFBPs) by porcine granulosa cells: identification of IGFBP-2 and -3 and regulation by hormones and growth factors. *Endocrinology* 127, 2298-2306.
- Mulheron, G. W., and Schomberg, D. W. (1990). Rat granulosa cells express transforming growth factor- β type 2 messenger ribonucleic acid which is regulatable by follicle-stimulating hormone *in vitro*. *Endocrinology* 126: 1777-1779.
- Mulheron, G. W., Danielpour, D., and Schomberg, D. W. (1991). Rat thecal/interstitial cells express transforming growth factor- β type 1 and 2, but only type 2 is regulated by gonadotropin *in vitro*. *Endocrinology* 129, 368-374.
- Mulheron, G. W., Bossert, N. L., Lapp, J. A., Walmer, D. K., and Schomberg, D. W. (1992). Human granulosa-luteal and cumulus cells express transforming growth factors-beta type 1 and 2 mRNA. *J. Clin. Endocrinol. Metab.* 74, 458-460.
- Nagy, F. (1972). Cell division kinetics and DNA synthesis in the immature Sertoli cells of the rat testis. *J. Reprod. Fertil.* 28, 389-395.

- Nakatani, A., Shimasaki, S., Erickson, G. F., and Ling, N. (1991). Tissue-specific expression of four insulin-like growth factor-binding proteins (1,2,3, and 4) in the rat ovary. *Endocrinology* 129, 1521-1529.
- Neufeld, G., Ferrara, N., Mitchell, R., Schweigerer, L., and Gospodarowicz, D. (1987). Granulosa cells produce basic fibroblast growth factor. *Endocrinology* 121, 597-603.
- Oliver, J. E., Aitman, T. J., Powell, J. R., Wilson, C. A., and Clayton, R. N. (1989). Insulin-like growth factor I gene expression in the rat ovary is confined to the granulosa cells of developing follicles. *Endocrinology* 124, 2671-2679.
- Phillips, H. S., Hains, J., Leung, D. W., and Ferrara, N. (1990). Vascular endothelial growth factor is expressed in rat corpus luteum. *Endocrinology* 127, 965-967.
- Ravindranath, N., Little-Ihrig, L., Phillips, H. S., Ferrara, N., and Zeleznik, A. J. (1992). Vascular endothelial growth factor messenger ribonucleic acid expression in the primate ovary. *Endocrinology* 131, 254-260.
- Ricciarelli, E., Hernandez, E. R., Hurwitz, A., Kokia, E., Rosenfeld, R. G., Schwander, J., and Adashi, E. Y. (1991). The ovarian expression of the antigonadotropic insulin-like growth factor binding protein-2 is theca-interstitial cell-selective: evidence for hormonal regulation. *Endocrinology* 129, 2266-2268.
- Ricciarelli, E., Hernandez, E. R., Tedeschi, C., Botero, L. F., Kokia, E., Rohan, R. M., Rosenfeld, R. G., Albiston, A. L., Herington, A. C., and Adashi, E. Y. (1992). Rat ovarian insulin-like growth factor binding protein-3: a growth hormone-dependent theca-interstitial cell-derived antigonadotropin. *Endocrinology* 130, 3092-3094.
- Richards, J. S. (1979). Hormonal control of ovarian follicular development: a 1978 perspective. *Rec. Prog. Horm. Res.* 35, 343-350.
- Roberts, A. B., and Sporn, M. B. (1988). Transforming growth factor beta. *Adv. Cancer Res.* 51, 107-145.
- Roberts, A. J., and Skinner, M. K. (1991). Transforming growth factors- α and - β differentially regulate growth and steroidogenesis of bovine thecal cells during antral follicle development. *Endocrinology* 129, 2041-2048.
- Roy, S. K., Ogren, C., Roy, C., and Lu, B. (1992). Cell-type-specific localization of transforming growth factor-beta 2 and transforming growth factor-beta 1 in the hamster ovary: differential regulation by follicle-stimulating hormone and luteinizing hormone. *Biol. Reprod.* 46, 595-606.
- Samares, S. E., Hagen, D. R., Shimasaki, S., Ling, N., and Hammond, J. M. (1992). Expression of insulin-like growth factor-binding protein-2 and -3 messenger ribonucleic acid in the porcine ovary: localization and physiological changes. *Endocrinology* 130, 2739-2744.
- Savion, N., Lui, G., Laherty, R., and Gospodarowicz, D. (1981). Factors controlling proliferation and progesterone production by bovine granulosa cells in serum-free medium. *Endocrinology* 109, 409-420.
- Schams, D., Koll, R., and Li, C. H. (1988). Insulin-like growth factor-I stimulates oxytocin and progesterone production by bovine granulosa cells in culture. *J. Endocrinol.* 116, 97-100.
- Skinner, M. K. (1990). Mesenchymal (stromal)-epithelial cell interactions in the testis and ovary which regulate gonadal function. *Reprod. Fertil. Dev.* 2, 237-243.
- Skinner, M. K., and Coffey, R. J. (1988). Regulation of ovarian cell growth through the local production of transforming growth factor- α by theca cells. *Endocrinology* 123, 2632-2638.
- Skinner, M. K., Keski, Oja, J., Osteen, K. G., and Moses, H. L. (1987a). Ovarian thecal cells produce transforming growth factor- β which can regulate granulosa cell growth. *Endocrinology* 121, 786-792.
- Skinner, M. K., Lobb, D., and Dorrington, J. H. (1987b). Ovarian thecal/interstitial cells produce an epidermal growth factor-like substance. *Endocrinology* 121, 1892-1899.
- Smith, E. P., Svoboda, M. E., VanWyk, J. J., Kierszenbaum, A. L., and Tres, L. T. (1987).

- Partial characterization of a somatomedin-like peptide from the medium of cultured rat Sertoli cells. *Endocrinology* 120, 186-193.
- Steinkampf, M. P., Mendelson, C. R., and Simpson, E. R. (1988). Effects of epidermal growth factor and insulin-like growth factor I on the levels of mRNA encoding aromatase cytochrome P-450 of human ovarian granulosa cells. *Mol. Cell. Endocrinol.* 59, 93-99.
- Thompson, N. L., Flanders, K. C., Smith, J. M., Ellingsworth, L. R., Roberts, A. B., and Sporn, M. B. (1989). Expression of transforming growth factor- β 1 in specific cells and tissues of adult and neonatal mice. *J. Cell Biol.* 108, 661-669.
- Tilly, J. L., and Johnson, A. L. (1990). Effect of several growth factors on plasminogen activator activity in granulosa and theca cells of the domestic hen. *Poultry Sci.* 69, 292-299.
- Tsafirri, A., Vale, W., and Hsueh, A. J. W. (1989). Effects of transforming growth factors and inhibin-related proteins on rat preovulatory Graafian follicles *in vitro*. *Endocrinology* 125, 1857-1862.
- Ui, M., Shimonaka, M., Shimasaki, S., and Ling, N. (1989). An insulin-like growth factor-binding protein in ovarian follicular fluid blocks follicle-stimulating hormone-stimulated steroid production by ovarian granulosa cells. *Endocrinology* 125, 912-916.
- Veldhuis, J. D., Rodgers, R. J., Dee, A., and Simpson, E. R. (1986). The insulin-like growth factor, somatomedin C, induces the synthesis of cholesterol side-chain cleavage cytochrome P-450 and adrenodoxin in ovarian cells. *J. Biol. Chem.* 261, 2499-2502.
- Veldhuis, J. D., Nestler, J. E., and Strauss, J. F. (1987). The insulin-like growth factor, somatomedin-C, modulates low density lipoprotein metabolism by swine granulosa cells. *Endocrinology* 121, 340-346.
- Yanker, B. A., and Shooter, E. M. (1982). The biology and mechanism of action of nerve growth factor. *Annu. Rev. Biochem.* 51, 845-868.
- Ying, S. T., Becker, A., Ling, N., Ueno, N., and Guillemin, R. (1986). Inhibin and beta type transforming growth factor (TGF β) have opposite modulating effects on the follicle stimulating hormone (FSH)-induced aromatase activity of cultured rat granulosa cells. *Biochem. Biophys. Res. Commun.* 136, 969-975.
- Zhiwen, Z., Findlay, J. K., Carson, R. S., Herington, A. C., and Burger, H. G. (1988). Transforming growth factor β enhances basal and FSH-stimulated inhibin production by rat granulosa cells *in vitro*. *Mol. Cell. Endocrinol.* 58, 161-166.
- Zsebo, K. M., Wypych, J., McNiece, I. K., Lu, H. S., Smith, K. A., Karkare, S. B., Sachdev, R. K., Yuschenkoff, V. N., Birkett, N. C., Williams, L. R., Satyagal, V. N., Tung, W., Bosselman, R. A., Mendiaz, E. A., and Langley, K. E. (1990a). Identification, purification, and biological characterization of hematopoietic stem cell factor from buffalo rat liver conditioned medium. *Cell* 6, 195-201.
- Zsebo, K. M., Williams, D. A., Geissler, E. N., Broudy, V. C., Martin, F. H., Atkins, H. L., Hsu, R., Birkett, N. C., Okino, K. H., Murdock, D. C., Jacobsen, F. W., Langley, K. E., Smith, K. A., Takeishi, T., Cattanach, B. M., Galli, S. J., and Snuggs, S. V. (1990b). Stem cell factor is encoded at the *Sl* locus of the mouse and is the ligand for the *c-kit* tyrosine kinase receptor. *Cell* 63, 213-224.