

# Developmentally distinct *in vivo* effects of FSH on proliferation and apoptosis during testis maturation

Sarah J Meachem<sup>1</sup>, Saleela M Ruwanpura<sup>1</sup>, Jessica Ziolkowski<sup>2</sup>, Jacquelyn M Ague<sup>3</sup>, Michael K Skinner<sup>3</sup> and Kate L Loveland<sup>2,4</sup>

<sup>1</sup>Prince Henry's Institute of Medical Research, Monash Medical Centre, 246 Clayton Road, Clayton, Victoria, 3168, Australia

<sup>2</sup>Monash Institute for Medical Research, Monash University, Clayton, Victoria, 3168, Australia

<sup>3</sup>Center for Reproductive Biology, School of Molecular Biosciences, Washington State University, Pullman, Washington, 99164, USA

<sup>4</sup>The Australian Research Council Centre of Excellence in Biotechnology and Development

(Requests for offprints should be addressed to S J Meachem; Email: sarah.meachem@phimr.monash.edu.au)

## Abstract

The critical influence of follicle stimulating hormone (FSH) on male fertility relates both to its impact on Sertoli cell proliferation in perinatal life and to its influence on the synthesis of Sertoli cell-derived products essential for germ cell survival and function in the developing adult testis. The nature and timing of this shift of germ cells to their reliance on specific Sertoli cell-derived products are not defined. Based on existing data, it is apparent that the dominant function of FSH shifts between 9 and 18 day postpartum (dpp) during the first wave of spermatogenesis from driving Sertoli cell proliferation to support germ cells. To enable comprehensive analysis of the impact of acute *in vivo* FSH suppression on Sertoli and germ cell development, FSH was selectively suppressed in Sprague–Dawley rats by passive immunisation for 2 days and/or 4 days prior to testis collection at 3, 9 and 18 dpp. The 3 dpp samples displayed no measurable changes, while 4 days of FSH suppression decreased Sertoli cell proliferation and numbers in 9 dpp, but not 18 dpp, animals. In contrast,

germ cell numbers were unaffected at 9 dpp but decreased at 18 dpp following FSH suppression, with a corresponding increase in germ cell apoptosis measured at 18 dpp. Sixty transcripts were measured as changed at 18 dpp in response to 4 days of FSH suppression, as assessed using Affymetrix microarrays. Some of these are known as Sertoli cell-derived FSH-responsive genes (e.g. StAR, cathepsin L, insulin-like growth factor binding protein-3), while others encode proteins involved in cell cycle and survival regulation (e.g. cyclin D1, scavenger receptor class B 1). These data demonstrate that FSH differentially affects Sertoli and germ cells in an age-dependent manner *in vivo*, promoting Sertoli cell mitosis at day 9, and supporting germ cell viability at day 18. This model has enabled identification of candidate genes that contribute to the FSH-mediated pathway by which Sertoli cells support germ cells.

*Journal of Endocrinology* (2005) **186**, 429–446

## Introduction

Sperm production in the adult male requires establishment of the Sertoli cell population and initiation of the first spermatogenic wave, events that occur during foetal and early postnatal life. In the rat, Sertoli cells proliferate until around 15 days after birth, setting the full complement of Sertoli cells. Surrounded by Sertoli cells, the gonocytes multiply in the foetal testis and then enter a period of quiescence lasting a week. They resume mitosis at 3 days after birth and migrate to the basement membrane at the seminiferous cord perimeter. Once they have contacted the basement membrane they are called spermatogonia. Spermatogonia subsequently pass through three developmental phases involving mitosis, meiosis and spermiogenesis to form spermatozoa, with the germ cells completing

the first wave of spermatogenesis at 43 days after birth in the rat (de Rooij 1998).

The mechanisms by which the Sertoli and germ cell populations are regulated during the first spermatogenic wave are under investigation and appear to be influenced by a complex network of interacting signals. It is known that pituitary-derived follicle stimulating hormone (FSH) exerts its effects on the developing germ cells indirectly, as only the Sertoli cells bear FSH receptors. A key role of FSH in setting the size of the Sertoli cell population in early postnatal life has been attributed to its stimulatory effect on Sertoli cell division (Orth 1984, Orth *et al.* 1988, Boitani *et al.* 1995). The response of the Sertoli cell to FSH changes during the first wave of spermatogenesis in rodents, and this influences the proliferation of Sertoli cells (Boitani *et al.* 1995, Meehan *et al.* 2000,

Buzzard *et al.* 2003). Synthesis of several Sertoli cell products also changes during development, however whether some of these are regulated by FSH *in vivo* remains to be determined (Munsie *et al.* 1997, Yan *et al.* 2001, Fragale *et al.* 2001, Migrenne *et al.* 2003, Pellegrini *et al.* 2003). FSH-regulated apoptosis during the first spermatogenic wave has not been examined, however in adult rats, germ cells are lost through apoptosis (Meachem *et al.* 1999), rather than through reduced proliferation when FSH is manipulated (McLachlan *et al.* 1995). However the regulation of adult spermatogenesis cannot be assumed to be mediated in the same manner as is the first spermatogenic wave, as shifts in the expression patterns of key regulatory genes (i.e., Bcl-2 family members and stem cell factor (SCF) have been observed as Sertoli cells mature and eventually mature germ cell populations emerge (Munsie *et al.* 1997, Huang *et al.* 1992, Meehan *et al.* 2001).

A large number of FSH responsive genes have been identified, however in most cases, their specific contribution to Sertoli cell and germ cell viability and function is undefined. The interaction of SCF with its receptor, c-kit, is essential for germ cell progression and survival in first wave and adult spermatogenesis (Yoshinaga *et al.* 1991, Packer *et al.* 1995), and expression of SCF in Sertoli cells has been shown to be regulated by FSH *in vitro* and *in vivo* (Rossi *et al.* 1993, McLean *et al.* 2002). Expression of the transcription factor DMRT1 in Sertoli cells can be elevated by FSH *in vivo* (Chen & Heckert 2001) and a role for DMRT1 in early postnatal testicular development relating to the termination of Sertoli cell proliferation has been deduced from the phenotype of *DMRT*<sup>-/-</sup> mice (Raymond *et al.* 2000). Some genes encoding apoptotic regulators in the Bcl-2 family of proteins have also been shown to be responsive to FSH, including *Bcl-w* and *Bok* (Yan *et al.* 2000, Suominen *et al.* 2001). A recent study using microarray analyses has defined 100–300 known transcripts that are regulated by FSH in cultures of Sertoli cells from 20 days post partum (dpp) rats (McLean *et al.* 2002). It is now timely to identify the FSH regulated genes involved in testicular development and spermatogenesis using an *in vivo* approach.

To more precisely understand the dynamic mechanisms by which FSH acts in immature rats, we set out to identify the cells that respond to *in vivo* changes in FSH levels during the first wave of spermatogenesis and to examine the functional changes in these cells. Administration of an antibody to FSH was performed to selectively suppress FSH for two or four days by immunoneutralisation. This acute FSH suppression differentially affected Sertoli and germ cell numbers during development as assessed by the, unbiased, optical disector stereological technique, corresponding with selective changes in proliferation and apoptosis in these cell types. Rat genomic microarrays enabled identification of candidate genes regulated by FSH from 14 to 18 dpp. The value of this model for

interrogating the changing roles of FSH in the first wave of spermatogenesis is discussed.

## Material and Methods

### Animals

Male outbred Sprague–Dawley rat pups with mothers were obtained from the Monash University Central Animal House (Clayton, Australia). They were maintained at 20 °C in a fixed 12 h light:12 h darkness cycle with free access to food and water in accordance with the Australian Code of Practice for Care and Use of Animals for Scientific Purposes (1997, National Health and Medical Research Council, Australia). This study was approved by the Monash Medical Centre Animal Ethics Committee.

### Passive immunisation against FSH

The ability of the polyclonal ovine antisera raised against rat FSH (FSHAb) to neutralise rat FSH *in vitro* and *in vivo* has been previously described (Meachem *et al.* 1998). The level of neutralisation of serum FSH achieved in adult rats was greater than 90% (reaching the limit of assay detection). Rat pups were immunised with the FSHAb or with normal sheep immunoglobulin (ConAb) at a daily dose of 10 mg/kg, a 5-fold higher dose than that previously administered to adult rats. Each animal received s.c. injections 2 days prior to death at 3 days dpp, and 2 and 4 days prior to death at 9 and 18 dpp. Ten rats were injected for each data point.

### Tissue collection and preparation

One h prior to death, each rat received 5-bromodeoxyuridine (BrdU; 50 mg/kg, s.c. Sigma) to enable proliferation analysis. Rats were killed 24 h after the final injection of the antibody, at 3, 9 or 18 dpp, by decapitation. Trunk blood was collected and allowed to clot overnight at 4 °C prior to serum collection for hormone assays. Due to the small amounts of serum collected for day 3 and 9 dpp rats, these samples were pooled in groups of three and two animals, respectively. The testes were then excised and weighed. The right testis of each animal was snap frozen on dry ice and stored at –75 °C for RNA preparation, while left testis was immersion-fixed with Bouin's solution for less than 5 h and sliced into 2-mm thick slabs orthogonal to the long axis of the testis. The testes from 9 and 18 dpp rats were divided into two, or four slices, respectively, and half of each processed into hydroxyethylmethacrylate resin (Technovit 7100; Kulzer and Co. GmbH, Friedrichsdorf, Germany) according to the manufacturer's instructions, while the other half was used for routine embedding into paraffin. The whole left testis from 3 dpp rats was embedded in resin and paraffin in every second

animal. Thick (25 µm) resin sections were serially cut (Supercut Microtome, Reichert Jung 2050, Nossloch, Germany), stained with the Periodic acid-Schiff's reaction reagents and counterstained with Mayer's Haematoxylin for the determination of cell number. Thin (5µm) paraffin sections were placed on Superfrost Plus slides for analysis of proliferation and apoptosis.

#### *Cell number estimates using the optical disector*

The optical disector stereological method (Wreford 1995) was used to determine the total number of cell nuclei per testis. All measurements were performed using a 100 × objective on an Olympus BX-50 microscope (Olympus, Tokyo, Japan). A microcator (D 8225; Heideinhain, Traunreut, Germany) that monitored scanned depth was attached to the microscope stage. The images were captured by a JVC TK-C1381 video camera coupled to a Pentium PC computer using a Screen Machine II fast multimedia video adapter (FAST, Hamburg, Germany). The software package, CASTGRID V1.60 (Olympus, Denmark, Germany), was used to generate an unbiased counting frame superimposed on video image. Fields were selected by a systematic uniform random sampling scheme as previously described (McLachlan *et al.* 1994, Wreford 1995) with the use of a motorized stage (Multi-control 2000; ITK, Lahnau, Germany). The final screen magnification was 2708-fold.

Sertoli cells were identified by their irregularly shaped nuclei, which were often positioned towards to the basement membrane and contained multiple nucleoli. Gonocytes were relatively large circular to ovoid cells with large circular nuclei, centrally located within the epithelium. Type A and B spermatogonia/preleptotene spermatocytes, leptotene/zygotene spermatocytes and pachytene spermatocytes were identified according to the characteristics described (Russell *et al.* 1990). Type B spermatogonia and preleptotene spermatocytes were not readily distinguishable at these stages of development and were counted as one. At least 300 Sertoli cell nuclei in total per testis were counted using the unbiased counting frame (175–430 µm<sup>2</sup>) and at least 80 gonocytes, spermatogonia (type A, B spermatogonia and preleptotene spermatocytes) or spermatocytes (leptotene, zygotene and pachytene spermatocytes) were counted employing a larger frame (430–1923 µm<sup>2</sup>). The frame size was selected based on cell frequency at different time points; less abundant cell types were counted in a larger counting frame. No correction for shrinkage was required (Meachem *et al.* 1996). Slides were masked prior to each type of quantitation (cell number, proliferation and apoptosis) to facilitate unbiased counting.

#### *Proliferation analysis*

BrdU incorporation into testicular cells at 3, 9 and 18 dpp was detected by immunohistochemistry as previously

described (Schlatt *et al.* 1999), with minor modifications as described below (see Fig 1A). In brief, slides bearing paraffin-embedded tissue sections were deparaffinized, rehydrated, and subjected to antigen retrieval in citrate buffer (0.1 M, pH 6; 90–95 °C for 10 min then room temperature for 20 min). Sections were treated successively with trypsin (0.25%, 1.5 min) and 3% hydrogen peroxide in methanol (20 min) and washed in PBS. All subsequent procedures were performed at room temperature, washes were in PBS, and incubations were in a humid chamber. Sections were treated with first with Superblock (40 min; DAKO, Carpinteria, CA, USA) and then with the monoclonal antibody to BrdU (clone BU-33, 4µl/ml in PBS, 1 h; Sigma). Following washes, biotinylated rabbit anti-mouse IgG was added (1:300 in PBS, 30 min; Zymed, California, USA). Sections were washed and incubated first in ABC complex (40 min; Vectastain Elite, Vector Laboratories, Burlingame, CA) and then with Tyramide Signal Amplification Biotin reagent (30 min; PerkinElmer Life Sciences, Inc, Boston, MA, USA) according to the manufacturers' instructions. After washing, diaminobenzidine (DAB) was added to reveal sites of antibody binding with a dark brown reaction product (2–3 min; DAKO Liquid DAB Substrate Chromogen System), and sections were counterstained with Mayer's Haematoxylin (3 min; Sigma), blued in Scott's tap water (1 min), and finally dehydrated and mounted in Depex (BDH Laboratory Suppliers, Poole, Dorset, UK) under glass coverslips. Cell types with anti-BrDU nuclear staining were identified on the basis of their location within the cord and the size and shape of cell nuclei, as described above.


The percentage of BrdU labelled cells was assessed using an unbiased counting frame of 430 µm<sup>2</sup>. To determine the proliferation index for each cell type, the total number of BrdU-labelled cells was divided by the total number of labelled and unlabelled cells.

#### *Apoptosis analysis*

Tissue sections (5 µm) were deparaffinized and rehydrated prior to the detection of DNA fragmentation (see Fig 1B). Apoptotic cells were detected using the terminal deoxynucleotidyl transferase (TdT)-mediated dUTP-biotin nick-end labelling (TUNEL) method (Meachem *et al.* 1999). Apoptotic cells were visualised using the chromogen DAB and processed as described for BrDU detection. On control sections, the TdT enzyme was omitted. TUNEL-positive cell types were identified based on their location within the cord, their size, and the shape of the cell nucleus.

The percentage of cells with TUNEL labelling was assessed using an unbiased counting frame of 430 µm<sup>2</sup> and the apoptotic index calculated by dividing the number of TUNEL labelled cells by the total number of labelled and unlabelled cells.





### Serum androgen levels

Serum androgen levels were measured by RIA using iodinated histamine–testosterone in combination with an acidic buffer (pH 5.1) to disrupt binding between testosterone and binding proteins in unextracted serum samples (O'Donnell *et al.* 1994). Serum samples (5–25 µl) were assayed in duplicate across two assays. Within assay variation for both assays was 3%, and between assay variation was 9%. Assay sensitivity was 0.6 ng/ml.

### Serum inhibin levels

Immunoreactive inhibin was measured by heterologous RIA as described previously (Robertson *et al.* 1988). Results are expressed in terms of an in-house rat ovarian extract calibrated against human recombinant (hr)-inhibin. Iodinated hr-inhibin was used as tracer. The antiserum used was rabbit antiserum (#1989) which is directed towards the  $\alpha$ -subunit thereby measuring both inhibin A and B, and cross-reacts 288% with pro-C, the prosequence of the inhibin alpha subunit (Robertson *et al.* 1989). Goat anti-rabbit IgG (GAR#12; Monash Institute of Reproduction & Development, Monash University, Melbourne, Australia) was used as second antibody. The assay buffer used was 0.01 M PBS containing 0.5% BSA (Sigma). Mouse serum pools diluted in a dose-dependent manner and were parallel to the standard curve (data not shown). The samples were run in a single assay in 20 µl duplicates. The within-assay variation was 7.2%. Assay sensitivity was 0.17 ng/ml.

### Oligonucleotide microarray hybridisation

To identify genes regulated by FSH at the 18 day time point, total RNA was isolated from total testes using acid phenol extraction (Chomczynski & Sacchi 1987). RNA from two individual animals treated from 14 dpp–18 dpp with either FSHAb or ConAb (4 samples in total) was used. The gene expression profile of FSHAb- and ConAb-treated testis were individually determined using Affymetrix RG\_U34A rat chips (Affymetrix Inc., Santa Clara, CA, USA) as previously described (McLean *et al.* 2002). The RNA from the FSHAb- and the

**Figure 1** Representative photomicrographs of cross sections of the testis from rats receiving normal sheep immunoglobulin for up to 4 days. Panel A. BrdU incorporation in the 3 dpp rat testis detected by immunohistochemistry. White arrows: nucleus of unlabelled Sertoli cell (SC) and gonocyte (G). Black arrow: nucleus of a labelled Sertoli cell (SC). Panel B. BrdU incorporation in the 9 dpp rat testis. White arrows: nucleus of unlabelled Sertoli cell (SC) and type B spermatogonia (B). Black arrows: nucleus of labelled Sertoli cell (SC) and type B spermatogonia (B). Panel C. DNA fragmentation indicative of cellular apoptosis detected by TUNEL in 18 dpp rat testis. White arrows: nucleus of unlabelled Sertoli cell (SC), type B spermatogonia (B) and pachytene spermatocyte (PSC). Black arrows: nucleus of labelled pachytene spermatocytes (PSC).

ConAb-treated testis was extracted and purified before biotin labelling and hybridisation to the microarray gene chip. Two replicates of each sample were hybridised to separate chips giving a total of four chips in the analysis. Each gene set on the Affymetrix chip is composed of 16 pairs of 24-mer oligonucleotides, with 8799 genes on each RG\_U34A chip. Included in each set is one anti-sense strand specific for the gene and one anti-sense strand with single point mutations is used as a comparative negative control. The labelled RNA was visualized on a Hewlett-Packard Gene Array Scanner (Hewlett-Packard Co., Palo Alto, CA, USA). An initial two way comparison was performed using the Affymetrix Suite software. In both experiments the FSHAb- and ConAb-treated chips were compared against each other.

#### *Bioinformatics and microarray statistics*

The bioinformatics and statistical analysis performed is as previously described (Eisen *et al.* 1998, Chaudhary *et al.* 2005, Kezele *et al.* 2002). Microarray output was examined visually for excessive background noise and physical anomalies. The default Microarray Suite (MAS, Silicon Genetics, Redwood City, CA, USA) statistical values were used for all analyses. An absolute analysis using MAS was performed to assess the relative abundance of the transcripts based on signal and detection (present, absent, or marginal) for the 16 different oligonucleotides per gene and comparison for analysis. The absolute analysis from MAS was imported into GeneSpring 5.1 software (Silicon Genetics, Redwood City, CA, USA). The data were normalized within GeneSpring using the default/recommended normalisation methods. These include setting of signal values below 0.01 to 0.01, total chip normalisation to the 50th percentile, and normalisation of each gene to the median. These normalisations allowed for the comparison of data based on relative abundance in any sample set rather than compared with a specific control value. Transcripts expressed differentially at a statistically significant level were determined using a one-way ANOVA parametric test with variances not assumed equal, and a *P*-value cutoff of 0.05. This was applied to all samples and considered all transcripts represented on the arrays. Two independent samples for each treatment group were analyzed was performed and allowed a 2 × 2 factorial comparison in the experiment. Subsequently, expression restrictions were applied to identify the transcripts expressed in a significant manner. These restrictions were designed so that the remaining transcripts met the following requirements in addition to being expressed in a significant manner: 1) each transcript have a signal value of at least 100 in the average of both samples, from at least one of the treatments tested and 2) had an average fold change of 1.5 or greater in signal intensity between treatments. Transcripts that passed these restrictions were considered for further analysis. Previous studies have

shown that microarray data correlates well with real-time quantitative PCR and Northern analysis (Eisen *et al.* 1998, Chaudhary *et al.* 2005, Kezele *et al.* 2002, Sadate-Ngatchou *et al.* 2004). Therefore, microarray data does not need to be confirmed as previously suggested (Shima *et al.* 2004). However, two selected genes were used in a real-time quantitative PCR procedure as previously described (McChlery & Clarke 2003) to help confirm the microarray procedure. The microarray chip data can be accessed at [www.skinner.wsu.edu](http://www.skinner.wsu.edu).

#### *Reverse transcription (RT) and real-time PCR analysis*

Real time PCR was used to measure the relative levels of two candidate FSH-regulated genes. Total RNA collected from two individual day 18 rats treated with either ConAb or with FSH Ab for 4 days was treated to remove residual genomic DNA (Ambion DNA-free Treatment Kit; Ambion, Austin TX, USA). RNA (500 ng) was converted to cDNA in a final volume of 20 µl using Superscript II according to the manufacturer's protocol (Invitrogen). For each sample, the absence of contaminating genomic DNA in cDNA samples was confirmed using reactions in which the RT enzyme was omitted. Quantitative RT-PCR analysis was performed using the Roche LightCycler (Roche, Mannheim, Germany) and the FastStart DNA Master SYBR-green 1 system (Roche). Oligonucleotide primer sequences for IGFBP-3, Smad3, and beta-actin were obtained either from published sources or Frodo.wi.mit.edu/cgi-bin/primer3/primer3 WWW.cgi;version 3.0 (Table 1). For PCR analysis, sample cDNA was diluted 1:10- to 1:50-fold, and PCR reaction conditions, including Mg<sup>2+</sup> concentration, primer concentration, anneal time and extension time were optimized for each primer pair as summarized in Table 1. For all PCR analyses, standard curves were generated using dilutions of an immature rat testicular cDNA preparation of arbitrary units (i.e. 17 dpp). PCR of all standards and samples were performed using duplicate reactions for approximately 40–45 cycles, after which a melting curve analysis was performed to monitor PCR product purity (see Table 1). In initial experiments, PCR product identities were verified by agarose gel electrophoresis and DNA sequencing (data not shown).

#### *Statistics*

A two-sample *t*-test was used to determine differences between FSHAb and ConAb treated samples with the assumption that data were normally distributed for all histological and hormone data. If data did not show normal distribution, then a Mann-Whitney test was carried out using Sigmapstat for Windows version 2.0 (Jandel Corporation, CA, USA). Data are expressed as mean ± S.E.M., *n*=7 rats per group. Statistical analysis of microarray data is described in Bioinformatics and microarray statistics.

**Table 1** Primer-specific LightCycler conditions used for PCR amplification of rat IGF-BP3, Smad 3 and beta-actin

Name [Accession No.]	Primer sequence	Product size (bp)	Primer concentration (pmol)	Mg <sup>2+</sup> concentration (mM)	Annealing temperature (°C)	Extension time (Sec)	Data temperature (°C)*	PCR product Tm (°C)**
Rat IGF-BP3 [NM_012588]	sense 5'-AATGCTGGGAGTGTGGAAG-3' antisense 5'-GCCGTATCTACTGGCTCTGC-3'	145	30	3.5	58	18	90	90
Rat Smad3 [U66479]	sense 5'-AAGGGCCAGCAGAACGGG-3' antisense 5'-GGGATGGAAATGGCTGTAGTC-3'	425	35	3.0	57	19	92	92
Rat β-actin [NM_031144]	sense 5'-CCGTAAGACCTCTATGCCAACA-3' antisense 5'-GCTAGGAGCCAGGCCAGTAATC-3'	103	50	2.5	67	5	80	86

\*Temperature at which the fluorescence of the PCR product was quantified during LightCycler analysis; \*\*PCR products have characteristic melting points (Tm). The Tm for a DNA product is defined as the temperature at which half of the DNA helical structure is lost, and is determined by melting temperature analysis on the LightCycler.

## Results

### Testicular weights and serum hormone measurements

Testicular weights were not significantly different following 2 days of FSH suppression in 3 dpp rats compared with their corresponding controls (Table 2), however testicular weights were reduced in 9 and 18 dpp animals after 2 (to 84% of control) and 4 days (to 78% of control) of FSH suppression. FSH suppression did not affect serum androgen levels at any time point compared with controls (Table 2). Serum inhibin levels following 2 days of FSH suppression in 3, 9 and 18 dpp rats were significantly reduced to 74%, 71% and 69%, and after 4 days 73% and 53% in 9 and 18 dpp rats compared with control values, respectively (Table 2). This reduction in circulating inhibin levels is presumed to reflect the successful neutralisation of FSH bioactivity through administration of the antibody raised against FSH, as FSH normally stimulates production of inhibin from Sertoli cells.

### Cell populations decreased in FSH withdrawn rats

At 3 dpp there were no differences in gonocyte ( $0.098 \pm 0.010$  vs  $0.091 \pm 0.054$  million/testis) or Sertoli cell numbers ( $3.50 \pm 2.4$  vs  $3.50 \pm 2.3$  million/testis) following 2 days of FSHAb treatment compared with ConAb treatment (Fig 2A), correlating no change in testis weight. At 9 dpp, there was a significant reduction (to 63% of control;  $P < 0.05$ ) in the number of Sertoli cells following 4 days of FSH suppression ( $16.8 \pm 1.7$  vs  $23.0 \pm 1.7$  million/testis), while spermatogonial numbers were unchanged ( $0.9 \pm 0.08$  vs  $1.1 \pm 0.4$  million/testis) (Fig 2B). At 18 dpp no difference in the number of Sertoli cells were observed in response to 4 days of FSHAb treatment ( $40.2 \pm 3.20$  vs  $44.2 \pm 4.18$  million/testis). A reduction in spermatogonia to 75% ( $16.9 \pm 1.8$  vs  $22.2 \pm 1.0$  million/testis,  $P < 0.05$ ) of control group values was observed following 4 days of FSH suppression. Spermatocyte number tended to be reduced to 80% ( $12.4 \pm 1.1$  vs  $15.5 \pm 1.0$  million/testis,  $P = 0.061$ ) of control group values was observed following 4 days of FSH suppression, although this did not achieve significance (Fig 2C).

### FSH suppression decreases Sertoli cell proliferation in day 9 rats

At 3 dpp, no differences in Sertoli cell proliferation were observed between FSHAb- and ConAb-treated rats ( $23.4\% \pm 3.2$  vs  $27.2\% \pm 1.8$ ); no BrdU labelled gonocytes were observed in these samples (Refer Fig 1). At 9 dpp, there was a significant reduction (to 66% of control,  $P < 0.01$ ) in the proliferation rate of Sertoli cells following FSH suppression ( $14.3\% \pm 0.6$  vs  $21.6\% \pm 1.2$ ) (Fig 2A), while no change was observed for spermatogonia ( $23.2\% \pm 2.4$  vs  $21.1\% \pm 1.6$ ,  $P = 0.11$ ) (Fig 3A). At



**Table 2** The testicular weights (mg), serum testosterone (ng/ml) and inhibin (ng/ml) levels at 3 dpp, 9 dpp and 18 dpp following 2 and 4 days of FSH suppression. Rats were treated with normal sheep immunoglobulin (ConAb) and with antibody raised against rat FSH (FSHAb) at 3 dpp, 9 dpp and 18 dpp following 2 and 4 days of treatment. The data are expressed as mean  $\pm$  S.E.M., with  $n=10$  rats/group.

Treatment	Testicular weight (mg)	Serum testosterone (ng/ml)	Serum inhibin (ng/ml)
3 days post partum			
+ConAb 2d	5.1 $\pm$ 0.2	0.64 $\pm$ 0.08	3.57 $\pm$ 0.21
+FSHAb 2d	5.1 $\pm$ 0.1	0.63 $\pm$ 0.11	2.65 $\pm$ 0.06**
9 days post partum			
+ConAb 2d	17.9 $\pm$ 0.9	1.36 $\pm$ 0.3	4.53 $\pm$ 0.3
+FSHAb 2d	15.2 $\pm$ 0.5*	1.18 $\pm$ 0.1	3.23 $\pm$ 0.2*
+ConAb 4d	18.7 $\pm$ 0.4	0.89 $\pm$ 0.2	4.17 $\pm$ 0.24
+FSHAb 4d	14.6 $\pm$ 0.5**	0.70 $\pm$ 0.1	3.03 $\pm$ 0.19*
18 days post partum			
+ConAb 2d	75.4 $\pm$ 3.0	0.92 $\pm$ 0.3	3.51 $\pm$ 0.4
+FSHAb 2d	63.2 $\pm$ 3.4*	0.92 $\pm$ 0.2	2.43 $\pm$ 0.2*
+ConAb 4d	74.1 $\pm$ 1.2	1.80 $\pm$ 0.5	5.75 $\pm$ 0.31
+FSHAb 4d	57.9 $\pm$ 2.1**	1.36 $\pm$ 0.2	3.09 $\pm$ 0.20**

Asterisks denote significant differences between rats receiving FSHAb and their corresponding controls. \* $P<0.01$  and \*\* $P<0.001$ .

18 dpp, in response to FSH suppression, there was no change in Sertoli cell (6.8%  $\pm$  2.3 vs 9.5%  $\pm$  1.0,  $P=0.30$ ), spermatogonial (19.6%  $\pm$  1.9 vs 23.3%  $\pm$  1.0,  $P=0.11$ ) or spermatocyte proliferation (11.9%  $\pm$  1.4 vs 10.8%  $\pm$  1.7,  $P=0.64$ ) (Fig 3B) compared with their corresponding controls. These data indicate that the reduction in testis weight due to FSH suppression at 9 dpp, but not at 18 dpp, reflects the lower number of Sertoli cells in these samples. The incidence of apoptosis was next investigated as an alternative basis for the reduced testis weights in the 18 dpp samples (Fig 3B).

#### *Germ cell apoptosis is affected by FSH suppression in 18 dpp rats*

No apoptosis was observed in Sertoli cells and gonocytes at 3 dpp (Refer Fig 1). No differences in Sertoli cell (0.6%  $\pm$  0.2 vs 0.9%  $\pm$  0.2,  $P=0.36$ ) or spermatogonial apoptosis (6.6%  $\pm$  1.1 vs 4.6%  $\pm$  0.9,  $P=0.20$ ) were observed following 4 days of FSH suppression at 9 dpp compared with controls (Fig 4A). However, at 18 dpp, there was a significant elevation of spermatogonial apoptosis (250% of control) in response to FSH suppression (15.5%  $\pm$  1.6 vs 6.2%  $\pm$  1.0,  $P<0.001$ ); spermatocyte apoptosis increased to 136% of control levels (16.1%  $\pm$  2.2 vs 11.8%  $\pm$  1.4,  $P=0.13$ ), although this did not achieve significance (Fig 4B). No change in Sertoli cell apoptosis was observed at any time point (Fig 4). These data indicate that a shift occurs between 9 and 18 dpp within the germ cell population towards reliance for support cues on FSH-derived products. The nature of this support system

and the effects of its removal were examined using a genomic microarray to identify genes regulated by altered levels of FSH at 18 dpp.


#### *Microarray analysis*

Thirty genes were identified in the 18 dpp samples as up-regulated and 30 as down-regulated (Table 3) following FSH suppression by treatment with FSHAb for 4 days. These 60 genes of interest met the selection criteria outlined previously; all give at least one signal intensity greater than 100, are flagged as changing, and have fold changes of 1.5 or greater.

Amongst these genes, 12 out of 60 (20%) have previously been reported to be regulated by FSH, predominantly through *in vitro* studies. Many of the genes on this list have not yet been identified as being expressed in the testis (26/60), and some of those known to be in the testis have not previously been assessed for regulation by FSH (14/34). This latter category of candidate *in vivo* FSH target genes have been reported to be expressed in Sertoli cells (4/14), germ cells (8/14) and/or non-Sertoli somatic cells (6/14); clearly the non-Sertoli cells expressed genes that would be regulated indirectly, through FSH regulation of Sertoli cell biosynthesis or other functions.

#### *RT-PCR analysis*

IGFBP-3 mRNA expression levels (measured in two individual animals) in FSHAb-treated rats were significantly elevated approximately 2.5-fold above the levels




**Figure 2** The number of Sertoli cells, spermatogonia and spermatocytes at 3 dpp (A), 9 dpp (B) and 18 dpp (C) following 2 and 4 days of FSH suppression. Cell numbers in rats treated with normal sheep immunoglobulin (black bars) and rats treated with antibody raised against rat FSH (white bars) at 3 dpp, 9 dpp and 18 dpp. Sertoli cell, spermatogonial and spermatocyte numbers are expressed as mean  $\pm$  S.E.M., where  $n=7$  rats/group. Asterisk identify significant differences between treated and control groups at  $P<0.05$ .

measured in two individual ConAb-treated rats. In contrast, Smad 3 mRNA expression was significantly reduced by approximately 6-fold following FSHAb treatment compared with ConAb-treated rats (Fig 5).

## Discussion

These data illustrate that the influence of FSH on Sertoli and germ cells are dynamic and cell-type specific during




**Figure 3** Proliferation of specific cell populations of the seminiferous epithelium at 9 dpp (A) and 18 dpp (B) following 4 days of FSH suppression. Percentage of cells incorporating BrdU in rats treated with normal sheep immunoglobulin (black bars) and rats treated with antibody raised against rat FSH (white bars) at 9 dpp and 18 dpp. The percentages of labelled versus unlabelled total cells, Sertoli cells, spermatogonia and spermatocytes are expressed as mean  $\pm$  S.E.M., where  $n=7$  rats/group. Asterisks denote significant differences between treated and control groups at  $P<0.01$ .

the first spermatogenic wave in the rat. We developed this model to enable a co-ordinated documentation of specific cellular changes, in association with the identification of testicular genes, for which expression is affected by variance in bioactive FSH levels. The results are in accord with previous observations that FSH and activin regulate Sertoli and germ cell proliferation in an age-dependent manner *in vitro* (Boitani *et al.* 1995, Fragale *et al.* 2001). Our findings expand on earlier analyses by measuring proliferation and apoptosis in the same samples and establishing a method for identification of FSH target genes *in vivo*. More specifically, this study demonstrated for the first time *in vivo* that acute FSH suppression reduces both the Sertoli and germ cell populations, first by inhibiting Sertoli cell proliferation from 5 to 9 dpp, and later by inducing germ cell apoptosis between 14 and 18 dpp (see Fig 6).

Acute suppression of FSH for 4 days resulted in decreased testis weights in 9 and 18 dpp rats. This decrease





**Figure 4** Apoptosis in specific cell populations of the seminiferous epithelium at 9 dpp (A) and 18 dpp (B) following 4 days of FSH suppression. The percentages of TUNEL labelled cells in rats treated with normal sheep immunoglobulin (black bars) and rats treated with antibody raised against rat FSH (white bars) at 9 dpp and 18 dpp. The percentages of labelled versus unlabelled total cells, Sertoli cells, spermatogonia and spermatocytes are expressed as mean  $\pm$  S.E.M., where  $n=7$  rats/group. Asterisks denote significant differences between treated and control groups at  $P<0.001$ .

in testicular weight was attributed, at least in part, to the decreased Sertoli cell population at 9 dpp and a decreased germ cell population in 18 dpp rats as assessed by the optical disector stereological technique. The Sertoli cell population decrease resulted from inhibition of proliferation and not from elevated apoptosis. In contrast, the reduction in spermatogonial and spermatocyte cell number at 18 dpp resulted from elevated apoptosis of germ cells.

#### Sertoli cell development

Our findings are in accord with other studies which show that FSH plays a central role in regulating the size of the Sertoli cell population (Meachem *et al.* 1996) by supporting Sertoli cell proliferation during early postnatal life (Orth 1984, Orth *et al.* 1988, Boitani *et al.* 1995). Whether FSH plays a role in terminating Sertoli cell proliferation around 15 days after birth is beyond the scope of this study, however data from other *in vitro* and *in vivo*

studies indicate a role for FSH in this process (Baker & O'Shaughnessy 2001, Buzzard *et al.* 2002). Survival of Sertoli cells in the postnatal rat does not appear to be influenced by FSH levels, as evidenced by the fact that FSH suppression did not induce Sertoli cell apoptosis, though the possibility remains that there are residual levels of serum FSH in this model which are sufficient to support Sertoli cell survival. We predict that a greater impact on testicular physiology could be apparent if complete FSH withdrawal were achieved.

Several other factors have been demonstrated to regulate Sertoli cell development in the time interval under investigation here, including thyroid hormone (Cooke & Meisami 1991, Van Haaster *et al.* 1992), activin (Boitani *et al.* 1995, Fragale *et al.* 2001, Buzzard *et al.* 2003), glial cell line-derived neurotrophic factor (GDNF) (Hu *et al.* 1999), interleukin-1 (Petersen *et al.* 2002), and transforming growth factor (TGF) and epidermal growth factor (Petersen *et al.* 2001). Interactions between these factors and FSH-mediated regulation of Sertoli cell function is not well understood, though two previously identified FSH regulated target genes, cyclin D2 (Buemer *et al.* 2000) and DMRT-1 (Raymond *et al.* 2000) are potential mediators.


In this study, the FSHAb dose administered was five times higher than that previously shown to neutralise greater than 90% of serum FSH in adult rats (Meachem *et al.* 1998, 1999). This dose was selected because serum FSH levels are 3–5 higher during postnatal life than in adulthood, as measured using RIA (Kirby *et al.* 1992). In our previous studies with this antibody, neutralisation of bioactive material was determined using an FSH *in vitro* bioassay, based on the induction of aromatase activity in immature rat Sertoli cells and assay of free FSH from rat serum using HPLC and RIA (Meachem *et al.* 1998, 1999). The significant reduction in serum inhibin levels following FSHAb administration in the present study are a positive indicator of the efficacy of FSH immunoneutralisation. Whether FSH regulates Sertoli cell development in the 3 dpp testis remains unclear. Our data suggest that Sertoli cell proliferation and apoptosis are not governed by FSH between 1 and 3 dpp in the rat. Consistent with this, Buzzard *et al.* (2003) reports that proliferation of purified Sertoli cells from rats younger than 6 dpp is not affected by FSH. On the other hand, Boitani *et al.* (1995) and Meehan *et al.* (2000) reported that Sertoli cell proliferation in 3 dpp rat testis fragment cultures was enhanced in the presence of FSH. Differences between our new data and that reported in earlier studies may reflect differences between *in vitro* and *in vivo* models, with the latter being uniquely employed in the present study. It is possible that removal of FSH for 2 days *in vivo* may be insufficient to observe a subtle but functionally important change in Sertoli cell proliferation at this age, due to this relatively small population of Sertoli cells present at this time (<4 million Sertoli cells/ 3 dpp testis). It may be necessary to extend the period of FSH suppression or to

**Table 3** Genes up and down regulated by FSH in day 18 postpartum rat testes. Day 15 rats were treated with normal sheep immunoglobulin or with an antibody raised against rat FSH for 4 days. Results are based on two microarray hybridizations of target synthesized from RNA from two independent rat testes from each treatment group. The Affymetrix microarray data set was analyzed by pairwise comparison using Gene Sifter. Sample sets with a signal at  $\geq 100$  detected in at least one position on the array and a fold change of 1.5 or greater are reported. References for those genes previously reported to be in the rat testis and/or regulated by FSH are in parentheses.

Genes up-regulated by FSH suppression	Accession number	Fold change	Average signal FSHAD	+/- s.e.m.	Testis expression previously reported	FSH regulation previously reported
Insulin-like growth factor binding protein 3	AI009405; M31837	6.47; 3.12	17.45	5.85; 1.99	Yes; Sertoli cells (Smith <i>et al.</i> 1990, Rappaport & Smith 1995, Khan <i>et al.</i> 2002)	Yes
Testin	U16858	3.97	30.65	0.90	Yes; Sertoli cells (Grima <i>et al.</i> 1998, Lui <i>et al.</i> 2003)	Yes
HNF-3/forkhead homolog-1	L13201	3.14	36.65	0.39	Yes; (Kaestner <i>et al.</i> 1998, Migrenne <i>et al.</i> 2003, Wolfrum <i>et al.</i> 2003)	No
2-4 kb repeat DNA right terminal region	X05472	2.62	53.0	1.99	No	No
Endothelin 1	M64711	2.46	36.15	0.46	Yes; Sertoli cells (Fantoni <i>et al.</i> 1993, Tripiciano <i>et al.</i> 1999)	Yes
cDNA clone rX01264 3'	AI639097	2.44	121.55	0.25	No	No
ATP-binding cassette, sub-family C sulfonylurea receptor 2B mRNA (SUR2B);	AF019628	2.33	20.75	1.01	Yes; Leydig cells (Schinkel & Jonker 2003, O'Shaughnessy <i>et al.</i> 2002)	No
Cyclin D1	AI231257	2.04	8.50	1.60	Yes; gonocytes & spermatogonia (Wolgemuth <i>et al.</i> 1998, Buemer <i>et al.</i> 2000)	No
cDNA, 3' end /clone=RKIBA73	AA892986	1.98	6.05	0.90	No	No
Lipase A, lysosomal acid	S81497	1.98; 1.84	55.95	0.63; 0.63	Yes; Guinea pig (Kabbaj <i>et al.</i> 2001)	No
SH3 domain binding protein CR16	U31159	1.93	10.8	0.63	No	No
G-protein-coupled receptor 105	U76206	1.81	8.85	0.56	No	No
Fibrinogen, beta polypeptide	M35602	1.80	7.85	0.52	No	No
Cyclic Protein-2=cathepsin L proenzyme	S85184	1.74; 1.67	58.15	0.12; 0.12	Yes; Sertoli cells (Penitilla <i>et al.</i> 1995, Wright <i>et al.</i> 2003)	Yes
Mast cell protease 7	U67910	1.74	20.6	0.85	No	No
UDP glycosyltransferase 1 family, polypeptide A7	AF039212	1.71	38.5	0.18	No	No
N-myc downstream-regulated gene 2	AA799560	1.70	185.55	0.10	No	No
Rat retinol-binding protein (RBP)	M10934	1.69	38.50	0.37	Yes; Sertoli cells (Eskild <i>et al.</i> 2000)	Yes
Crystallin, alpha B	M55534	1.67	43.90	0.84	Yes; spermatocytes (Kato <i>et al.</i> 1991)	No
Similar to microsomal glutathione S-transferase 3 (LOC289197)	AA892234	1.66	358.95	0.25	No	No
Ribonuclease 7 precursor; eosinophil cationic protein	D88586	1.63	30.7	0.27	No	No
Mitogen activated protein kinase kinase 1	U48596	1.58	321.3	0.55	No	No
Cadherin 2; N-Cadherin	AF097593	1.57	65.15	0.30	Yes; Sertoli cells (Lampa <i>et al.</i> 1999)	Yes
Rat nuclear receptor co-regulator 1	M81766	1.56	75.85	0.09	Yes; Sertoli cells (Dufour & Kim 1999, Kasiner <i>et al.</i> 1996)	No
Reelin	AA893471	1.55	39.25	0.44	Yes; (Ikeda & Terashima 1999, Andersen <i>et al.</i> 2003)	No
Vdulp1; upregulated by 1,25-dihydroxyvitamin D-3	AI014169	1.55	490.7	0.20	No	No
PC4; interferon-related developmental regulator 1	AI014163	1.54	78.05	0.37	Yes; germ cells (Mbikay <i>et al.</i> 1997)	No
N-acetylglucosaminyltransferase I	D16302	1.51	132.45	0.09	Yes; (Nozaki <i>et al.</i> 2003)	No
mRNA for MHC class II antigen RT1.B-1 beta-chain	M36151	1.51	50.90	2.25	Yes; germ cells (Mori <i>et al.</i> 2000)	No
<i>Rattus norvegicus</i> transcribed sequences	AA891962	1.51	11.6	0.16	No	No

Table 3 Continued

Accession number	Fold change	Average signal FSHAb	+/- S.E.M.	Testis expression previously reported	FSH regulation previously reported
<b>Genes up-regulated by FSH suppression</b>					
Steroidogenic acute regulatory protein	4.34; 4.21	59.85; 81.05	6.81; 1.36	Yes; Sertoli cells (Gregory & De Phillip 1998, McLean et al. 2002, Manna et al. 2003)	Yes
Aminolevulinic acid synthase 2	3.00	188.20	0.64	No	No
Phosphodiesterase 4B	2.55; 1.80	14.45	0.51; 0.42	Yes; Sertoli cells (Swinnen et al. 1991)	Yes
MAD homolog 3 ( <i>Drosophila</i> )	2.39	40.65	0.12	Yes; Sertoli, Leydig cells & spermatocytes (Xu et al. 2003)	No
Melanoma inhibitory activity; cartilage derived retinoic acid sensitive protein	2.27	357.45	1.12	No	No
Syndecan 1	2.24	82.10	0.48	Yes; Sertoli cells (Brucato et al. 2002)	Not tested
75 kDa glucose regulated protein	2.08	534.40	0.23	No	No
Gap junction membrane channel protein alpha 1 (gja-1); connexin43	2.06	50.10	0.38	Yes; Sertoli, Leydig & germ cells (Sommersberg et al. 2000, Bravo-Moreno et al. 2002)	Yes
Neuronal activity-regulated pentraxin	2.05	253.4	1.51	No	No
Protein C inhibitor	2.03	6.76.65	0.28	Yes; Leydig & germ cells (Uhrin et al. 2000, Odet et al. 2003)	No
Organic anion transporter protein 3 (OAT3); solute carrier family 21 (fatty acid transporter), member 7	1.92	27.30	0.45	Yes	No
Amyloid beta (A4) precursor-like protein 2	1.87	431.55	0.20	Yes; von der Kammer et al. 1994)	No
<i>R. norvegicus</i> cDNA clone rx05024, 3'	1.82	52.90	0.44	No	No
<i>R. norvegicus</i> similar to Esophagus cancer-related gene-2 protein precursor (ECRG-2) (LOC301016), mRNA	1.82	336.75	0.23	No	No
Adducin 3, gamma	1.79	127.8	0.00	No	No
Monocarboxylate transporter (MCT1); solute Carrier family 16, member 1	1.77	45.8	0.20	Yes; germ cells (Jackson et al. 1997, Boussouar et al. 2003, Goddard et al. 2003)	No
Scavenger receptor class B, member 1	1.77	345.00	0.35	Yes; Sertoli & Leydig cells (Reaven et al. 2000, Kawasaki et al. 2002)	No
Lysyl oxidase	1.76	44.3	0.16	Yes; (Asuncion et al. 2001, Maki et al. 2001)	No
Transcribed sequence with moderate similarity to protein ref: NP_061921.1 ( <i>Homo sapiens</i> ) hypothetical protein FLJ20752 ( <i>H. sapiens</i> )	1.67	174.65	0.42	Yes; (Ericsson et al. 1993)	No
Proteolipid protein	1.67	169.75	0.20	Yes; (Li et al. 2002)	No
IP63 protein	1.66	80.35	0.08	No	No
Fumarate hydratase 1	1.63	61.45	0.18	No, but it is ubiquitous	No
cAMP responsive element modulator	1.63	36.85	0.74	Yes; (Walker et al. 1994, West et al. 1994, Monaco et al. 1995)	Yes
Lamin B1	1.63	41.55	0.59	Yes; Sertoli, Leydig, peritubular & germ cells (Machiels et al. 1997)	No
Cyclin D2	1.60	279.55	0.45	Yes; Sertoli cells & spermatogonia (Wolgemuth et al. 1998, Buemer et al. 2000, Strothmann et al. 2004)	Yes
Androgen Binding Protein (ABP); sex hormone binding globulin	1.59	873.95	0.11	Yes; Sertoli cells (Skinner et al. 1989, Danzo 1995, Larriba et al. 1995, Joseph et al. 1997)	Yes
Superoxide dismutase 3	1.57	147.15	0.21	No	No
Chondroitin sulfate proteoglycan	1.56	17.85	0.34	Yes; Leydig cells (Ungefroren et al. 1995)	No
Nicotinamide nucleotide transhydrogenase	1.53	405.45	0.30	Yes; (Arklblad et al. 2002)	No
Transcribed sequence with moderate similarity to protein sp: P00722 ( <i>E. coli</i> ) BGAL_EC01 Beta-galactosidase (Lactase)	1.52	600.8	0.39	No	No



**Figure 5** Expression levels of rat Smad 3 and IGF-BP3 by real time PCR. Two rats treated with normal sheep immunoglobulin (black bars) and treated with antibody raised against rat FSH (white bars) were tested to validate data obtained from the microarray analysis. Each sample was measured over three separate experiments in terms of ng of mRNA normalised to beta actin. Data are mean  $\pm$  S.D.

increase the dose of neutralising antibody if in fact residual levels of FSH exist. The Sertoli cell numbers measured in control antibody-treated groups in this study are in accord with our previous study (Meachem *et al.* 1996) and that of others (Simorangkir *et al.* 1997), reporting numbers quantified by the optical disector stereological technique, in rats of similar ages.


#### Germ cell development

These data provide no evidence that FSH supports the maturation of gonocytes into type A spermatogonia. Studies of congenitally deficient hypogonadal mice (hpg) lacking both FSH and LH show that gonocyte number is

not altered in these newborn mice compared with wild-type controls, however gonocyte number is significantly reduced at 5 dpp, suggesting that gonocyte survival may be in part regulated by gonadotrophins. Whether this is an effect specific to FSH is yet to be elucidated (Baker & O'Shaughnessy 2001). Several other recent studies highlight the impact of locally produced members of the TGF $\beta$  superfamily on germ cell development, including bone morphogenetic protein 4 (BMP4) (Pellegrini *et al.* 2003), activin (Boitani *et al.* 1995, Meehan *et al.* 2000) and GDNF (Meng *et al.* 2000, Yomogida *et al.* 2003), and production of both inhibin (an activin antagonist) and GDNF is regulated by FSH (Tadokoro *et al.* 2002). Meehan *et al.* (2000) provides *in vitro* evidence, using 1 and 3 day cultures of 3 dpp rat testis fragments, that FSH plus follistatin mediates gonocyte maturation into type A spermatogonia, but that FSH alone does not support gonocytes, as assessed on a per Sertoli cell basis. Meehan's study reported that activin independently elevated gonocyte number 4-fold above that of DMEM treated fragments and that follistatin prevented this activin-induced rise, giving rise to the concept that balance between activin and follistatin can influence germ cell maturation at the onset of spermatogenesis. Another facet of first wave initiation, gonocyte migration, relies at least in part on the interaction of c-kit and its FSH-regulated ligand, SCF (Orth *et al.* 2000), and it is apparent that the population of migratory gonocytes have stem cell activity, while those lacking pseudopods are undergoing apoptosis (Orwig *et al.* 2002). Thus influence of FSH at this age may reflect a complex set of interacting signals.

Several lines of evidence indicate that FSH enables spermatogonial survival in the adult rat, (Shetty *et al.* 1996, Meachem *et al.* 1999) without directly enhancing germ cell proliferation *in vivo* (McLachlan *et al.* 1995). In immature rats rendered FSH- and LH-replete by hypophysectomy, administration of FSH reduced the number of degenerating cells (Russell *et al.* 1987) and prevented germ cell apoptosis (Tapanainen *et al.* 1993). Evidence from culture of adult rat seminiferous tubules (Henriksen *et al.* 1996) and of immature rat testis fragments (Boitani *et al.* 1993) indicates that FSH also enhances proliferation of differentiated spermatogonia *in vitro*. The significant reduction in the spermatogonial population after 4 days of FSH suppression in the current study provides evidence that FSH influences survival of germ cell populations from 14–18 dpp. The reduction in germ cell number reduction appears attributable to a 2.5-fold higher apoptotic index compared with control levels, not to a change in proliferation rate. We also observed that spermatogonial numbers were not influenced by FSH at 9 days after birth, and this finding is inconsistent with *in vitro* data (Boitani *et al.* 1995). Low levels of circulating FSH may be present in the animals examined in this study; this may be sufficient to support spermatogonial development, as small increases in FSH can significantly impact the recovery of





**Figure 6** Dynamic effects of FSH on Sertoli and germ cell development during first wave in rats assessed through *in vivo* and *in vitro* analyses. Time points listed illustrate progressive maturation of germ cells and Sertoli cells in the first wave of spermatogenesis in rat. At 3 dpp, Sertoli cells are proliferating and gonocytes are re-entering the cell cycle and migrating to the basement membrane of the seminiferous cord to become undifferentiated and stem cell spermatogonia. At 9 dpp, proliferating Sertoli cells and differentiating spermatogonia are observed. By 18 dpp, Sertoli cell proliferation has ceased, and the cords have transformed into seminiferous tubules as tight junctions form between Sertoli cells and vectorial secretion from this epithelium commences. Germ cells that have progressed to the pachytene stage of meiosis are predominant in the seminiferous tubules. In this study, suppression of FSH by passive immunisation for 2 and 4 days revealed the age-dependent differential *in vivo* affects of FSH on Sertoli and germ cells: promotion of Sertoli cell mitosis at 9 dpp and increasing germ cell viability at 18 dpp. These data are in agreement with previous *in vitro* observations that FSH regulates Sertoli cell proliferation in an age-dependent manner, interacting with activin to promote Sertoli proliferation and germ cell development at discrete ages. This comparison also highlights differences that may specifically reflect the use of *in vitro* versus *in vivo* models.

spermatogonial number in adult rats (Meachem *et al.* 1998). In the present study, spermatogonial number was assessed using a sensitive, unbiased optical disector method appropriate for counting these cells. Effects of lowered FSH levels may have been masked due to factors associated with counting a small population of cells. Spermatogonial and spermatocyte numbers in control antibody-treated rats in this study were in accord with our (Meachem *et al.* 1996) and that of others (Simorangkir *et al.* 1997) reporting data that were assessed in rats of similar ages by the using the optical disector technique.

Age dependent changes in cell survival resulting from FSH suppression may be due to developmentally regulated changes in the cellular synthesis of apoptotic regulators, such that documented for members of the Bcl-2 family during the first wave of mouse spermatogenesis (Meehan *et al.* 2001). The *Bcl-w* (Yan *et al.* 2000) and *Bok* (Suominen *et al.* 2001) mRNAs were shown to be up- and down-regulated respectively by FSH *in vitro*, and the *bcl-w* knockout mouse has elevated germ cell apoptosis following

14 dpp, indicating the potential influence of Bcl-2 family members on the cellular responses in this model.

*Identification of genes regulated by FSH in vivo*

Interrogation of a rat gene microarray with RNA from 18 dpp rats in this study identified 60 genes that were regulated by 4 days of FSH suppression *in vivo*. Twenty percent of these differentially expressed genes have been previously identified as FSH-regulated, predominantly through *in vitro* methods, a finding which indicates the validity of this model and experimental approach. For example, in an analysis of 20 dpp rat Sertoli cells using the same microarray platform, FSH-responsive mRNAs were identified at 2, 4, 8 and 24 h of culture (McLean *et al.* 2002). In accord with the present analysis, that *in vitro* analysis identified the genes encoding steroidogenic acute regulatory (StAR) protein and endothelin as down- and up-regulated, respectively, in the absence of FSH. FSH has now been identified through both *in vitro*

(McLean *et al.* 2002) and *in vivo* studies to be one of the wide variety of transcriptional regulators of StAR mRNA synthesis (Gregory & De Phillip 1998, Manna *et al.* 2003). Cyclin D2 and androgen binding protein (ABP) are additional Sertoli cell proteins up-regulated by FSH that are required for normal testicular function (Skinner *et al.* 1989, Danzo 1995, Larriba *et al.* 1995, Sicinski *et al.* 1996, Joseph *et al.* 1997), while protein C inhibitor, also required for spermatogenesis (Uhrin *et al.* 2000, Odet *et al.* 2003), appears to be synthesized only in germ cells and Leydig cells (Odet *et al.* 2003). *In vivo* confirmation of the negative regulation by FSH of genes encoding insulin-like growth factor binding protein-3 (IGFBP-3) (Smith *et al.* 1990, Rappaport & Smith 1995, Khan *et al.* 2002), HNF-3/forkhead (Wolfrum *et al.* 2003), transferrin (Kaestner *et al.* 1998), endothelin-1 (Fantoni *et al.* 1993, Tripiciano *et al.* 1999) and cathepsin-L (Penttila *et al.* 1995, Wright *et al.* 2003) is presented. Members of the HNF-3/forkhead transcription factor family are part of a signal transduction cascade from Akt/protein kinase B that regulates transcription of apoptosis-related genes (Wolfrum *et al.* 2003). HNF-gamma is known to regulate transferrin (Kaestner *et al.* 1998), a gene that is itself regulated by FSH (Migrenne *et al.* 2003); we hypothesize that HNF-3 represents an intermediate for FSH regulation of several genes.

The high concordance of the present data with previous reports provides validation of the methodological approach used to identify *in vivo* targets of FSH regulation, though some differences may be noted. In this study, FSH suppression led to elevated testicular N-cadherin mRNA, but in a previous 2 day *in vitro* analysis, FSH alone did not affect N-cadherin expression; FSH in combination with testosterone stimulated it (Lampa *et al.* 1999). This most likely reflects a specific difference between *in vitro* and *in vivo* models regarding hormone responses; alternatively, the duration of suppression or addition may underpin the difference between these two results.

In addition, novel candidates for FSH regulation have been uncovered, some of which may be indirectly regulated by altered levels of available FSH or represent bona fide novel FSH-regulated genes. Targets not previously identified in the testis include aminolevulinic acid synthase 2 and melanoma inhibitory activity.

Several categories of encoded gene products within the array of candidate FSH target genes may aid identification of the pathways and processes that underpin the reduction in germ cell survival that was measured after 4 days of treatment with the FSHAb. Most striking are the genes relating to cell cycle regulation and apoptosis (e.g. lamin, cyclin D1, cyclin D2, Scavenger Receptor B1). Products of other FSH target genes comprise or influence the architecture of the seminiferous epithelium (e.g. cadherin, cathepsin L), while others are known to function in signal transduction (e.g. HNF3, MAPKKK, Smad 3, rat retinol-binding protein) and to participate in regulation of hor-

monal inputs (StAR and androgen binding protein). However, because the microarray analysis presented in this study addresses a single time point, early and acutely affected FSH gene targets may be undetected. The cellular responses identified in this study (i.e. germ cell apoptosis) would have been initiated before the time point of analysis, so future studies will address this in order to more fully comprehend the pathways through which FSH levels influence Sertoli cell function and germ cell survival.

The limit for microarray signal changes considered to be significant was a 1.5-fold difference between the control and treatment groups. The highest difference detected was 6.57-fold, with several others showing around a 4-fold difference. While this level of change is low in comparison to systems where a single cell type is tested, these data do demonstrate that this approach can identify FSH-regulated genes in a complex cell mixture (i.e. total testis). This is, in fact, comparable to the situation that might be required for analysis of clinical samples, where the small sample size would preclude the use of cell separation methods prior to analysis. The fact that up- and down-regulation of genes is measurable following 4 days of treatment indicates that in conditions where FSH suppression has been long term, FSH target gene expression interrogation may be a useful tool for identifying the basis of pathological change. It is important to note that some alterations in gene expression may be in part attributed to the reduction in testicular weight as a consequence of cell loss. However, we believe the disproportionate change in gene expression (1.5–6 fold) compared with the 22% reduction in testis weight following FSH suppression indicates that the changes in gene expression are primarily due to specific FSH effects.

In conclusion, the model used in this study has successfully identified distinct developmental responses to FSH in Sertoli and germ cells in the postnatal rat and identified candidate genes that may underpin these responses at 18 dpp. These data illustrate the switch in Sertoli cell function during the first spermatogenic wave, as they cease proliferation and establish a niche for support of germ cell development. In addition, the emerging dependence of germ cells on Sertoli cell-derived products can be explored through identification of FSH-target genes and the pathways in which they function.

## Acknowledgements

We thank Liana Nagley for the preparation of RNA prior to microarray analysis, Sue Hayward for measurement of inhibin levels and Dr Ingrid Sadler-Riggelman for her technical assistance.

## Funding

Supported by the Wellcome Trust Fellow Scheme (#058479 to S M), and The National Health and Medical

Research Council of Australia, Program Grants (#050387 to S J M; #1147386 to K L L) and Research Fellowship (#143792 to K L L). The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

## References

- Andersen OM, Yeung CH, Vorum H, Wellner M, Andreassen TK, Erdmann B, Mueller EC, Herz J, Otto A, Cooper TG & Willnow TE 2003 Essential role of the apolipoprotein E receptor-2 in sperm development. *Journal of Biological Chemistry* **278** 23989–23995.
- Arklblad EL, Egorov M, Shakhparonov M, Romanova L, Polzikov M & Rydstrom J 2002 Expression of proton-pumping nicotinamide nucleotide transhydrogenase in mouse, human brain and C elegans. *Comparative Biochemistry and Physiology. Part B, Biochemistry and Molecular Biology* **133** 13–21.
- Asuncion L, Fogelgren B, Fong KS, Fong SF, Kim Y & Csiszar K 2001 A novel human lysyl oxidase-like gene (LOXL4) on chromosome 10q24 has an altered scavenger receptor cysteine rich domain. *Matrix Biology* **20** 487–491.
- Baker PJ & O'Shaughnessy PJ 2001 Role of gonadotrophins in regulating numbers of Leydig and Sertoli cells during fetal and postnatal development in mice. *Reproduction* **122** 227–234.
- Beumer TL, Roepers-Gajadien HL, Gademan IS, Kal HB & de Rooij DG 2000 Involvement of the D-type cyclins in germ cell proliferation and differentiation in the mouse. *Biology of Reproduction* **63** 1893–1898.
- Boitani C, Politi MG & Menna T 1993 Spermatogonial cell proliferation in organ culture of immature rat testis. *Biology of Reproduction* **48** 761–767.
- Boitani C, Stefanini M, Fragale A & Morena AR 1995 Activin stimulates Sertoli cell proliferation in a defined period of rat testis development. *Endocrinology* **136** 5438–5444.
- Boussouar F, Mauduit C, Tabone E, Pellerin L, Magistretti PJ & Benahmed M 2003 Developmental and hormonal regulation of the monocarboxylate transporter 2 (MCT2) expression in the mouse germ cells. *Biology of Reproduction* **69** 1069–1078.
- Bravo-Moreno JF, Diaz-Sanchez V, Montoya-Flores JG, Lamoyi E, Saez JC & Perez-Armendariz EM 2001 Expression of connexin43 in mouse Leydig, Sertoli and germinal cells at different stages of postnatal development. *Anatomical Record* **264** 13–24.
- Bruce S, Bocquet J & Villers C 2002 Regulation of glypican-1, syndecan-1 and syndecan-4 mRNAs expression by follicle-stimulating hormone, cAMP increase and calcium influx during rat Sertoli cell development. *European Journal of Biochemistry* **269** 3461–3469.
- Buemer TL, Roepers-Gajadien HL, Gademan IS, Kal HB & de Rooij DG 2000 Involvement of the D-type cyclins in germ cell proliferation and differentiation in the mouse. *Biology of Reproduction* **63** 1893–1898.
- Buzzard JJ, Wreford NG & Morrison JR 2002 Marked extension of proliferation of rat Sertoli cells in culture using recombinant human FSH. *Reproduction* **124** 633–634.
- Buzzard JJ, Farnworth PG, De Kretser DM, O'Connor AE & Wreford NG 2003 Proliferative phase Sertoli cells display a developmentally regulated response to activin *in vitro*. *Endocrinology* **144** 474–483.
- Chaudhary J, Sadler-Riggelman I, Ague JM & Skinner MK 2005 The Helix-Loop-Helix inhibitor of differentiation proteins induce post-mitotic terminally differentiated Sertoli cells to re-enter the cell cycle and proliferate. *Biology of Reproduction* **72** 1205–1217.
- Chen JK & Heckert LL 2001 Dmrt1 expression is regulated by follicle-stimulating hormone and phorbol esters in postnatal Sertoli cells. *Endocrinology* **142** 1167–1178.
- Chomczynski P & Sacchi N 1987 Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry* **162** 156–159.
- Cooke PS & Meisami E 1991 Early hypothyroidism in rats causes increased adult testis and reproductive organ size but does not change testosterone levels. *Endocrinology* **129** 237–243.
- Danzo BJ 1995 The effects of a gonadotropin-releasing hormone antagonist on androgen-binding protein distribution and other parameters in the adult male rat. *Endocrinology* **136** 4004–4011.
- Dufour JM & Kim KH 1999 Cellular and subcellular localisation of six retinoid receptors in rat testis during postnatal development: identification of potential heterodimeric receptors. *Biology of Reproduction* **61**:1300–1308.
- Eisen MB, Spellman PT, Brown PO & Botstein D 1998 Cluster analysis and display of genome-wide expression patterns *PNAS* **95** 14863–14888.
- Ericsson J, Runquist M, Thelin A, Andersson M, Chojnacki T & Dallner G 1993 Distribution of prenyltransferases in rat tissues. Evidence for a cytosolic all-trans-geranylgeranyl diphosphate synthase. *Journal of Biological Chemistry* **268** 832–838.
- Eskild W, Troen G, Blaner WS, Nilsson A & Hansson V 2000 Evidence for independent control at the mRNA and protein levels of cellular retinol binding protein 1 in rat Sertoli cells. *Journal of Reproduction and Fertility* **119** 101–109.
- Fantoni G, Morris PL, Forti G, Vannelli GB, Orlando C, Barni T, Sestini R, Danza G & Maggi M 1993 Endothelin-1: a new autocrine/paracrine factor in rat testis. *American Journal of Physiology – Endocrinology and Metabolism* **265** E267–E274.
- Fragale A, Puglisi R, Morena AR, Stefanini M & Boitani C 2001 Age-dependent activin receptor expression pinpoints activin A as a physiological regulator of rat Sertoli cell proliferation. *Molecular Human Reproduction* **71** 107–114.
- Goddard I, Florin A, Mauduit C, Tabone E, Contard P, Bars R, Chuzel F & Benahmed M 2003 Alteration of lactate production and transport in the adult rat testis exposed in utero to flutamide. *Molecular Cell Endocrinology* **206** 137–146.
- Gregory CW & De Phillip RM 1998 Detection of steroidogenic acute regulatory protein (StAR) in mitochondria of cultured rat Sertoli cells incubated with follicle-stimulating hormone. *Biology of Reproduction* **58** 470–474.
- Grima J, Wong CC, Zhu LJ, Zong SD & Cheng CY 1998 Testin secreted by Sertoli cells is associated with the cell surface, and its expression correlates with the disruption of Sertoli-germ cell junctions but not the inter-Sertoli tight junction. *Journal of Biological Chemistry* **273** 21040–21053.
- Henriksen K, Kulmala J, Toppari J, Mehrotra K & Parvinen M 1996 Stage-specific apoptosis in the rat seminiferous epithelium: quantification of irradiation effects. *Journal of Andrology* **17** 394–402.
- Hu J, Shima H & Nakagawa H 1999 Gial cell line-derived neurotrophic factor stimulates Sertoli cell proliferation in the early postnatal period of rat testis development. *Endocrinology* **140** 3416–3421.
- Huang HF, Pogach L, Giglio W, Nathan E & Seebode J 1992 GnRH-A induced arrest of spermiogenesis in rats is associated with altered androgen binding protein distribution in the testis and epididymis. *Journal of Andrology* **13** 153–159.
- Ikeda Y & Terashima T 1999 Expression of reelin, the gene responsible for the reeler mutation, in embryonic development and adulthood in the mouse. *Developmental Dynamics* **210** 157–172.
- Jackson VN, Price NT, Carpenter L & Halestrap AP 1997 Cloning of the monocarboxylate transporter isoform MCT2 from rat testis provides evidence that expression in tissues is species-specific and may involve post-transcriptional regulation. *Biochemical Journal* **324** 447–453.
- Joseph DR, O'Brien DA, Sullivan PM, Becchis M, Tsuruta JK & Petrusz P 1997 Overexpression of androgen-binding protein/sex

- hormone-binding globulin in male transgenic mice: tissue distribution and phenotypic disorders. *Biology of Reproduction* **56** 21–32.
- Kabbaj O, Holm C, Vitale C & Pelletier RM 2001 Expression, activity and subcellular localization of testicular hormone-sensitive lipase during postnatal development in the guinea pig. *Biology of Reproduction* **65** 601–612.
- Kaestner KH, Hiemisch H & Schutz G 1998 Targeted disruption of the gene encoding hepatocyte nuclear factor 3 gamma results in reduced transcription of hepatocyte-specific genes. *Molecular Cell Biology* **18** 4245–4251.
- von der Kammer H, Hanes J, Klaudiny J & Scheit KH 1994 A human amyloid precursor-like protein is highly homologous to a mouse sequence-specific DNA-binding protein. *DNA Cell Biology* **13** 1137–1143.
- Kastner P, Mark M, Leid M, Gansmuller A, Chin W, Grondona JM, Decimo D, Krezel W, Dierich A & Chambon P 1996 Abnormal spermatogenesis in RXR beta mutant mice. *Genes and Development* **10** 80–92.
- Kato K, Shinohara H, Kurobe N, Inaguma Y, Shimizu K & Ohshima K 1991 Tissue distribution and developmental profiles of immunoreactive alpha B crystallin in the rat determined with a sensitive immunoassay system. *Biochimica et Biophysica Acta* **1074** 201–208.
- Kawasaki Y, Nakagawa A, Nagaosa K, Shiratsuchi A & Nakanishi Y 2002 Phosphatidylserine binding of class B scavenger receptor type I, a phagocytosis receptor of testicular Sertoli cells. *Journal of Biological Chemistry* **277** 27559–27566.
- Kezele PR, Ague JM, Nilsson E & Skinner MK 2002 Alterations in the ovarian transcriptome during primordial follicle assembly and development. *Biology of Reproduction* **72** 241–255.
- Khan SA, Ndjountche L, Pratchard L, Spicer LJ & Davis JS 2002 Follicle-stimulating hormone amplifies insulin-like growth factor I-mediated activation of AKT/protein kinase B signaling in immature rat Sertoli cells. *Endocrinology* **143** 2259–2267.
- Kirby JD, Jetton AE, Cooke PS, Hess RA, Bunick D, Ackland JF, Turek FW & Schwartz NB 1992 Developmental hormonal profiles accompanying the neonatal hypothyroidism-induced increase in adult testicular size and sperm production in the rat. *Endocrinology* **131** 559–565.
- Lampa J, Hoogerbrugge JW, Baarends WM, Stanton PG, Perryman KJ, Grootegoed JA & Robertson DM 1999 Follicle-stimulating hormone and testosterone stimulation of immature and mature Sertoli cells *in vitro*: inhibin and N-cadherin levels and round spermatid binding. *Journal of Andrology* **20** 399–406.
- Larriba S, Esteban C, Toran N, Gerard A, Audi L, Gerard H & Reventos J 1995 Androgen binding protein is tissue-specifically expressed and biologically active in transgenic mice. *Journal of Steroid Biochemistry and Molecular Biology* **53** 573–578.
- Li S, Moore CL, Dobretsova A & Wight PA 2002 Myelin proteolipid protein (Plp) intron 1 DNA is required to temporally regulate Plp gene expression in the brain. *Journal of Neurochemistry* **83** 193–201.
- Lui WY, Mruk D, Lee WM & Cheng CY 2003 Sertoli cell tight junction dynamics: their regulation during spermatogenesis. *Biology of Reproduction* **68** 1087–1097.
- Machiels BM, Ramaekers FC, Kuijpers HJ, Groenewoud JS, Oosterhuis JW & Looijenga LH 1997 Nuclear lamin expression in normal testis and testicular germ cell tumours of adolescents and adults. *Journal of Pathology* **182** 197–204.
- Maki JM, Tikkanen H & Kivirikko KI 2001 Cloning and characterization of a fifth human lysyl oxidase isoenzyme: the third member of the lysyl oxidase-related subfamily with four scavenger receptor cysteine-rich domains. *Matrix Biology* **20** 493–496.
- Manna P, Wang X-J & Stocco DM 2003 Involvement of multiple transcription factors in the regulation of steroidogenic acute regulatory protein gene expression. *Steroids* **68** 1125–1134.
- Mbakay M, Tadros H, Ishida N, Lerner CP, De Lamirande E, Chen A, El-Alfy M, Clermont Y, Seidah NG, Chretien M, Gagnon C & Simpson EM 1997 Impaired fertility in mice deficient for the testicular germ-cell protease PC4. *PNAS* **94** 6842–6846.
- McClery SM & Clarke SC 2003 The use of hydrolysis and hairpin probes in real-time PCR. *Molecular Biotechnology* **25** 267–274.
- McLachlan RI, Wreford NG, De Kretser DM & Robertson DM 1995 The effects of recombinant follicle-stimulating hormone on the restoration of spermatogenesis in the gonadotropin-releasing hormone-immunized adult rat. *Endocrinology* **136** 4035–4043.
- McLachlan RI, Wreford NG, Meachem SJ, De Kretser DM & Robertson DM 1994 Effects of testosterone on spermatogenic cell populations in the adult rat. *Biology of Reproduction* **51** 945–955.
- McLean DJ, Friel PJ, Pouchnik D & Griswold MD 2002 Oligonucleotide microarray analysis of gene expression in follicle-stimulating hormone-treated rat Sertoli cells. *Molecular Endocrinology* **16** 2780–2792.
- Meachem SJ, McLachlan RI, de Kretser DM, Robertson DM & Wreford NG 1996 Neonatal exposure of rats to recombinant human follicle-stimulating hormone increases adult Sertoli and spermatogenic cell numbers. *Biology of Reproduction* **54** 36–44.
- Meachem SJ, Wreford NG, Stanton PG, Robertson DM & McLachlan RI 1998 Follicle-stimulating hormone is required for the initial phase of spermatogenic restoration in adult rats following gonadotrophin suppression. *Journal of Andrology* **19** 725–735.
- Meachem SJ, McLachlan RI, Stanton PG, Robertson DM & Wreford NG 1999 FSH immunoneutralization acutely impairs spermatogonial development in normal adult rats. *Journal of Andrology* **20** 756–762.
- Meehan T, Schlatt S, O'Bryan MK, de Kretser DM & Loveland KL 2000 Regulation of germ cell and Sertoli cell development by activin, follistatin and FSH. *Developmental Biology* **220** 225–237.
- Meehan T, Loveland KL, de Kretser D, Cory S & Print CG 2001 Developmental regulation of the bcl-2 family during spermatogenesis: Insights into the sterility of bcl-w<sup>-/-</sup> male mice. *Cell Death and Differentiation* **8** 225–233.
- Meng X, Lindahl M, Hyvonen ME, Parvinen M, de Rooij DG, Hess MW, Raatikainen-Ahokas A, Sainio K, Rauvala H, Lakso M, Pichel JG, Westphal H, Saarma M & Sariola H 2000 Regulation of cell fate decision of undifferentiated spermatogonia by GDNF. *Science* **287** 1489–1493.
- Migrenne S, Racine C, Guillou F & Habert R 2003 Pituitary hormones inhibit the function and differentiation of fetal Sertoli cells. *Endocrinology* **144** 2617–2622.
- Monaco L, Foulkes NS & Sassone-Corsi P 1995 Pituitary follicle-stimulating hormone (FSH) induces CREM gene expression in Sertoli cells: involvement in long-term desensitization of the FSH receptor. *PNAS* **92** 10673–10677.
- Mori T, Guo MW, Sato E, Baba T, Takasaki S & Mori E 2000 Molecular and immunological approaches to mammalian fertilization. *Journal of Reproductive Immunology* **47** 139–158.
- Munsie M, Schlatt S, deKretser DM & Loveland KL 1997 Expression of stem cell factor in the postnatal rat testis. *Molecular Reproduction and Development* **47** 19–25.
- Nozaki H, Matsuzawa T, Nakamura T, Arai I & Urashima T 2003 Are there two forms of beta 2-N-acetylglucosaminyltransferase I in rat testicular and epididymal fluids? *Biochimica et Biophysica Acta* **1649** 140–145.
- O'Donnell L, McLachlan RI, Wreford NG & Robertson DM 1994 Testosterone promotes the conversion of round spermatids between stages VII and VIII of the rat spermatogenic cycle. *Endocrinology* **135** 2608–2614.
- Odet F, Guyot R, Leduque P & Le Magueresse-Battistoni B 2003 Evidence for Similar Expression of Protein C Inhibitor and the Urokinase-Type Plasminogen Activator System during Mouse Testis Development. *Endocrinology* **145** 1481–1489.



- Orth JM 1984 The role of follicle-stimulating hormone in controlling Sertoli cell proliferation in testes of foetal rats. *Endocrinology* **115** 1248–1255.
- Orth JM, Gunsalus GL & Lamperti AA 1988 Evidence from Sertoli cell-depleted rats indicates that spermatid number in adults depends on numbers of Sertoli cells produced during perinatal development. *Endocrinology* **122** 787–794.
- Orth JM, Jester WF, Li LH & Laslett AL 2000 Gonocyte-Sertoli cell interactions during development of the neonatal rodent testis. *Current Topics in Developmental Biology* **50** 103–124.
- Orwig KE, Ryu BY, Avarbock MR & Brinster RL 2002 Male germ-line stem cell potential is predicted by morphology of cells in neonatal rat testes. *PNAS* **99** 11706–11711.
- O'Shaughnessy PJ, Willerton L & Baker PJ 2002 Changes in Leydig cell gene expression during development in the mouse. *Biology of Reproduction* **66** 966–975.
- Packer AI, Besmer P & Bachvarova RF 1995 Kit ligand mediates survival of type A spermatogonia and dividing spermatocytes in postnatal mouse testes. *Molecular Reproduction and Development* **42** 303–310.
- Pellegrini M, Grimaldi P, Rossi P, Geremia R & Dolci S 2003 Developmental expression of BMP4/ALK3/SMAD5 signaling pathway in the mouse testis: a potential role of BMP4 in spermatogonia differentiation. *Journal of Cell Science* **116** 3363–3372.
- Penttilä TL, Hakovirta H, Mali P, Wright WW & Parvinen M 1995 Follicle-stimulating hormone regulates the expression of cyclic protein-2/cathepsin L messenger ribonucleic acid in rat Sertoli cells in a stage-specific manner. *Molecular and Cellular Endocrinology* **113** 175–181.
- Petersen C, Boitani C, Froya B & Soder O 2001 Transforming growth factor- $\alpha$  stimulates proliferation of rat Sertoli cells. *Molecular and Cellular Endocrinology* **181** 221–227.
- Petersen C, Boitani C, Froya B & Soder O 2002 Interleukin-1 is a potent growth factor for immature rat Sertoli cells. *Molecular and Cellular Endocrinology* **186** 37–47.
- Rappaport MS & Smith EP 1995 Insulin-like growth factor (IGF) binding protein 3 in the rat testis: follicle-stimulating hormone dependence of mRNA expression and inhibition of IGF-I action on cultured Sertoli cells. *Biology of Reproduction* **52** 419–425.
- Raymond CS, Murphy MW, O'Sullivan MG, Bardwell VJ & Zarkower D 2000 Dmrt1, a gene related to worm and fly sexual regulators is required for mammalian testis differentiation. *Genes and Development* **14** 2587–2595.
- Reaven E, Zhan L, Nomoto A, Leers-Sucheta S & Azhar S 2000 Expression and microvillar localization of scavenger receptor class B, type I (SR-BI) and selective cholesteryl ester uptake in Leydig cells from rat testis. *Journal of Lipid Research* **41** 343–356.
- Robertson DM, Hayward S, Irby D, Jacobsen J, Clarke L, McLachlan RI & de Kretser DM 1988 Radioimmunoassay of rat serum inhibin: changes after PMSG stimulation and gonadectomy. *Molecular Cellular Endocrinology* **58** 1–8.
- Robertson DM, Giacometti, M, Foulds L, Lahnstein J, Goss NH, Hearn MTW & de Kretser DM 1989 Isolation of inhibin  $\alpha$ -subunit precursor proteins from bovine follicular fluid. *Endocrinology* **125** 2141–2149.
- de Rooij DG 1998 Stem cells in testis. *International Journal of Experimental Pathology* **79** 67–80.
- Rossi P, Dolci S, Albanesi C, Grimaldi P, Ricca R & Geremia R 1993 Follicle-stimulating hormone induction of steel factor (SLF) mRNA in mouse Sertoli cells and stimulation of DNA synthesis in spermatogonia by soluble SLF. *Developmental Biology* **155** 68–74.
- Russell LD, Alger LE & Nequin LG 1987 Hormonal control of pubertal spermatogenesis. *Endocrinology* **120** 1615–1632.
- Russell LD, Ettlín RA, Sinha Hikim AP & Clegg ED 1990 *Histological and histopathological evaluation of the testis*. Ed. LD Russell. Clearwater, Florida: Cache River Press.
- Sadate-Ngatchou PI, Pouchnik DJ & Griswold MD 2004 Identification of testosterone-regulated genes in testes of hypogonadal mice using oligonucleotide microarray *Molecular Endocrinology* **18** 422–433.
- Schinkel AH & Jonker JW 2003 Mammalian drug efflux transporters of the ATP binding cassette (ABC) family: an overview. *Advanced Drug Delivery Reviews* **55** 3–29.
- Schlatt S, Zhengwei Y, Meehan T, de Kretser DM & Loveland KL 1999 Application of morphometric techniques to postnatal rat testes in organ culture: insights into testis growth. *Cell and Tissue Research* **298** 335–343.
- Shetty J, Marathe GK & Dighe RR 1996 Specific immunoneutralization of FSH leads to apoptotic cell death of the pachytene spermatocytes and spermatogonial cells in the rat. *Endocrinology* **137** 2179–2182.
- Shima JE, McLean DJ, McCarrey JR & Griswold MD 2004 The murine testicular transcriptome: Characterizing gene expression in the testis during the Progression of spermatogenesis. *Biology of Reproduction* **71** 319–330.
- Sicinski P, Donaher JL, Geng Y, Parker SB, Gardner H, Park MY, Robker RL, Richards JS, McGinnis LK, Biggers JD, Eppig JJ, Bronson RT, Elledge SJ & Weinberg RA 1996 Cyclin D2 is an FSH-responsive gene involved in gonadal cell proliferation and oncogenesis. *Nature* **384** 470–474.
- Simorangkir DR, Wreford NG & de Kretser DM 1997 Impaired germ cell development in the testes of immature rat with neonatal hypothyroidism. *Journal of Andrology* **18** 186–193.
- Skinner MK, Schlitz SM & Anthony CT 1989 Regulation of Sertoli cell differentiated function: testicular transferrin and androgen-binding protein expression. *Endocrinology* **124** 3015–3024.
- Smith EP, Dickson BA & Chernauek SD 1990 Insulin-like growth factor binding protein-3 secretion from cultured rat Sertoli cells: dual regulation by follicle stimulating hormone and insulin-like growth factor-I. *Endocrinology* **127** 2744–2751.
- Sommersberg B, Bulling A, Salzer U, Frohlich U, Garfield RE, Amsterdam A & Mayerhofer A 2000 Gap junction communication and connexin 43 gene expression in a rat granulosa cell line: regulation by follicle-stimulating hormone. *Biology of Reproduction* **63** 1661–1668.
- Strothmann K, Simoni M, Mathur P, Siakhamary S, Nieschlag E & Gromoll J 2004 Gene expression profiling of mouse Sertoli cell lines. *Cell Tissue Research* **315** 249–257.
- Suominen JS, Yan W, Toppari J & Kaipia A 2001 The expression and regulation of Bcl-2-related ovarian killer (Bok) mRNA in the developing and adult rat testis. *European Journal of Endocrinology* **145** 771–778.
- Swinnen JV, Tsikalas KE & Conti M 1991 Properties and hormonal regulation of two structurally related cAMP phosphodiesterases from the rat Sertoli cell. *The Journal of Biological Chemistry* **266** 18370–18377.
- Tadokoro Y, Yomogida K, Ohta H, Tohda A & Nishimune Y 2002 Homeostatic regulation of germinal stem cell proliferation by the GDNF/FSH pathway. *Mechanisms of Development* **113** 29–39.
- Tapanainen JS, Tilly JL, Vihko KK & Hsueh AJ 1993 Hormonal control of apoptotic cell death in the testis: gonadotropins and androgens as testicular cell survival factors. *Molecular Endocrinology* **76** 43–50.
- Tripiciano A, Peluso C, Morena AR, Palombi F, Stefanini M, Ziparo E, Yanagisawa M & Filippini A 1999 Cyclic expression of endothelin-converting enzyme-1 mediates the functional regulation of seminiferous tubule contraction. *Journal of Cell Biology* **145** 1027–1038.
- Uhrin P, Dewerchin M, Hilpert M, Chrenek P, Schofer C, Zechmeister-Machhart M, Kronke G, Vales A, Carmeliet P, Binder BR & Geiger M 2000 Disruption of the protein C inhibitor gene results in impaired spermatogenesis and male infertility. *Journal of Clinical Investigation* **106** 1531–1539.

- Ungefroren H, Ergun S, Krull NB & Holstein AF 1995 Expression of the small proteoglycans biglycan and decorin in the adult human testis. *Biology of Reproduction* **52** 1095–1105.
- Van Haaster LH, De Jong FH, Docter R & de Rooij DG 1992 The effect of hypothyroidism on Sertoli cell proliferation and differentiation and hormone levels during testicular development in the rat. *Endocrinology* **131** 1574–1576.
- Walker WH, Sanborn BM & Habener JF 1994 An isoform of transcription factor CREM expressed during spermatogenesis lacks the phosphorylation domain and represses cAMP-induced transcription. *PNAS* **91** 12423–12427.
- West AP, Sharpe RM & Saunders PT 1994 Differential regulation of cyclic adenosine 3',5'-monophosphate (cAMP) response element-binding protein and cAMP response element modulator messenger ribonucleic acid transcripts by follicle-stimulating hormone and androgen in the adult rat testis. *Biology of Reproduction* **50** 869–881.
- Wolfrum C, Besser D, Luca E & Stoffel M 2003 Insulin regulates the activity of forkhead transcription factor Hnf-3 beta/Foxa-2 by Akt-mediated phosphorylation and nuclear/cytosolic localization. *PNAS* **100** 11624–11629.
- Wolgemuth DJ, Besset V, Liu D, Zhang Q & Rhee K 1998 Role of cell-cycle genes in the regulation of mammalian meiosis. In *Germ Cell Development, Division, Disruption and Death*, pp 49–60. Ed Zirkin B. New York: Springer-Verlag.
- Wreford NG 1995 Theory and practice of stereological techniques applied to the estimation of cell number and nuclear volume in the testis. *Microscopy Research and Technique* **32** 423–436.
- Wright WW, Smith L, Kerr C & Charron M 2003 Mice that express enzymatically inactive cathepsin L exhibit abnormal spermatogenesis. *Biology of Reproduction* **68** 680–687.
- Xu J, Beyer AR, Walker WH & McGee EA 2003 Developmental and stage-specific expression of Smad2 and Smad3 in rat testis. *Journal of Andrology* **24** 192–200.
- Yan W, Kero J, Suominen J & Toppari J 2001 Differential expression and regulation of the retinoblastoma family of proteins during testicular development and spermatogenesis: roles in the control of germ cell proliferation, differentiation and apoptosis. *Oncogenetics* **20** 1343–1356.
- Yan W, Suominen J, Samson M, Jegou B & Toppari J 2000 Involvement of Bcl-2 family proteins in germ cell apoptosis during testicular development in the rat and pro-survival effect of stem cell factor on germ cells *in vitro*. *Molecular and Cellular Endocrinology* **165** 115–129.
- Yomogida K, Yagura Y, Tadokoro Y & Nishimune Y 2003 Dramatic Expansion of Germinal Stem Cells by Ectopically- Expressed Human Glial Cell Line-Derived Neurotrophic Factor in Mouse Sertoli Cells. *Biology of Reproduction* **69** 1303–1307.
- Yoshinaga K, Nishikawa S, Ogawa M, Hayashi S, Kunisada T, Fujimoto T & Nishikawa S 1991 Role of c-kit in mouse spermatogenesis: identification of spermatogonia as a specific site of c-kit expression and function. *Development* **113** (Suppl) 689–699.

Received 16 May 2005

Accepted 27 June 2005