

Spring 2020 – Systems Biology of Reproduction
Lecture Outline – Epigenetics and Transgenerational Reproductive Disease
Michael K. Skinner – Biol 475/575
CUE 418, 10:35-11:50 am, Tuesday & Thursday
March 10, 2020
Week 9

Epigenetics and Transgenerational Reproductive Disease

- Environmental Induced Pathology
- Transgenerational Disease Phenotype
- Compound Specificity
- Epigenetic Mechanism
- Epimutations and Exposure Specificity
- Transgenerational Transcriptome
- Transgenerational Testis Disease
- Transgenerational Ovary Disease
- Broader Impact

REQUIRED READING

Nilsson EE, Sadler-Riggelman I, Skinner MK. Environmentally induced epigenetic transgenerational inheritance of disease. *Environ Epigenet*. 2018 Jul 17;4(2):1-13, dvy016.

REFERENCES

- Duempelmann L, Skribbe M, Bühler M. (2020) Small RNAs in the Transgenerational Inheritance of Epigenetic Information. *Trends Genet*. 2020 Jan 14. doi: 10.1016/j.tig.2019.12.001. [Epub ahead of print]
- Patel NJ, Hogan KJ, Rizk E, Stewart K, Madrid A, et al. (2020) Ancestral Folate Promotes Neuronal Regeneration in Serial Generations of Progeny. *Mol Neurobiol*. 2020 Jan 10. doi: 10.1007/s12035-019-01812-5. [Epub ahead of print]
- Legoff L, D'Cruz SC, Tevosian S, Primig M, Smagulova F. (2019) Transgenerational Inheritance of Environmentally Induced Epigenetic Alterations during Mammalian Development. *Cells* 3;8(12).
- Meyer DN, Crofts EJ, Akemann C, Gurdziel K, Farr R, Baker BB, Weber D, Baker TR. (2019) Developmental exposure to Pb²⁺ induces transgenerational changes to zebrafish brain transcriptome. *Chemosphere* 2;244:125527.
- King SE, Nilsson E, Beck D, Skinner MK. (2019) Adipocyte epigenetic alterations and potential therapeutic targets in transgenerationally inherited lean and obese phenotypes following ancestral exposures. *Adipocyte*. 8(1):362-378.
- Bodden C, Hannan AJ, Reichelt AC. (2020) Diet-Induced Modification of the Sperm Epigenome Programs Metabolism and Behavior. *Trends Endocrinol Metab*. 31(2):131-149.

- Cavalli G, Heard E. (2019) Advances in epigenetics link genetics to the environment and disease. *Nature*. 571(7766):489-499.
- Reece AS, Hulse GK. (2019) Impacts of cannabinoid epigenetics on human development: reflections on Murphy et. al. 'cannabinoid exposure and altered DNA methylation in rat and human sperm' epigenetics 2018; 13: 1208-1221. *Epigenetics*. 14(11):1041-1056.
- Cong W, Miao Y, Xu L, Zhang Y, Yuan C, et al. (2019) Transgenerational memory of gene expression changes induced by heavy metal stress in rice (*Oryza sativa* L.). *BMC Plant Biol*. 27;19(1):282.
- Sarkies P. (2020) Molecular mechanisms of epigenetic inheritance: Possible evolutionary implications. *Semin Cell Dev Biol*. 97:106-115.
- King SE, McBirney M, Beck D, Sadler-Riggelman I, Nilsson E, Skinner MK. (2019) Sperm epimutation biomarkers of obesity and pathologies following DDT induced epigenetic transgenerational inheritance of disease. *Environ Epigenet*. 27;5(2):dvz008.
- Skinner MK, Nilsson E, Sadler-Riggelman I, Beck D, Ben Maamar M, McCarrey JR. (2019) Transgenerational sperm DNA methylation epimutation developmental origins following ancestral vinclozolin exposure. *Epigenetics*. 14(7):721-739.
- Ben Maamar M, Nilsson E, Sadler-Riggelman I, Beck D, McCarrey JR, Skinner MK. (2019) Developmental origins of transgenerational sperm DNA methylation epimutations following ancestral DDT exposure. *Dev Biol*. 15;445(2):280-293.
- Lacal I, Ventura R. (2018) Epigenetic Inheritance: Concepts, Mechanisms and Perspectives. *Front Mol Neurosci*. 28;11:292.
- Nilsson E, Klukovich R, Sadler-Riggelman I, Beck D, Xie Y, Yan W, Skinner MK. Environmental toxicant induced epigenetic transgenerational inheritance of ovarian pathology and granulosa cell epigenome and transcriptome alterations: ancestral origins of polycystic ovarian syndrome and primary ovarian insufficiency. *Epigenetics*. 2018;13(8):875-895.
- Gold HB, Jung YH, Corces VG. Not just heads and tails: The complexity of the sperm epigenome. *J Biol Chem*. 2018 Sep 7;293(36):13815-13820.
- Ben Maamar M, King SE, Nilsson E, Beck D, Skinner MK. Epigenetic transgenerational inheritance of parent-of-origin allelic transmission of outcross pathology and sperm epimutations. *Dev Biol*. 2020 Feb 1;458(1):106-119.
- Sadler-Riggelman I, Klukovich R, Nilsson E, Beck D, Xie Y, Yan W, Skinner MK. Epigenetic transgenerational inheritance of testis pathology and Sertoli cell epimutations: generational origins of male infertility. *Environ Epigenet*. 2019 Aug 29;5(3):dvz013.
- Kubsad D, Nilsson EE, King SE, Sadler-Riggelman I, Beck D, Skinner MK. Assessment of Glyphosate Induced Epigenetic Transgenerational Inheritance of Pathologies and Sperm Epimutations: Generational Toxicology. *Sci Rep*. 2019 Apr 23;9(1):6372.
- Baxter FA, Drake AJ. Non-genetic inheritance via the male germline in mammals. *Philos Trans R Soc Lond B Biol Sci*. 2019 Apr 15;374(1770):20180118.
- Klukovich R, Nilsson E, Sadler-Riggelman I, Beck D, Xie Y, Yan W, Skinner MK. Environmental Toxicant Induced Epigenetic Transgenerational Inheritance of Prostate Pathology and Stromal-Epithelial Cell Epigenome and Transcriptome Alterations: Ancestral Origins of Prostate Disease. *Sci Rep*. 2019 Feb 18;9(1):2209.
- Lecoutre S, Petrus P, Rydén M, Breton C. Transgenerational Epigenetic Mechanisms in Adipose Tissue Development. *Trends Endocrinol Metab*. 2018 Oct;29(10):675-685.
- Duempelmann L, Skribbe M, Bühler M. Small RNAs in the Transgenerational Inheritance of Epigenetic Information. *Trends Genet*. 2020 Jan 14. pii: S0168-9525(19)30259-8. doi: 10.1016/j.tig.2019.12.001. [Epub ahead of print]

- Stermer AR, Wilson SK, Klein D, Hall SJ, Boekelheide K. Trichloroethylene exposure alters dimethylated histone three lysine four in protein kinase A signaling pathway chromatin of rat sperm†. *Biol Reprod.* 2019 Nov 21;101(5):875-877.
- van Steenwyk G, Roszkowski M, Manuella F, Franklin TB, Mansuy IM. Transgenerational inheritance of behavioral and metabolic effects of paternal exposure to traumatic stress in early postnatal life: evidence in the 4th generation. *Environ Epigenet.* 2018 Oct 16;4(2):dvy023.
- Beck D, Sadler-Riggleman I, Skinner MK. (2017) Generational comparisons (F1 versus F3) of vinclozolin induced epigenetic transgenerational inheritance of sperm differential DNA methylation regions (epimutations) using MeDIP-Seq. *Environ Epigenet.* 3(3). pii: dvx016.
- Holder LB, Haque MM, Skinner MK. (2017) Machine learning for epigenetics and future medical applications. *Epigenetics.* 12(7):505-514.
- Craig JR, Jenkins TG, Carrell DT, Hotaling JM. (2017) Obesity, male infertility, and the sperm epigenome. *Fertil Steril.* 107(4):848-859.
- Tikoo K, Vikram A, Shrivastava S, Jena G, Shah H, Chhabra R. (2017) Parental High-Fat Diet Promotes Inflammatory and Senescence-Related Changes in Prostate. *Oxid Med Cell Longev.* 2017;2017:4962950.
- Bruner-Tran KL, Gnecco J, Ding T, Glore DR, Pensabene V, Osteen KG. (2017) Exposure to the environmental endocrine disruptor TCDD and human reproductive dysfunction: Translating lessons from murine models. *Reprod Toxicol.* 68:59-71.
- Ho SM, Cheong A, Adgent MA, Veevers J, Suen AA, Tam NNC, Leung YK, Jefferson WN, Williams CJ. (2017) Environmental factors, epigenetics, and developmental origin of reproductive disorders. *Reprod Toxicol.* 68:85-104.
- Siddeek B, Lakhdari N, Inoubli L, et al. (2016) Developmental epigenetic programming of adult germ cell death disease: Polycomb protein EZH2-miR-101 pathway. *Epigenomics.* 8(11):1459-1479.
- Mykhalchenko K, Lizneva D, Trofimova T, Walker W, Suturina L, Diamond MP, Azziz R. (2017) Genetics of polycystic ovary syndrome. *Expert Rev Mol Diagn.* 17(7):723-733.
- Filippou P, Homburg R. (2017) Is foetal hyperexposure to androgens a cause of PCOS? *Hum Reprod Update.* 1;23(4):421-432.
- Lawrenson K, Kar S, McCue K, et al. (2016) Functional mechanisms underlying pleiotropic risk alleles at the 19p13.1 breast-ovarian cancer susceptibility locus. *Nat Commun.* 7:12675.
- Kar SP, Beesley J, Amin Al Olama A, et al. (2016) Genome-Wide Meta-Analyses of Breast, Ovarian, and Prostate Cancer Association Studies Identify Multiple New Susceptibility Loci Shared by at Least Two Cancer Types. *Cancer Discov.* 6(9):1052-67.
- Cucchiara V, Yang JC, Mirone V, Gao AC, Rosenfeld MG, Evans CP. (2017) Epigenomic Regulation of Androgen Receptor Signaling: Potential Role in Prostate Cancer Therapy. *Cancers (Basel).* 16;9(1). pii: E9. doi: 10.3390/cancers9010009.
- Sampath Kumar A, Seah MK, Ling KY, Wang Y, Tan JH, Nitsch S, Lim SL, Lorthongpanich C, Wollmann H, Low DH, Guccione E, Messerschmidt DM. (2017) Loss of maternal Trim28 causes male-predominant early embryonic lethality. *Genes Dev.* 31(1):12-17.
- van den Berg GJ, Pinger PR. (2016) Transgenerational effects of childhood conditions on third generation health and education outcomes. *Econ Hum Biol.* 23:103-120.
- Vrooman LA, Bartolomei MS. (2017) Can assisted reproductive technologies cause adult-onset disease? Evidence from human and mouse. *Reprod Toxicol.* 68:72-84.
- Reichetzeder C, Dwi Putra SE, Li J, Hocher B. (2016) Developmental Origins of Disease - Crisis Precipitates Change. *Cell Physiol Biochem.* 39(3):919-38.

- Illum LR, Bak ST, Lund S, Nielsen AL. (2017) DNA methylation in epigenetic inheritance of metabolic diseases through the male germ line. *J Mol Endocrinol*. 2017 Dec 4. pii: JME-17-0189. doi: 10.1530/JME-17-0189.
- McBirney M, King SE, Pappalardo M, Houser E, Unkefer M, Nilsson E, Sadler-Riggelman I, Beck D, Winchester P, Skinner MK. (2017) Atrazine induced epigenetic transgenerational inheritance of disease, lean phenotype and sperm epimutation pathology biomarkers. *PLoS One* 12(9):e0184306.
- McCarrey JR, Lehle JD, Raju SS, Wang Y, Nilsson EE, Skinner MK. (2016) Tertiary Epimutations - A Novel Aspect of Epigenetic Transgenerational Inheritance Promoting Genome Instability. *PLoS One*. 11(12):e0168038.
- Blake GE, Watson ED. (2016) Unravelling the complex mechanisms of transgenerational epigenetic inheritance. *Curr Opin Chem Biol*. 33:101-7.
- Sailer C, Schmid B, Grossniklaus U. (2016) Apomixis Allows the Transgenerational Fixation of Phenotypes in Hybrid Plants. *Curr Biol*. 26(3):331-7.
- Stuppia L, Franzago M, Ballerini P, Gatta V, Antonucci I. (2015) Epigenetics and male reproduction: the consequences of paternal lifestyle on fertility, embryo development, and children lifetime health. *Clin Epigenetics*. 11;7:120.
- Keil KP, Vezina CM. (2015) DNA methylation as a dynamic regulator of development and disease processes: spotlight on the prostate. *Epigenomics*. 7(3):413-25.
- Cahenzli F, Wenk BA, Erhardt A. (2015) Female butterflies adapt and allocate their progeny to the host-plant quality of their own larval experience. *Ecology*. 96(7):1966-73.
- Huo X, Chen D, He Y, Zhu W, Zhou W, Zhang J. (2015) Bisphenol-A and Female Infertility: A Possible Role of Gene-Environment Interactions. *Int J Environ Res Public Health*. 7;12(9):11101-16.
- Erwin AA, Galdos MA, Wickersheim ML, et al. (2015) piRNAs Are Associated with Diverse Transgenerational Effects on Gene and Transposon Expression in a Hybrid Dysgenic Syndrome of *D. virilis*. *PLoS Genet*. 2015 Aug 4;11(8):e1005332.
- Jobson MA, Jordan JM, Sandrof MA, et al. (2015) Transgenerational Effects of Early Life Starvation on Growth, Reproduction, and Stress Resistance in *Caenorhabditis elegans*. *Genetics*. 201(1):201-12.
- Quinnies KM, Doyle TJ, Kim KH1, Rissman EF. (2015) Transgenerational Effects of Di-(2-Ethylhexyl) Phthalate (DEHP) on Stress Hormones and Behavior. *Endocrinology*. 156(9):3077-83.
- Fénichel P, Brucker-Davis F, Chevalier N. (2015) The history of Distilbène® (Diethylstilbestrol) told to grandchildren--the transgenerational effect. *Ann Endocrinol (Paris)*. 76(3):253-9.
- Bale TL. (2015) Epigenetic and transgenerational reprogramming of brain development. *Nat Rev Neurosci*. 16(6):332-44.
- Ziv-Gal A, Wang W, Zhou C, Flaws JA. (2015) The effects of in utero bisphenol A exposure on reproductive capacity in several generations of mice. *Toxicol Appl Pharmacol*. 1;284(3):354-62.
- Tortiglione C. (2014) The heritable effects of nanotoxicity. *Nanomedicine (Lond)*. 9(18):2829-41.
- VandeVoort CA, Grimsrud KN, Midic U, et al. (2015) Transgenerational effects of binge drinking in a primate model: implications for human health. *Fertil Steril*. 103(2):560-9.
- Wei Y, Schatten H, Sun QY. (2015) Environmental epigenetic inheritance through gametes and implications for human reproduction. *Hum Reprod Update*. 21(2):194-208.
- Shama LN, Wegner KM. (2014) Grandparental effects in marine sticklebacks: transgenerational plasticity across multiple generations. *J Evol Biol*. 27(11):2297-307.

- Baker TR, King-Heiden TC, Peterson RE, Heideman W. (2014) Dioxin induction of transgenerational inheritance of disease in zebrafish. *Mol Cell Endocrinol.* 398(1-2):36-41
- Bruner-Tran KL, Ding T, Yeoman KB, Archibong A, Arosh JA, Osteen KG. (2014) Developmental exposure of mice to dioxin promotes transgenerational testicular inflammation and an increased risk of preterm birth in unexposed mating partners. *PLoS One.* 15;9(8):e105084.
- Crews D, Gillette R, Miller-Crews I, Gore AC, Skinner MK. (2014) Nature, nurture and epigenetics. *Mol Cell Endocrinol.* 398(1-2):42-52.
- Kawashima T, Berger F. (2014) Epigenetic reprogramming in plant sexual reproduction. *Nat Rev Genet.* 15(9):613-24.
- Nilsson EE, Skinner MK. (2015) Environmentally induced epigenetic transgenerational inheritance of disease susceptibility. *Transl Res.* 165(1):12-7.
- Grandjean P, Barouki R, Bellinger DC, et al. (2015) Life-Long Implications of Developmental Exposure to Environmental Stressors: New Perspectives. *Endocrinology.* 156(10):3408-15.
- Wei Y, Schatten H, Sun QY. (2015) Environmental epigenetic inheritance through gametes and implications for human reproduction. *Hum Reprod Update.* 21(2):194-208.
- Bao J, Bedford MT. (2016) Epigenetic regulation of the histone-to-protamine transition during Spermiogenesis. *Reproduction.* 2016 Feb 5. pii: REP-15-0562. [Epub ahead of print]
- Tseng YT, Liao HF, Yu CY, Mo CF, Lin SP. (2015) Epigenetic factors in the regulation of prospermatogonia and spermatogonial stem cells. *Reproduction.* 150(3):R77-91.
- Katsi V, Felekos I, Siristatidis C, et al. (2015) Preeclampsia: What Does the Father Have to Do with It? *Curr Hypertens Rep.* 17(8):60.
- Beaujean N. (2014) Epigenetics, embryo quality and developmental potential. *Reprod Fertil Dev.* 27(1):53-62.
- Ge ZJ, Schatten H, Zhang CL, Sun QY. (2015) Oocyte ageing and epigenetics. *Reproduction.* 2015 Mar;149(3):R103-14.
- Vaiserman AM. (2015) Epigenetic programming by early-life stress: Evidence from human populations. *Dev Dyn.* 244(3):254-65.
- Campos EI, Stafford JM, Reinberg D. (2014) Epigenetic inheritance: histone bookmarks across generations. *Trends Cell Biol.* 2014 Nov;24(11):664-74.
- She W, Baroux C. (2014) Chromatin dynamics during plant sexual reproduction. *Front Plant Sci.* 2014 Jul 24;5:354.
- Ding GL, Huang HF. (2013) Paternal transgenerational glucose intolerance with epigenetic alterations in second generation offspring of GDM. *Asian J Androl.* 15(4):451-2.
- Di Giacomo M, Comazzetto S, et al. (2013) Multiple epigenetic mechanisms and the piRNA pathway enforce LINE1 silencing during adult spermatogenesis. *Mol Cell.* 23;50(4):601-8.
- Guerrero-Bosagna C, Savenkova M, Haque MM, Nilsson E, Skinner MK. (2013) Environmentally induced epigenetic transgenerational inheritance of altered Sertoli cell transcriptome and epigenome: molecular etiology of male infertility. *PLoS One.* 8(3):e59922.
- Tracey R, Manikkam M, Guerrero-Bosagna C, Skinner MK. (2013) Hydrocarbons (jet fuel JP-8) induce epigenetic transgenerational inheritance of obesity, reproductive disease and sperm epimutations. *Reprod Toxicol.* 36:104-16.
- Feng X, Zilberman D, Dickinson H. (2013) A conversation across generations: soma-germ cell crosstalk in plants. *Dev Cell.* 11;24(3):215-25.
- Soubry A, Schildkraut JM, et al. (2013) Paternal obesity is associated with IGF2 hypomethylation in newborns: results from a Newborn Epigenetics Study (NEST) cohort. *BMC Med.* 6;11:29.

- Chamorro-García R, Sahu M, et al. (2013) Transgenerational inheritance of increased fat depot size, stem cell reprogramming, and hepatic steatosis elicited by prenatal exposure to the obesogen tributyltin in mice. *Environ Health Perspect.* 121(3):359-66.
- Raychaudhuri N, Dubruille R, et al. (2012) Transgenerational propagation and quantitative maintenance of paternal centromeres depends on Cid/Cenp-A presence in *Drosophila* sperm. *PLoS Biol.* 10(12):e1001434.
- González-Recio O, Ugarte E, Bach A. (2012) Trans-generational effect of maternal lactation during pregnancy: a Holstein cow model. *PLoS One.* 7(12):e51816.
- Hackett JA, Sengupta R, et al. (2013) Germline DNA demethylation dynamics and imprint erasure through 5-hydroxymethylcytosine. *Science.* 25;339(6118):448-52.
- Guerrero-Bosagna C, Covert TR, et al. (2012) Epigenetic transgenerational inheritance of vinclozolin induced mouse adult onset disease and associated sperm epigenome biomarkers. *Reprod Toxicol.* 34(4):694-707.
- Roseboom TJ, Watson ED. (2012) The next generation of disease risk: are the effects of prenatal nutrition transmitted across generations? Evidence from animal and human studies. *Placenta.* 33 Suppl 2:e40-4.
- Uzumcu M, Zama AM, Oruc E. (2012) Epigenetic mechanisms in the actions of endocrine-disrupting chemicals: gonadal effects and role in female reproduction. *Reprod Domest Anim.* 47 Suppl 4:338-47.
- Wolstenholme JT, Edwards M, et al. (2012) Gestational exposure to bisphenol a produces transgenerational changes in behaviors and gene expression. *Endocrinology.* 153(8):3828-38.
- Govorko D, Bekdash RA, Zhang C, Sarkar DK. (2012) Male germline transmits fetal alcohol adverse effect on hypothalamic proopiomelanocortin gene across generations. *Biol Psychiatry.* 1;72(5):378-88.
- Zeh JA, Bonilla MM, et al. (2012) From father to son: transgenerational effect of tetracycline on sperm viability. *Sci Rep.* 2:375.
- Ding GL, Wang FF, et al. (2012) Transgenerational glucose intolerance with Igf2/H19 epigenetic alterations in mouse islet induced by intrauterine hyperglycemia. *Diabetes.* 61(5):1133-42.
- Manikkam M, Guerrero-Bosagna C, Tracey R, Haque MM, Skinner MK. (2012) Transgenerational actions of environmental compounds on reproductive disease and identification of epigenetic biomarkers of ancestral exposures. *PLoS One.* 7(2):e31901.
- van Montfoort AP, Hanssen LL, et al. (2012) Assisted reproduction treatment and epigenetic inheritance. *Hum Reprod Update.* 18(2):171-97.
- Halliday J. (2012) Outcomes for offspring of men having ICSI for male factor infertility. *Asian J Androl.* 14(1):116-20.
- Burdge GC, Hoile SP, Uller T, et al. (2011) Progressive, transgenerational changes in offspring phenotype and epigenotype following nutritional transition. *PLoS One.* 6(11):e28282.
- Low FM, Gluckman PD, Hanson MA. (2011) Developmental plasticity and epigenetic mechanisms underpinning metabolic and cardiovascular diseases. *Epigenomics.* 3(3):279-94.
- Chan D, Delbès G, Landry M, Robaire B, Trasler JM. (2012) Epigenetic alterations in sperm DNA associated with testicular cancer treatment. *Toxicol Sci.* 125(2):532-43.
- Guerrero-Bosagna C, Skinner MK. (2012) Environmentally induced epigenetic transgenerational inheritance of phenotype and disease. *Mol Cell Endocrinol.* 6;354(1-2):3-8.
- Calle A, Fernandez-Gonzalez R, et al. (2012) Long-term and transgenerational effects of in vitro culture on mouse embryos. *Theriogenology.* 1;77(4):785-93.

- Matthews SG, Phillips DI. (2012) Transgenerational inheritance of stress pathology. *Exp Neurol*. 2012 Jan;233(1):95-101.
- Radtke KM, Ruf M, et al. (2011) Transgenerational impact of intimate partner violence on methylation in the promoter of the glucocorticoid receptor. *Transl Psychiatry*. 19;1:e21.
- Stouder C, Paoloni-Giacobino A. (2011) Specific transgenerational imprinting effects of the endocrine disruptor methoxychlor on male gametes. *Reproduction*. 141(2):207-16.
- Skinner MK, Manikkam M, Guerrero-Bosagna C. (2011) Epigenetic transgenerational actions of endocrine disruptors. *Reprod Toxicol*. 31(3):337-43.
- Pentinat T, Ramon-Krauel M, et al. (2010) Transgenerational inheritance of glucose intolerance in a mouse model of neonatal overnutrition. *Endocrinology*. 151(12):5617-23.
- Yazbek SN, Spiezio SH, Nadeau JH, Buchner DA. (2010) Ancestral paternal genotype controls body weight and food intake for multiple generations. *Hum Mol Genet*. 1;19(21):4134-44.
- Zama AM, Uzumcu M. (2010) Epigenetic effects of endocrine-disrupting chemicals on female reproduction: an ovarian perspective. *Front Neuroendocrinol*. 31(4):420-39.
- Dunn GA, Morgan CP, Bale TL. (2011) Sex-specificity in transgenerational epigenetic programming. *Horm Behav*. 59(3):290-5.
- Skinner MK, Manikkam M, Guerrero-Bosagna C. (2010) Epigenetic transgenerational actions of environmental factors in disease etiology. *Trends Endocrinol Metab*. 21(4):214-22.
- Guerrero-Bosagna CM, Skinner MK. (2009) Epigenetic transgenerational effects of endocrine disruptors on male reproduction. *Semin Reprod Med*. 27(5):403-8.
- Jablonka E, Raz G. (2009) Transgenerational epigenetic inheritance: prevalence, mechanisms, and implications for the study of heredity and evolution. *Q Rev Biol*. 84(2):131-76.
- Wang H, Chai Y, Chu X, et al. (2009) Molecular characterization of a rice mutator-phenotype derived from an incompatible cross-pollination reveals transgenerational mobilization of multiple transposable elements and extensive epigenetic instability. *BMC Plant Biol*. 29;9:63.
- Barber RC, Hardwick RJ, et al. (2009) The effects of in utero irradiation on mutation induction and transgenerational instability in mice. *Mutat Res*. 12;664(1-2):6-12.
- Katz DJ, Edwards TM, Reinke V, Kelly WG. (2009) A *C. elegans* LSD1 demethylase contributes to germline immortality by reprogramming epigenetic memory. *Cell*. 17;137(2):308-20.
- Godmann M, Lambrot R, Kimmins S. (2009) The dynamic epigenetic program in male germ cells: Its role in spermatogenesis, testis cancer, and its response to the environment. *Microsc Res Tech*. 72(8):603-19.
- Youngson NA, Whitelaw E. (2008) Transgenerational epigenetic effects. *Annu Rev Genomics Hum Genet*. 9:233-57.
- Schneider S, Kaufmann W, et al. (2008) Vinclozolin--the lack of a transgenerational effect after oral maternal exposure during organogenesis. *Reprod Toxicol*. 2008 Apr;25(3):352-60.
- Nilsson EE, Anway MD, et al. (2008) Transgenerational epigenetic effects of the endocrine disruptor vinclozolin on pregnancies and female adult onset disease. *Reproduction*. 135(5):713-21.
- Anway MD, Skinner MK. (2008) Transgenerational effects of the endocrine disruptor vinclozolin on the prostate transcriptome and adult onset disease. *Prostate*. 1;68(5):517-29.
- Anway MD, Rekow SS, Skinner MK. (2008) Transgenerational epigenetic programming of the embryonic testis transcriptome. *Genomics*. 91(1):30-40.
- Skinner MK. (2008) What is an epigenetic transgenerational phenotype? F3 or F2. *Reprod Toxicol*. 25(1):2-6.
- Crews D, Gore AC, Hsu TS, et al. (2007) Transgenerational epigenetic imprints on mate preference. *Proc Natl Acad Sci U S A*. 3;104(14):5942-6.

- Jirtle RL, Skinner MK. (2007) Environmental epigenomics and disease susceptibility. *Nat Rev Genet.* 8(4):253-62.
- Gluckman PD, Hanson MA, Beedle AS. (2007) Non-genomic transgenerational inheritance of disease risk. *Bioessays.* 29(2):145-54.
- Price TM, Murphy SK, Younglai EV. (2007) Perspectives: the possible influence of assisted reproductive technologies on transgenerational reproductive effects of environmental endocrine disruptors. *Toxicol Sci.* 96(2):218-26.
- Anway MD, Leathers C, Skinner MK. (2006) Endocrine disruptor vinclozolin induced epigenetic transgenerational adult-onset disease. *Endocrinology.* 147(12):5515-23.
- Anway MD, Memon MA, Uzumcu M, Skinner MK. (2006) Transgenerational effect of the endocrine disruptor vinclozolin on male spermatogenesis. *J Androl.* 27(6):868-79.
- Crews D, McLachlan JA. (2006) Epigenetics, evolution, endocrine disruption, health, and disease. *Endocrinology.* 147(6 Suppl):S4-10.
- Anway MD, Skinner MK. (2006) Epigenetic transgenerational actions of endocrine disruptors. *Endocrinology.* 147(6 Suppl):S43-9.
- Magnúsdóttir E, Surani MA. (2014) How to make a primordial germ cell. *Development.* 141(2):245-52.
- Colaneri A, Wang T, et al. (2013) A minimal set of tissue-specific hypomethylated CpGs constitute epigenetic signatures of developmental programming. *PLoS One.* 12;8(9):e72670.
- Skinner MK, Haque CG, Nilsson E, et al. (2013) Environmentally induced transgenerational epigenetic reprogramming of primordial germ cells and the subsequent germ line. *PLoS One.* 15;8(7):e66318.
- Wang H, Wang X, Xu X, Zwaka TP, Cooney AJ. (2013) Epigenetic Re-programming of the Germ Cell Nuclear Factor Gene is Required for Proper Differentiation of Induced Pluripotent Cells. *Stem Cells.* 2013 Mar 14. [Epub ahead of print]
- Cantone I, Fisher AG. (2013) Epigenetic programming and reprogramming during development. *Nat Struct Mol Biol.* 20(3):282-9.
- Park YJ, Herman H, et al. (2012) Sequences sufficient for programming imprinted germline DNA methylation defined. *PLoS One.* 7(3):e33024.
- Niles KM, Chan D, et al. (2011) Critical period of nonpromoter DNA methylation acquisition during prenatal male germ cell development. *PLoS One.* 6(9):e24156.
- Eguizabal C, Shovlin TC, et al. (2009) Generation of primordial germ cells from pluripotent stem cells. *Differentiation.* 78(2-3):116-23.
- Hemberger M, Dean W, Reik W. (2009) Epigenetic dynamics of stem cells and cell lineage commitment: digging Waddington's canal. *Nat Rev Mol Cell Biol.* 10(8):526-37.
- Murphey P, Yamazaki Y, et al. (2009) Epigenetic regulation of genetic integrity is reprogrammed during cloning. *Proc Natl Acad Sci U S A.* 24;106(12):4731-5.
- Shao C, Li Q, Chen S, et al. (2014) Epigenetic modification and inheritance in sexual reversal of fish. *Genome Res.* Feb 2. [Epub ahead of print].
- Sharma A. (2014) Novel transcriptome data analysis implicates circulating microRNAs in epigenetic inheritance in mammals. *Gene.* Jan 29. [Epub ahead of print].
- Wei Y, Yang CR, et al. (2014) Paternally induced transgenerational inheritance of susceptibility to diabetes in mammals. *Proc Natl Acad Sci U S A.* 4;111(5):1873-8.
- Soubry A, Hoyo C, Jirtle RL, Murphy SK. (2014) A paternal environmental legacy: Evidence for epigenetic inheritance through the male germ line. *Bioessays.* Jan 16. [Epub ahead of print].

- Zheng X, Chen L, Li M, et al. (2013) Transgenerational variations in DNA methylation induced by drought stress in two rice varieties with distinguished difference to drought resistance. *PLoS One*. 11;8(11):e80253.
- Skinner MK, Manikkam M, et al. (2013) Ancestral dichlorodiphenyltrichloroethane (DDT) exposure promotes epigenetic transgenerational inheritance of obesity. *BMC Med*. 23;11:228.
- Blewitt M, Whitelaw E. (2013) The use of mouse models to study epigenetics. *Cold Spring Harb Perspect Biol*. 2013 Nov 1;5(11):a017939. doi: 10.1101/cshperspect.a017939.
- Guérin TM, Palladino F, Robert VJ. (2013) Transgenerational functions of small RNA pathways in controlling gene expression in *C. elegans*. *Epigenetics*. 25;9(1). [Epub ahead of print]
- Berghof TV, Parmentier HK, Lammers A. (2013) Transgenerational epigenetic effects on innate immunity in broilers: an underestimated field to be explored? *Poult Sci*. 92(11):2904-13.
- Aiken CE, Ozanne SE. (2014) Transgenerational developmental programming. *Hum Reprod Update*. 20(1):63-75.
- Padmanabhan N, Jia D, et al. (2013) Mutation in folate metabolism causes epigenetic instability and transgenerational effects on development. *Cell*. 26;155(1):81-93.
- Bond DM, Baulcombe DC. (2014) Small RNAs and heritable epigenetic variation in plants. *Trends Cell Biol*. 24(2):100-7.
- Vassoler FM, Sadri-Vakili G. (2013) Mechanisms of transgenerational inheritance of addictive-like behaviors. *Neuroscience*. S0306-4522(13)00658-1. [Epub ahead of print]
- Ayala-García B, López-Santibáñez Guevara M, et al. (2013) Speciation, phenotypic variation and plasticity: what can endocrine disruptors tell us? *Int J Endocrinol*. ;2013:862739.
- Kiani J, Grandjean V, Liebers R, et al. (2013) RNA-mediated epigenetic heredity requires the cytosine methyltransferase Dnmt2. *PLoS Genet*. 9(5):e1003498.
- Feng X, Guang S. (2013) Small RNAs, RNAi and the inheritance of gene silencing in *Caenorhabditis elegans*. *J Genet Genomics*. 20;40(4):153-60.
- Suter L, Widmer A. (2013) Environmental heat and salt stress induce transgenerational phenotypic changes in *Arabidopsis thaliana*. *PLoS One* 9;8(4):e60364.
- Shiota H, Goudarzi A, Rousseaux S, Khochbin S. (2013) Transgenerational inheritance of chromatin states. *Epigenomics*. 5(2):121-2.
- Guerrero-Bosagna C, Savenkova M, et al. (2013) Environmentally induced epigenetic transgenerational inheritance of altered Sertoli cell transcriptome and epigenome: molecular etiology of male infertility. *PLoS One*. ;8(3):e59922.
- Tricker PJ, López CM, (2013) Transgenerational, dynamic methylation of stomata genes in response to low relative humidity. *Int J Mol Sci*. 26;14(4):6674-89.
- Grossniklaus U, Kelly WG, et al. (2013) Transgenerational epigenetic inheritance: how important is it? *Nat Rev Genet*. 14(3):228-35.
- Lim JP, Brunet A. (2013) Bridging the transgenerational gap with epigenetic memory. *Trends Genet*. 29(3):176-86.
- Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner MK. (2013) Plastics derived endocrine disruptors (BPA, DEHP and DBP) induce epigenetic transgenerational inheritance of obesity, reproductive disease and sperm epimutations. *PLoS One*. 8(1):e55387.
- Wan M, Gu H, Wang J, et al. (2013) Inducible mouse models illuminate parameters influencing epigenetic inheritance. *Development*. 140(4):843-52.
- Yan B, Ma L. (2012) The role for maternal piRNA in transgenerational inheritance. *Epigenomics*. 4(5):484-5.

- Sharma A. (2013) Transgenerational epigenetic inheritance: focus on soma to germline information transfer. *Prog Biophys Mol Biol.* 113(3):439-46.
- Bohacek J, Gapp K, Saab BJ, Mansuy IM. (2013) Transgenerational epigenetic effects on brain functions. *Biol Psychiatry.* 15;73(4):313-20.
- Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner MK. (2012) Dioxin (TCDD) induces epigenetic transgenerational inheritance of adult onset disease and sperm epimutations. *PLoS One.* 7(9):e46249.
- Skinner MK, Mohan M, et al. (2012) Epigenetic transgenerational inheritance of somatic transcriptomes and epigenetic control regions. *Genome Biol.* 3;13(10):R91.
- Ou X, Zhang Y, Xu C, et al. (2012) Transgenerational inheritance of modified DNA methylation patterns and enhanced tolerance induced by heavy metal stress in rice (*Oryza sativa* L.). *PLoS One.* 2012;7(9):e41143.
- Manikkam M, Tracey R, et al. (2012) Pesticide and insect repellent mixture (permethrin and DEET) induces epigenetic transgenerational inheritance of disease and sperm epimutations. *Reprod Toxicol.* 34(4):708-19.
- Becker C, Weigel D. (2012) Epigenetic variation: origin and transgenerational inheritance. *Curr Opin Plant Biol.* 15(5):562-7.
- Ashe A, Sapetschnig A, et al. (2012) piRNAs can trigger a multigenerational epigenetic memory in the germline of *C. elegans*. *Cell.* 6;150(1):88-99.
- Crews D, Gillette R, et al (2012) Epigenetic transgenerational inheritance of altered stress responses. *Proc Natl Acad Sci U S A.* 5;109(23):9143-8.
- Kovalchuk I. (2012) Transgenerational epigenetic inheritance in animals. *Front Genet.* 9;3:76.
- Nilsson E, Larsen G, Manikkam M, et al. (2012) Environmentally induced epigenetic transgenerational inheritance of ovarian disease. *PLoS One.* 7(5):e36129.
- Grentzinger T, Armenise C, et al. (2012) piRNA-mediated transgenerational inheritance of an acquired trait. *Genome Res.* 22(10):1877-88.
- Goerlich VC, Nätt D, et al. (2012) Transgenerational effects of early experience on behavioral, hormonal and gene expression responses to acute stress in the precocial chicken. *Horm Behav.* 61(5):711-8.
- Braunschweig M, Jagannathan V, et al. (2012) Investigations on transgenerational epigenetic response down the male line in F2 pigs. *PLoS One.* 7(2):e30583.
- Daxinger L, Whitelaw E. (2012) Understanding transgenerational epigenetic inheritance via the gametes in mammals. *Nat Rev Genet.* 31;13(3):153-62.
- Berger SL. (2012) Transgenerational inheritance of longevity: epigenetic mysteries abound. *Cell Metab.* 4;15(1):6-7.
- Benayoun BA, Brunet A. (2012) Epigenetic memory of longevity in *Caenorhabditis elegans*. *Worm.* 1;1(1):77-81.
- Nelson VR, Nadeau JH. (2010) Transgenerational genetic effects. *Epigenomics.* ;2(6):797-806.
- Guerrero-Bosagna C, Skinner MK. (2012) Environmentally induced epigenetic transgenerational inheritance of phenotype and disease. *Mol Cell Endocrinol.* 6;354(1-2):3-8.
- Greer EL, Maures TJ, et al. (2011) Transgenerational epigenetic inheritance of longevity in *Caenorhabditis elegans*. *Nature.* 19;479(7373):365-71.
- Skinner MK. (2011) Environmental epigenetic transgenerational inheritance and somatic epigenetic mitotic stability. *Epigenetics.* 6(7):838-42.
- Hauser MT, Aufsatz W, Jonak C, Luschnig C. (2011) Transgenerational epigenetic inheritance in plants. *Biochim Biophys Acta.* 1809(8):459-68.

- Skinner MK. (2011) Role of epigenetics in developmental biology and transgenerational inheritance. *Birth Defects Res C Embryo Today*. 93(1):51-5.
- Paszkowski J, Grossniklaus U. (2011) Selected aspects of transgenerational epigenetic inheritance and resetting in plants. *Curr Opin Plant Biol*. 14(2):195-203.
- Matthews SG, Phillips DI. (2012) Transgenerational inheritance of stress pathology. *Exp Neurol*. 233(1):95-101.
- Walker DM, Gore AC. (2011) Transgenerational neuroendocrine disruption of reproduction. *Nat Rev Endocrinol*. 7(4):197-207.
- Carone BR, Fauquier L, et al. (2010) Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell*. 143(7):1084-96.
- Pembrey ME. (2010) Male-line transgenerational responses in humans. *Hum Fertil (Camb)*. 13(4):268-71.
- Arai JA, Feig LA. (2011) Long-lasting and transgenerational effects of an environmental enrichment on memory formation. *Brain Res Bull*. 25;85(1-2):30-5.
- Daxinger L, Whitelaw E. (2010) Transgenerational epigenetic inheritance: more questions than answers. *Genome Res*. 20(12):1623-8.
- Pentinat T, Ramon-Krauel M, et al. (2010) Transgenerational inheritance of glucose intolerance in a mouse model of neonatal overnutrition. *Endocrinology*. 151(12):5617-23.
- Guerrero-Bosagna C, Settles M, Lucker B, Skinner MK. (2010) Epigenetic transgenerational actions of vinclozolin on promoter regions of the sperm epigenome. *PLoS One*. 30;5(9). pii: e13100.
- Skinner MK, Guerrero-Bosagna C. (2009) Environmental signals and transgenerational epigenetics. *Epigenomics*. 1(1):111-117.
- Dunn GA, Morgan CP, Bale TL. (2011) Sex-specificity in transgenerational epigenetic programming. *Horm Behav*. 59(3):290-5.
- Sharma A, Singh P. (2009) Detection of transgenerational spermatogenic inheritance of adult male acquired CNS gene expression characteristics using a *Drosophila* systems model. *PLoS One*. 2;4(6):e5763.
- Slatkin M. (2009) Epigenetic inheritance and the missing heritability problem. *Genetics*. 182(3):845-50.

REVIEW ARTICLE

Environmentally induced epigenetic transgenerational inheritance of disease

Eric E. Nilsson, Ingrid Sadler-Riggleman and Michael K. Skinner*

Center for Reproductive Biology, School of Biological Sciences, Washington State University, Pullman, WA 99164-4236, USA

*Correspondence address. Center for Reproductive Biology, School of Biological Sciences, Washington State University, Pullman, WA 99164-4236, USA.
Tel: +1-509-335-1524; Fax: +1-509-335-2176; E-mail: skinner@wsu.edu
Managing Editor: Isabelle Mansuy

Abstract

Ancestral environmental exposures such as toxicants, abnormal nutrition or stress can promote the epigenetic transgenerational inheritance of disease and phenotypic variation. These environmental factors induce the epigenetic reprogramming of the germline (sperm and egg). The germline epimutations can in turn increase disease susceptibility of subsequent generations of the exposed ancestors. A variety of environmental factors, species and exposure specificity of this induced epigenetic transgenerational inheritance of disease is discussed with a consideration of generational toxicology. The molecular mechanisms and processes involved in the ability of these inherited epimutations to increase disease susceptibility are discussed. In addition to altered disease susceptibility, the potential impact of the epigenetic inheritance on phenotypic variation and evolution is considered. Observations suggest environmentally induced epigenetic transgenerational inheritance of disease is a critical aspect of disease etiology, toxicology and evolution that needs to be considered.

Key words: epigenetics; transgenerational; non-genetic inheritance; disease etiology; evolution; review

Introduction

The term epigenetics was originally coined by C.H. Waddington in the 1940s in relation to his studies of gene–environment interactions involving non-Mendelian inherited phenomena [1, 2]. More recent molecular oriented definitions are that epigenetics refers to ‘the molecular factors and processes around the DNA that regulate genome activity independent of DNA sequence, and that are mitotically stable’ [3] (Table 1). These molecular factors include DNA methylation [4], histone modifications [5], non-coding RNAs [6, 7], chromatin structure [8], and RNA methylation [9] (Fig. 1). The complex integration of epigenetic modifications is referred to as the ‘epigenome’. The first whole epigenome analysis was accomplished in 2005, mapping

histone acetylation and methylation in yeast [10]. Epigenetic processes are critical for allowing an organism to respond to its environment with changes in gene expression. In addition, epigenetic mechanisms allow a stem cell type to develop into a differentiated cell type [3, 11, 12] (Fig. 2). Therefore, epigenetic processes are an integral part of normal biology.

Molecular Epigenetic Mechanisms

There are a variety of epigenetic factors that act around the DNA in a cell to regulate gene expression and genome activity. DNA methylation is the most extensively studied epigenetic factor. DNA methylation involves a small (methyl) chemical


Received 8 February 2018; revised 1 May 2018; accepted 15 May 2018

© The Author(s) 2018. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Table 1: glossary terms and definitions

Glossary term	Definition
Epigenetics	Molecular factors and processes around DNA that regulate genome activity independent of DNA sequence, and are mitotically stable
Epigenetic transgenerational inheritance	Germline mediated inheritance of epigenetic information between generations in the absence of continued direct environmental influences
Multigenerational	Direct exposure of multiple generations
Epimutation	Environmentally induced differential presence of epigenetic alterations that can lead to altered genome activity when compared to organisms not having the exposure



EPIGENETIC MECHANISMS AND MARKS

- DNA Methylation
- Histone Modifications
- Chromatin Structure
- Non-coding RNA
- RNA methylation

Figure 1: epigenetic mechanisms and processes (marks). Modified from [122]

group being attached to DNA, primarily at the cytosine base when it is adjacent to a guanine residue [4, 13] to produce 5-methylcytosine (5mC). Other chemical modifications of cytosine bases in DNA have also been described. The TET (ten-eleven translocation) family of enzymes can oxidize 5mC to 5-hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC), and 5-carboxylcytosine (5caC) [14]. In broad terms, the presence of 5mC often represses DNA transcription, while 5hmC is permissive to transcription [15, 16]. However, one of the main functions appears in the DNA methylation erasure during early development [17]. N(6)-methyladenine is an epigenetic modification to the adenine base of DNA that was once thought to only be present in prokaryotic organisms, but has now been described in mammalian embryonic stem cells [18].

The histone proteins that DNA is wrapped around create the nucleosome and can be chemically modified to alter gene expression. There are many different histone post-translational modifications including lysine acetylation, lysine and arginine methylation, arginine citrullination, lysine ubiquitination, lysine sumoylation, ADP-ribosylation, proline isomerization, and serine/threonine/tyrosine phosphorylation [19]. These modifications can change chromatin structure or recruit transcriptional cofactors to DNA in order to regulate gene expression.

Alternatively, they can act as repressive marks to reduce gene expression in major regions of the genome such as heterochromatin. In broad terms, histone acetylation can increase transcription, while methylation can be repressive to transcription.

Non-coding RNA molecules can act as epigenetic factors [20]. These are small and long RNA molecules that do not code for a protein, but rather function as RNA to regulate gene expression. The non-coding RNA molecules that act as epigenetic factors are not DNA sequence dependent, so the majority do not rely on having a nucleotide sequence that is complementary to a specific DNA or RNA region in order to function. Long non-coding RNAs (lncRNAs) [21] and transfer RNA-derived small RNAs (tsRNAs) [22] are examples of RNA classes that are present in sperm and can act as epigenetic factors that affect subsequent generations [23].

RNA molecules can themselves be epigenetically modified and so affect translation and gene expression [24]. The most prevalent reversible modification to the internal sequence of mRNA is methylation of adenosine to form N(6)-methyladenosine (m(6)A). m(6)A mRNA methylation is associated with post-transcriptional regulation [25, 26]. Cytosine methylation (m3C) in both mRNA and tRNA also occurs [27, 28]. Methylation of tRNA inhibits processing of tRNA into tsRNA halves, which

EPIGENETIC AND GENETIC CASCADE OF EVENTS INVOLVED IN DEVELOPMENT




Figure 2: epigenetic and genetic cascade of events involved in development. Cascade of genetic and epigenetic stages interacting to promote differentiated cells. The critical window of exposure allows environmental factors to alter the epigenetic cascade to obtain a modified differentiated site and to cause altered transcriptomes to increase disease susceptibility and phenotypic variation. Modified from [3]

themselves affect transcription [22, 29, 30]. Therefore, RNA methylation is the most recent epigenetic molecular factor identified.

The coiling, looping, and general structure of DNA is termed chromatin structure and is also an epigenetic factor [8]. The three-dimensional structure of DNA can make certain regions of the genome accessible to transcription machinery, such as being enhancer regions near gene promoters to affect gene expression. Therefore, epigenetic molecular processes include DNA methylation, histone modifications, non-coding RNAs, RNA methylation, and chromatin structure.

Epigenetic Transgenerational Inheritance

The definition of epigenetic transgenerational inheritance is 'germline-mediated inheritance of epigenetic information between generations in the absence of continued direct environmental influences that leads to phenotypic variation' [3, 31] (Table 1). Multigenerational exposures, in contrast, refer to observed effects in subsequent generations that are the result of direct exposure [31] (Table 1; Fig. 3). Direct environmental exposure of the parents, considered to be the F0 generation, can also affect the germline (sperm or eggs) of either parent. Therefore, the next generation (F1) derived from this germline is still considered exposed, and so is not truly transgenerational. For pre-conception parental exposures the F2 generation offspring is considered the first transgenerational unexposed generation (Fig. 3). The situation is different when a gestating female is exposed, because then the fetus and the fetus' germline are directly exposed as well. In that case, the F3 generation is the first unexposed transgenerational offspring [31] (Fig. 3).

The Agouti mouse model is an example of multigenerational inheritance [32–34]. When pregnant Agouti mice are exposed to a methyl donor in their diet, they experience increased methylation on an allele of their Agouti gene, which leads to a coat color change in their offspring. Generally, this change is not

passed on to future generations. Instead the normal process of epigenetic reprogramming in the germline and early embryo returns the DNA methylation state to its original setting.

An increasing number of examples of transgenerational inheritance of disease are present in the literature (Table 2). Some of the first experiments to establish the potential for epigenetic transgenerational inheritance were performed by Conrad Waddington, who coined the term 'epigenetic' [1, 35]. In these studies, it was found that a heat shock induced wing structure change in *Drosophila melanogaster* persisted for more than seven generations [35]. An even earlier study in Guinea pigs demonstrated transgenerational inheritance of decreased fertility and increased mortality for four generations after ancestral exposure to ethanol vapor, although this was not attributed to epigenetic inheritance at the time [36]. One of the first studies to associate molecular epigenetic changes with transgenerational inheritance of disease in mammals was an investigation of the effects of treating pregnant rats with the agricultural fungicide vinclozolin [37]. The F3 generation (great-grand offspring) demonstrated reproductive abnormalities such as increased testicular germ cell apoptosis and decreased sperm motility. These transgenerational phenotypes were correlated with changes in DNA methylation in the F3 generation sperm [37].

Several environmental toxicants including vinclozolin, DDT (dichlorodiphenyltrichloroethane), methoxychlor, plastic derived compounds, hydrocarbons, atrazine, tributyltin have been shown to promote the transgenerational inheritance of increased disease susceptibility in rodent models [38, 39] (Table 2). The diseases that were increased transgenerationally included testis, prostate and kidney disease, obesity, polycystic ovaries, reduced oocyte number in the ovaries, and cancer [39]. For the purposes of this review, more recently published investigations of epigenetic transgenerational inheritance of disease will be highlighted, (Table 2). Exposure of mice to the phthalate plastic derived compound DEHP (di(2-ethylhexyl) phthalate) has been shown to result in transgenerational changes to stress




Figure 3: environmentally induced transgenerational epigenetic inheritance. Schematic of multigenerational versus transgenerational environmental exposures. Modified from [31]

Table 2: examples of transgenerational inheritance from specific exposures and specific effects

Exposure	Effects	Reference
Environmental toxicants		
Vinclozolin	Impaired male fertility; prostate, kidney disease, tumors, immune and reproductive pathologies	[37, 78, 94]
Vinclozolin	Gender-specific changes in anxiety-like behavior	[85]
Methoxychlor	Impaired male fertility; kidney disease, ovary disease, and obesity	[37, 86]
Permethrin/DEET	Prostate, kidney disease	[81]
Dioxin	Prostate, kidney disease, reduced fertility, negative effects on pregnancy outcome	[80, 123]
BPA/phthalates	Prostate, kidney disease; obesity	[43]
Hydrocarbon mixture (jet fuel)	Prostate, kidney disease; obesity; immune and reproductive pathologies	[46]
Vinclozolin, permethrin/DEET, plastics, dioxin, jet fuel	Polycystic ovaries, reduced primordial follicle pool	[82]
DDT	Obesity	[45]
Phthalate	Disruption of testicular germ cell organization and spermatogonial stem cell function, changes in hormones and behavior	[40, 124]
Phthalate	Disrupted ovarian function	[41]
Tributyltin	Increase in fat depot size	[38]
BPA	Cardiac disease; reduced fertility	[48, 72]
BPA	Changes in social behavior and neural gene expression	[42]
Atrazine	Testicular disease, early puberty, lean phenotype	[125]
Benzo[a]pyrene	Behavioral and physiological deficits	[50]
Mercury	Behavior change	[49]
Other exposures		
Caloric restriction	Cardiovascular mortality	[56, 77]
High-fat diet	Increased body size; reduced insulin sensitivity, increased mammary cancer	[57–59]
Folate	Congenital malformations	[126]
Stress	Reduced social interaction; increased stress resilience; disrupted neural connectivity; physiology changes; increased anxiety	[51–55]
Drought	DNA methylation changes	[127]
Heat/salt stress	Accelerated flowering, increased salt tolerance	[128]
Prediabetes/diabetes	Impaired glucose tolerance; reduced insulin sensitivity, male subfertility	[61, 62]
Smoking	Abnormal pulmonary function	[129]
Ethanol	Neurological defects; decreased fertility	[36, 47, 130]
Heat stress	Increased Hsp70 production and tolerance to heat stress; wing structure changes	[131, 132]

hormones, behavior [40], and ovarian function [41]. Earlier studies in mice [42] showed that ancestral exposure to the plastic derived compound bisphenol A (BPA) caused changes in social behavior in juvenile mice and changes in expression of neural

genes such as oxytocin and vasopressin. Earlier studies in rats have shown that exposure to a mixture of BPA and phthalates induces transgenerational increases in pubertal abnormalities, testis disease, and ovarian disease [43]. Ancestral exposure of

ENVIRONMENTALLY INDUCED EPIGENETIC TRANSGENERATIONAL INHERITANCE

Environmental Toxicants

Vinclozolin (Agricultural Fungicide)
Methoxychlor (Agricultural Pesticide)
Dioxin/TCDD (Industrial Contaminant)
Plastic Compounds (BPA & Phthalates)
Mercury

Permethrin & DEET (Insect Repellants)
DDT (Pesticide)
Tributyltin (Industrial Toxicant & Biocide)
Hydrocarbons (Jet Fuel)
Atrazine

Other Types Exposures

Nutrition (High Fat or Caloric Restriction)

Smoking & Alcohol

Temperature & Drought (Plant Health & Flowering)

Stress (Behavioral)



Plants



Flies



Worms



Fish



Bird



Rodents



Pigs



Humans

Figure 4: environmentally induced epigenetic transgenerational inheritance. Various exposures and species investigated

mice to the toxicant tributyltin results in a transgenerational increase in obesity [38, 44]. Earlier investigations in pregnant rat exposures to DDT, jet fuel hydrocarbons, or a BPA/phthalates mix will also increase obesity transgenerationally [43, 45, 46]. Other recently published investigations indicate that ethanol exposure of pregnant mice can cause transgenerational neurological changes in descendants that resemble those of Fetal Alcohol Spectrum Disorders [47]. In zebrafish, BPA exposure of males can result in the transgenerational inheritance of heart disorders in the F2 generation [48]. Zebrafish exposure to mercury [49] or to the industrial pollutant benzopyrene [50] induces the transgenerational inheritance of abnormal neurobehaviors that are correlated with epigenetic changes (i.e. epimutations) in sperm (Table 2) [49, 50].

Exposure to environmental factors other than toxicants can also induce transgenerational inheritance (Table 2). The stress of maternal separation in mice transgenerationally disrupts functional connectivity throughout the brain [51], as well as both impairing social interactions and cognition and making the descendant mice more stress resilient [52]. Mice subjected to restraint stress also transmitted reduced anxiety levels to their transgenerational descendants [53]. Conversely, social hierarchy stress in mice was shown to increase anxiety behaviors transgenerationally [54]. This raises the possibility that several psychological stressors can induce different transgenerational effects. In pregnant rats, the stressors of forced swim and restraint induce transgenerational inheritance of physiological changes such as alterations in catecholamine biosynthesis and immune response [55].

Other examples of transgenerational inheritance have been observed with caloric restriction or high fat diets. The Överkalix study by Bygren *et al.* [56] shows how cardiovascular mortality in humans can be influenced by reduced childhood and adolescent food supply. Effects were shown to reach into the second generation. Maternal high fat diet in mice can increase body size and reduce insulin sensitivity in F3 generation female offspring [57], although Masuyama *et al.* [58] demonstrated that a normal diet in utero for three subsequent generations can return glucose and lipid metabolism to normal. In addition,

a maternal high fat diet in mice can transgenerationally increase mammary cancer risk [59]. Previous studies with rats demonstrated that exposure of pregnant animals to the environmental toxicant vinclozolin also promoted a transgenerational increase in tumors [60]. Interestingly, diabetes in mice can induce transgenerational inheritance of male subfertility [61]. A paternal prediabetic condition in mice can be inherited transgenerationally as shown by impaired glucose tolerance and decreased insulin sensitivity [62]. Similarly, male rats fed a high fat diet promoted transgenerational inheritance of impaired glucose tolerance in F2 generation offspring [63].

Species Diversity of Epigenetic Transgenerational Inheritance

Epigenetic transgenerational inheritance has been identified to occur in a wide variety of organisms (Fig. 4). This review focuses on examples of epigenetic transgenerational inheritance of disease or abnormalities in different animal species. A number of studies have demonstrated the environment (e.g. heat and drought) can induce the epigenetic transgenerational inheritance of phenotypic variation in plants [64]. In the nematode worm *Caenorhabditis elegans* increased longevity that is associated with the histone modification H3K4me3 methylation can be transgenerationally inherited for up to three generations [65]. As mentioned previously, Waddington performed early experiments using the model insect species *D. melanogaster* and demonstrated that a heat shock induced wing structure changes that persisted for more than seven generations [1, 35] and now for hundreds of generations in today's stocks. In more recent examples, it has been found that a high-sugar maternal fly diet can alter the larval body composition for the next two generations [66]. Similarly, a high fat larval diet in fruit flies can cause transgenerational alterations to F2 generation pupal and egg size [67]. Manipulations of the protein levels in the diet of fruit flies can affect longevity and reproduction for three subsequent generations, and this effect is associated with histone

modifications [68, 69]. In another arthropod species, the crustacean *Daphnia magna*, exposure to the toxicant 5-azacytidine results in decreased body length and reduced levels of DNA methylation in non-exposed subsequent generations [70].

Several species of fish have shown epigenetic transgenerational inheritance of disease. Zebrafish exposed to the environmental toxicants benzo(a)pyrene [50], methylmercury [49] or dioxin [71] transmit to their grand-offspring behavioral changes, visual defects, increased body mass, skeletal abnormalities and/or decreased fertility, sometimes associated with changes in DNA methylation. Medaka exposed to the endocrine disruptors BPA or ethinylestradiol produce grand-offspring and great-grand-offspring with reduced fertility [72].

Some bird species have shown evidence of epigenetic transgenerational inheritance. In a study with quail eggs exposed to the environmental estrogen genistein [73] the great-grand offspring age at which the first egg was laid was significantly greater. In ducks, feeding a methionine-deficient diet produces grand-offspring with altered weight gain and changes in metabolic parameters [74].

In mammals most studies of epigenetic transgenerational inheritance have occurred in rodents [75]. Another experimental mammal involves pigs and abnormal nutritional induced epigenetic transgenerational inheritance [76]. Examples of transgenerational inheritance of increased susceptibility to diseases have been outlined above for rats, mice and Guinea pigs [36, 37, 41, 44, 45]. Evidence of epigenetic transgenerational inheritance of disease in humans comes from retrospective studies such as those including the Dutch and Swedish famines [56, 77]. As previously mentioned, the descendants of people exposed to famine conditions as children 9–12 years of age in Sweden were investigated and it was found that men whose grandfathers had been exposed to famine had an increased risk of mortality due to diabetes, and similarly women whose grandmothers were exposed had increased risk [31]. Due to the conservation of environmentally induced epigenetic transgenerational inheritance from plants to humans all organisms will utilize epigenetic inheritance to facilitate environmental adaptation and response.

Phenotypic Diversity of Transgenerationally Inherited Diseases

Studies of the effects of ancestral exposure to an array of toxicants (Table 2) demonstrate epigenetic transgenerational inheritance of a variety of diseases and abnormalities, including testis disease [37], prostate and kidney disease [43, 46, 78–82], mammary tumors [78], immune and reproductive pathologies [46, 78, 83, 84], obesity [45, 46], behavioral effects [85] and many others listed in Table 2. The disease phenotypes observed in these experiments often depend on the specific exposure of the F0 generation. For example, increased obesity risk in rats is inherited transgenerationally after ancestral exposure to DDT, plastic compounds, hydrocarbons and methoxychlor [43, 45, 86], but not dioxins. Jet fuel hydrocarbons induce an elevated rate of luteal ovarian cyst formation in F3 females [46, 82], a phenotype not observed with other exposures. On the other hand, some ovarian disorders such as polycystic ovaries and reduction of the primordial follicle pool size have been shown to be inherited transgenerationally after exposure of the F0 generation to many of the toxicants studied [84, 87]. The explanation for this phenomenon may be that some developmental processes, in this case ovarian follicle development, are more

sensitive to epigenetic and gene expression changes in their developmental regulatory networks, and so will be more easily affected than those of other cells and tissues (Fig. 2).

Epigenetic processes are major mechanisms by which organisms respond and adapt to their environment. Therefore, how can environmental epigenetic insults result in transgenerational inheritance of increased disease susceptibility? Since this is a maladaptive response one possible answer may be seen in the predictive adaptive response hypothesis [88]. In this hypothesis an environmental stressor like famine may epigenetically promote an adaptive (thrifty) phenotype in subsequent generations. If the current environment of those descendants has more than adequate nutrients, diseases such as diabetes and obesity are promoted. Another possibility is that an environmental insult, such as exposure to a toxicant, may interfere with the normal molecular epigenetic machinery and result in stochastic and/or directed epigenetic changes that could be considered epimutations. The term epimutation is defined as ‘the environmentally induced differential presence of epigenetic alterations that can lead to altered genome activity, when compared to organisms not having exposure’ (Table 1). If these epimutations occur in germ cells they can lead to transgenerational inheritance of a wider range of phenotypes in the progeny. Some of those phenotypes may be poorly adapted and develop disease. This would explain an increase in disease susceptibility in organisms whose ancestors were exposed to environmental insults. However, phenotypic variation is the ‘raw material’ upon which natural selection acts. Therefore, the increased phenotypic variation may also result in some individuals who are better adapted to an altered environment which can facilitate natural selection and evolution [89].

Developmental Etiology of Epigenetic Transgenerational Inheritance

A number of reproductive processes involve DNA methylation changes that normally will be reset by genome-wide DNA methylation reprogramming events. The two main developmental periods are in the early embryo after fertilization and during germ cell specification at gonadal sex determination [90, 91] (Fig. 5). This phenomenon allows embryonic stem cells to develop by removing epigenetic constraints to promote pluripotency. Some parental epigenetic changes, such as imprinted genes, are protected from being reprogrammed during these developmental periods. In contrast, some parent specific imprints are established during this epigenetic reprogramming [92]. Environmentally induced DNA methylation alterations called differential DNA methylation regions (DMRs) [93] present in germ cells behave as imprinted-like genes in the way their methylation patterns persist. By definition, true imprinted genes display ‘parent-of-origin allelic transmission with monoallelic gene expression’. DMRs often demonstrate parent-of-origin allelic transmission, but monoallelic gene expression has not been demonstrated. Differentially methylated sites connected with transgenerational inheritance are called ‘imprinted-like’ [94]. The transmission of epigenetic information to future generations via germ cells can alter the epigenome of the developing embryonic stem cells which would be expected to promote changes to the epigenetic and transcriptional programming of all derived somatic cells [95]. Those tissues that are sensitive to alterations in their epigenomes and transcriptomes may show increased susceptibility and prevalence of disease development [93, 96] (Fig. 2). Normal biology




Figure 5: epigenetic reprogramming during primordial germ cell development at gonadal sex determination and following fertilization in the early embryo. Modified from [94]

requires alterations of epigenetics for the development of stem cell populations and subsequent somatic cell differentiation. The epigenetic transgenerational inheritance molecular process is directly linked to these epigenetic reprogramming processes in the germline and the developing embryo.

Germline Epimutations

It is a prerequisite for environmentally induced epigenetic transgenerational inheritance that there be epigenetic changes (i.e. epimutations) in the germline, because the germ cells (sperm and egg) are the only cells that can transmit molecular information between generations from the parents to their offspring. Early studies investigating transgenerational epimutations in germ cells used pregnant rats exposed to vinclozolin during the period of gonadal sex determination when epigenetic reprogramming of the fetal germ cells occurs. A genome-wide promoter analysis was applied to look for epigenetic changes in the sperm DNA and approximately 50 differential DMRs were identified in gene promoters in vinclozolin lineage F3 generation sperm DNA versus control lineage [97]. Similar experiments have been performed in rats using a number of additional toxicants including dioxin [80], a mixture of permethrin and DEET (*N,N*-diethyl-meta-toluamide) [81], BPA and phthalates [43] and jet fuel (hydrocarbons) [46]. All these toxicants were found to promote transgenerational inheritance of both disease and sperm DMRs. Interestingly, it was observed that each toxicant produced an exposure-specific set of DNA methylation changes in the sperm, and comparisons between the different toxicant exposures demonstrated negligible overlap between them [82]. This raises the possibility that these 'epigenetic signatures' may be used in the future as a diagnostic tool to determine if an individual has had a particular environmental toxicant exposure in their ancestry. The examination of the genomic features of all these DMRs identified a low CpG density termed CpG deserts [98] and a DNA sequence motif called Environmental Induced Differential Methylation Consensus Sequence 1 (EDM1). Nearly all the DMRs identified with numerous exposures had these genomic features [39]. A machine

learning analysis used this data to identify approximately 40 000 potential genome-wide DMR sites susceptible to environmental alterations [99]. Further studies are needed to determine the utility of these potential epimutation sites as biomarkers for exposure and disease.

Comparisons have been made of the DMRs induced in the direct exposure F1 generation and transgenerational F3 generation vinclozolin lineage male sperm [100]. As described above, when the gestating female is directly exposed to a toxicant the F1 generation fetus is also directly exposed, as are the developing germ cells within the F1 generation fetus that will generate the F2 generation. The F3 generation animals are the first non-exposed transgenerational descendants (Fig. 3). Therefore, the molecular mechanisms of inducing epigenetic changes is different in the direct exposure F1 generation, and in the F1 generation germ cells (sperm) that will produce the F2 generation, when compared with mechanisms by which epimutations are induced in the transgenerational F3 generation. In a study involving vinclozolin exposure of gestating female rats there was a distinct set of DNA methylation changes in the F1 generation sperm that was different from the set of methylation changes in the transgenerational F3 generation sperm [100]. This suggests that the direct exposure induced F1 generation sperm epimutations promote epigenetic alterations during germ cell development in subsequent generations that lead to the different DMRs in the F3 generation. This mechanism appears to be associated with altered early embryonic development of the stem cells.

In addition to DNA methylation, other epigenetic factors such as non-coding RNA (ncRNA) can also contribute to epigenetic transgenerational inheritance. Small ncRNAs of the microRNA class are altered in the sperm of stressed vs. unstressed mice and have been shown experimentally to promote a change in the hypothalamic-pituitary-adrenal stress axis reactivity of offspring [101]. Another class of small non-coding RNAs associated with transgenerational sperm are 5' halves of tRNAs [102]. These tRNA 5' halves and microRNAs are transgenerationally altered in the F3 generation sperm of rats ancestrally exposed to vinclozolin during pregnancy [102]. A number

of studies have demonstrated the potential role of ncRNA in epigenetic transgenerational inheritance [103].

Another epigenetic factor present in sperm and associated with transgenerational inheritance is the retention of histone proteins [104]. During spermatogenesis in vertebrates the histone cores that DNA is wrapped around in most somatic cell types are replaced by protamines, allowing for more tightly compacted DNA in sperm heads [105]. However, 1–10% of histones are retained in mammals, depending on species [106]. These retained sperm histones have been implicated in regulating gene expression in the resulting offspring [107]. In a recent transgenerational study using rats, Ben Mammar *et al.* [108] demonstrated that a specific set of histones are retained in F3 generation control lineage sperm. This same set of histones is retained in F3 generation rats ancestrally exposed to vinclozolin or DDT, but additional sites of histone retention are induced in the vinclozolin and DDT lineage sperm [108]. Therefore, histone retention also appears to be associated with sperm mediated transgenerational inheritance of disease following ancestral DDT or vinclozolin exposure [104, 108].

Since post-translational modifications of histones are known to be an epigenetic factor that regulates gene expression studies have investigated histone modifications present in sperm. Histone H3 methylation changes in retained sperm histones have been correlated with fertility in humans [109] and with survival of offspring in mice [110]. Histone modifications have been correlated with epigenetic transgenerational inheritance of altered phenotypes in *C. elegans* [111], *Drosophila* [112], and recently in mammals [104].

Previous transgenerational studies have focused on epigenetic factors and epimutations in sperm due to the relative ease of obtaining large numbers of sperm cells. Several studies have shown that epigenetic transgenerational inheritance is mediated through the female germline [45, 86]. Eggs cannot be obtained in large enough quantity to allow traditional molecular analysis. Future studies with single cell analyses may be needed to document the role of epimutations in eggs. Epigenetic factors in eggs appear to play an equally important role in epigenetic inheritance, but remain to be investigated. The epigenetic transgenerational inheritance of disease following environmental exposures will likely be mediated by the integrated actions and combination of different epigenetic factors present in gametes. A recent study in rats demonstrated that after treatment of gestating females with DDT or vinclozolin there were concurrent transgenerational alterations in F3 generation sperm in DNA methylation, histone retention, and non-coding RNAs [108, 113]. Therefore, transgenerational alterations in all the different epigenetic processes appear to be involved in the epigenetic transgenerational inheritance phenomenon.

Transgenerational Gene Expression Changes

Transgenerational inheritance of environmentally induced epigenetic changes requires transmission through the germ line from parents to future generations. However, epigenetic changes themselves would not cause disease, rather they must manifest as changes in gene expression. Ensuing disease such as cancer, prostate or kidney abnormalities, and obesity are brought on by disturbances in gene expression in the pertinent somatic cells. The hypothesis is that the epimutations in the germline alter the epigenome of the embryonic stem cells that then affect all subsequent somatic cell epigenomes and transcriptomes [87, 95] (Fig. 2). These cell and tissue specific epimutations promote tissue specific alterations in transcriptomes

[96]. These aberrant transcriptomes could then lead to a susceptibility for physiological abnormalities and disease (Fig. 2).

Exposure of F0 generation animals to environmental toxicants will affect and change the transcriptomes of potentially all tissues in future generations [96]. In a study of rats ancestrally exposed to vinclozolin the transcriptomes of 11 different tissue types from adult male and female animals were examined [96]. It was found that there were gene expression differences between control and vinclozolin lineage animals in the different tissues with minimal overlap in the differentially expressed genes between tissues. However, there was significant overlap in the physiological pathways and cellular processes that were affected by gene expression changes in different tissues. For example, both prostate and liver tissues were enriched for genes in transcription and focal adhesion processes, but the specific genes altered were not the same in each tissue [96]. These observations warranted a closer look at the genomic locations of epimutations and differentially expressed genes. Looking across the different tissue types it was found that there were regions of the genome that had statistically over-represented clusters of gene expression changes [96]. These regions in the genome were called epigenetic control regions (ECR). These ECR are 2–5 megabase in size and have clusters of genes. Within these ECRs are DNA methylation epimutations and long non-coding RNA (ncRNA) expression sites [114]. The long ncRNAs play a role in regulation of distal gene transcription and epigenetic regulation [115, 116]. Observations suggest that within an ECR many of the genes are epigenetically regulated up or down as a block. Therefore, in one cell type those genes within the ECR normally expressed would be regulated while in another cell type a different set of genes within the ECR normally expressed would be affected. Epigenetic alterations within the ECR can influence gene expression in a variety of cell types differently [96]. Interestingly, the location of ECRs has been shown to co-localize with clusters of transgenerational epimutations (e.g. DMRs) found after ancestral toxicant exposures [117].

Several studies have suggested how the molecular mechanisms of environmentally induced transgenerational inheritance may lead to tissue specific disease occurrence. As mentioned earlier, two ovarian disorders, polycystic ovarian syndrome, and primary ovarian insufficiency (premature reduction of the primary follicle pool) were both induced transgenerationally by a number of environmental toxicants [83]. Analysis of this phenomenon involved the isolation of a specific cell type from the tissue that is associated with the disease in the vinclozolin lineage animals. The granulosa cells were isolated from the ovarian follicles of young female rats prior to disease onset. The epigenomes and transcriptomes of these granulosa cells from control and vinclozolin lineages were analyzed and compared [87]. Granulosa cells from F3 generation vinclozolin lineage rats had differences in both the epigenome and the transcriptome compared with the control lineage. Interestingly, some of the affected genes had been previously shown to be associated with polycystic ovarian syndrome and primary ovarian insufficiency [87]. Similar results were obtained when the molecular basis of transgenerational male infertility in rats was examined. As above, changes in the epigenome and transcriptome were found in testicular Sertoli cells of F3 generation rats after ancestral vinclozolin exposure [95]. Several of the differentially regulated genes identified were known to be associated with male infertility, such as HDAC1 and HSP90AA1 [118, 119]. In addition, a number of Sertoli cell genes associated with pyruvate production were down-regulated and this is

known to impact spermatogenic cell survival and promote germ cell apoptosis, which is one of the testis pathology phenotypes observed [95]. Therefore, the environmentally induced transgenerational changes in the somatic cell epigenomes are associated with transgenerational changes in gene expression, which are related to the increases in disease development observed.

Experimental and Technical Approach Limitations

One of the main experimental design issues and limitations is a consideration of what constitutes a multigenerational or intergenerational direct exposure versus a true non-exposed transgenerational generation. A number of past studies have referred to F1 generation fetal exposure or F1 generation germline that will generate the F2 generation as transgenerational experiments (Fig. 3). Many previous reports have not carefully considered this issue and misinterpreted the results as transgenerational. A multigenerational or intergenerational exposure experiment is important and helps elucidate risk of exposure on multiple generations physiology and pathology. However, the mechanisms involved are distinct and impacts different than transgenerational generations [31]. This non-genetic form of inheritance needs to be distinguished from multiple generation exposure that is due to direct exposures and toxicities.

Another experimental design issue is the use of mixed cell populations for an epigenetic analysis [120]. Every cell type in the body has the same DNA sequence, so for genetic analysis a mixed cell population does not affect the data or observations. In contrast, each cell type in the organism has a very distinct epigenome to allow the cell type to have its unique cell biology and physiology. The reason a neuron is distinct from a hepatocyte is not the genetic sequence, but the epigenetic differences between the cell types that regulate the unique gene expression. Therefore, an epigenetic analysis of mixed cell populations are influenced by small changes in specific cell population numbers which will alter the epigenetic data experimentally observed without an actual change in molecular epigenetics [120]. A number of epigenetic studies have used whole blood which contains over 20 different cell populations to do epigenetics. Twin studies using this approach have not been revealing due to the variation in cell populations in the blood and inability to dissect out specific epigenetic changes. Purifying a specific cell type such as monocytes from the blood will be far more useful for epigenetic analyses than use of the mixed cell population. Therefore, epigenetic analysis optimally requires purified cell populations [120].

Epigenetic molecular procedures have dramatically developed over the past decade to provide greater accuracy and precision. The technology of next generation sequencing is superior to array technology and previous biochemical procedures. The current procedures for DNA methylation, ncRNA and histone modifications use next generation sequencing which should be considered the optimal approach for any genome-wide analysis. If a few selected sites are examined then the array technology or biochemical approach can be used and are less costly. For the genome-wide approaches, the different DNA methylation approaches are methylated DNA immunoprecipitation (MeDIP) sequencing (MeDIP-Seq) and bisulfite sequencing (BS-Seq). The MeDIP-Seq is biased to low density CpG <20% while the BS-Seq is biased to high density CpG. All these

procedures are efficient, but the limitation in CpG bias needs to be considered in the interpretation of the data obtained. The RNA-Seq and chromatin immunoprecipitation ChIP-Seq approaches are the optimal procedures currently available with few alterations. Third generation sequencing that may be able to assess epigenetic modifications during the sequencing will be a future technology to elucidate the DNA methylation CpG density bias, but remains to be optimized. The rate at which epigenetic technology is developing suggests within the next five years we will likely be using new technologies. The research in this area needs to consider the limitations of some of the technology currently used.

Conclusions

Research in the area of environmentally induced epigenetic transgenerational inheritance of disease and phenotypic variation has provided evidence of transgenerational inheritance of epimutations in plants, worms, flies, fish, birds, pigs, mice, rats, and humans [121] (Fig. 4). Ancestral exposure to environmental influences such as toxicants, abnormal nutrition, or stress can induce changes in the germline epigenome that are transmitted to descendants. These epimutations caused by individual exposures must occur in the germline in order to be transmitted. When these germline epigenetic changes become imprinted-like and escape the normal processes of epigenetic reprogramming, then epigenetic transgenerational inheritance can occur. Since the embryonic stem cells develop an altered epigenome, these epimutations subsequently induce somatic cell alterations in the epigenome and transcriptome, which will increase disease susceptibility in the offspring. Therefore, these ancestral exposures to environmental toxicants can lead to transgenerational changes in the epigenome and transcriptome of future generations and lead to an increased incidence of disease. Although DNA methylation is the most thoroughly studied epigenetic mechanism, other epigenetic processes are equally important. Future research will need to investigate the multiple epigenetic mechanisms and how they integrate. The developmental aspects of how the epigenetic transgenerational inheritance of disease develops are still unclear. How epimutations in sperm result in epigenetic changes in the resultant embryo needs to be investigated. How the derived embryonic stem cell changes can lead to epigenetic and transcriptome changes in the function of an adult organ associated with disease also remain to be elucidated on a molecular level. The potential role these ancestral exposures and epigenetic transgenerational inheritance have on disease etiology needs to be seriously considered. In addition, it may be clinically useful to determine what epimutation patterns or signatures are associated with specific disease and/or ancestral exposures in humans. Epigenetic biomarker signatures may be used in the future as a diagnostic tool to assess if an individual has a specific disease susceptibility or environmental toxicant exposures. This will facilitate preventive medicine and therapeutic approaches to mitigate associated disease risks.

Acknowledgements

We apologize to authors whose studies on the topic were not presented or referenced, but the increase in research in the area has grown significantly. We acknowledge the assistance of Dr Millissia Ben Maamar for critical review of the

manuscript and Ms Heather Johnson in preparing the manuscript. This was supported by a John Templeton Foundation grant (50183) and NIH (ES012974) grant to M.K.S.

Conflict of interest statement. None declared.

References

- Waddington CH. *Organisers and Genes*. Cambridge: Cambridge University Press, 1940.
- Van Speybroeck L. From epigenesis to epigenetics: the case of C. H. Waddington. *Ann N Y Acad Sci* 2002;**981**:61–81.
- Skinner MK. Environmental epigenetic transgenerational inheritance and somatic epigenetic mitotic stability. *Epigenetics* 2011;**6**:838–42.
- Holliday R, Pugh JE. DNA modification mechanisms and gene activity during development. *Science* 1975;**187**:226–32.
- Turner BM. Histone acetylation as an epigenetic determinant of long-term transcriptional competence. *Cell Mol Life Sci* 1998;**54**:21–31.
- Jodar M, Selvaraju S, Sandler E, Diamond MP, Krawetz SA. The presence, role and clinical use of spermatozoal RNAs. *Hum Reprod Update* 2013;**19**:604–24.
- Mattick JS. The genetic signatures of noncoding RNAs. *PLoS Genet* 2009;**5**:e1000459.
- Yaniv M. Chromatin remodeling: from transcription to cancer. *Cancer Genet* 2014;**207**:352–7.
- Schaefer M, Pollex T, Hanna K, Tuorto F, Meusburger M, Helm M, Lyko F. RNA methylation by Dnmt2 protects transfer RNAs against stress-induced cleavage. *Genes Dev* 2010;**24**:1590–5.
- Pokholok DK, Harbison CT, Levine S, Cole M, Hannett NM, Lee TI, Bell GW, Walker K, Rolfe PA, Herbolzheimer E, Zeitlinger J, Lewitter F, Gifford DK, Young RA. Genome-wide map of nucleosome acetylation and methylation in yeast. *Cell* 2005;**122**:517–27.
- Boland MJ, Nazor KI, Loring JF. Epigenetic regulation of pluripotency and differentiation. *Circ Res* 2014;**115**:311–24.
- Avgustinova A, Benitah SA. Epigenetic control of adult stem cell function. *Nat Rev Mol Cell Biol* 2016;**17**:643–58.
- Singer J, Roberts-Ems J, Riggs AD. Methylation of mouse liver DNA studied by means of the restriction enzymes msp I and hpa II. *Science* 1979;**203**:1019–21.
- Kriaucionis S, Tahiliani M. Expanding the epigenetic landscape: novel modifications of cytosine in genomic DNA. *Cold Spring Harb Perspect Biol* 2014;**6**:a018630.
- An J, Rao A, Ko M. TET family dioxygenases and DNA demethylation in stem cells and cancers. *Exp Mol Med* 2017;**49**:e323.
- Mellen M, Ayata P, Heintz N. 5-hydroxymethylcytosine accumulation in postmitotic neurons results in functional demethylation of expressed genes. *Proc Natl Acad Sci USA* 2017;**114**:E7812–21.
- Tang WW, Dietmann S, Irie N, Leitch HG, Floros VI, Bradshaw CR, Hackett JA, Chinnery PF, Surani MA. A unique gene regulatory network resets the human germline epigenome for development. *Cell* 2015;**161**:1453–67.
- Wu TP, Wang T, Seetin MG, Lai Y, Zhu S, Lin K, Liu Y, Byrum SD, Mackintosh SG, Zhong M, Tackett A, Wang G, Hon LS, Fang G, Swenberg JA, Xiao AZ. DNA methylation on N(6)-adenine in mammalian embryonic stem cells. *Nature* 2016;**532**:329–33.
- Rothbart SB, Strahl BD. Interpreting the language of histone and DNA modifications. *Biochim Biophys Acta* 2014;**1839**:627–43.
- Kornfeld JW, Bruning JC. Regulation of metabolism by long, non-coding RNAs. *Front Genet* 2014;**5**:57.
- Wei JW, Huang K, Yang C, Kang CS. Non-coding RNAs as regulators in epigenetics (Review). *Oncol Rep* 2017;**37**:3–9.
- Chen Q, Yan M, Cao Z, Li X, Zhang Y, Shi J, Feng GH, Peng H, Zhang X, Zhang Y, Qian J, Duan E, Zhai Q, Zhou Q. Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* 2016;**351**:397–400.
- Chen Q, Yan W, Duan E. Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. *Nat Rev Genet* 2016;**17**:733–43.
- Sibbritt T, Patel HR, Preiss T. Mapping and significance of the mRNA methylome. *Wiley Interdiscip Rev RNA* 2013;**4**:397–422.
- Yue Y, Liu J, He C. RNA N6-methyladenosine methylation in post-transcriptional gene expression regulation. *Genes Dev* 2015;**29**:1343–55.
- Fu Y, Dominissini D, Rechavi G, He C. Gene expression regulation mediated through reversible m(6)A RNA methylation. *Nat Rev Genet* 2014;**15**:293–306.
- Xu L, Liu X, Sheng N, Oo KS, Liang J, Chionh YH, Xu J, Ye F, Gao YG, Dedon PC, Fu XY. Three distinct 3-methylcytidine (m(3)C) methyltransferases modify tRNA and mRNA in mice and humans. *J Biol Chem* 2017;**292**:14695–703.
- Lyons SM, Fay MM, Akiyama Y, Anderson PJ, Ivanov P. RNA biology of angiogenin: current state and perspectives. *RNA Biol* 2017;**14**:171–8.
- Hussain S, Sajini AA, Blanco S, Dietmann S, Lombard P, Sugimoto Y, Paramor M, Gleeson JG, Odom DT, Ule J, Frye M. NSun2-mediated cytosine-5 methylation of vault noncoding RNA determines its processing into regulatory small RNAs. *Cell Rep* 2013;**4**:255–61.
- Saikia M, Hatzoglou M. The many virtues of tRNA-derived stress-induced RNAs (tiRNAs): discovering novel mechanisms of stress response and effect on human health. *J Biol Chem* 2015;**290**:29761–8.
- Skinner MK. What is an epigenetic transgenerational phenotype? F3 or F2. *Reprod Toxicol* 2008;**25**:2–6.
- Blewitt ME, Vickaryous NK, Paldi A, Koseki H, Whitelaw E. Dynamic reprogramming of DNA methylation at an epigenetically sensitive allele in mice. *PLoS Genet* 2006;**2**:e49.
- Waterland RA, Travisano M, Tahiliani KG. Diet-induced hypermethylation at agouti viable yellow is not inherited transgenerationally through the female. *FASEB J* 2007;**21**:3380–5.
- Blewitt M, Whitelaw E. The use of mouse models to study epigenetics. *Cold Spring Harb Perspect Biol* 2013;**5**:a017939.
- Waddington CH. Canalisation of development and the inheritance of acquired characters. *Nature* 1942;**150**:563–5.
- Stockard CR, Papanicolaou GN. Further studies on the modification of the germ-cells in mammals: the effect of alcohol on treated Guinea pigs and their descendants. *J Exp Zool* 1918;**26**:119–226.
- Anway MD, Cupp AS, Uzumcu M, Skinner MK. Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science* 2005;**308**:1466–9.
- Chamorro-Garcia R, Sahu M, Abbey RJ, Laude J, Pham N, Blumberg B. Transgenerational inheritance of increased fat depot size, stem cell reprogramming, and hepatic steatosis elicited by prenatal exposure to the obesogen tributyltin in mice. *Environ Health Perspect* 2013;**121**:359–66.
- Skinner MK. Endocrine disruptor induction of epigenetic transgenerational inheritance of disease. *Mol Cell Endocrinol* 2014;**398**:4–12.

40. Quinnes KM, Doyle TJ, Kim KH, Rissman EF. Transgenerational effects of di-(2-ethylhexyl) phthalate (DEHP) on stress hormones and behavior. *Endocrinology* 2015; **156**:3077–83.
41. Rattan S, Brehm E, Gao L, Niermann S, Flaws JA. Prenatal exposure to di(2-ethylhexyl) phthalate disrupts ovarian function in a transgenerational manner in female mice. *Biol Reprod* 2018; **98**:130–45.
42. Wolstenholme JT, Edwards M, Shetty SR, Gatewood JD, Taylor JA, Rissman EF, Connelly JJ. Gestational exposure to bisphenol A produces transgenerational changes in behaviors and gene expression. *Endocrinology* 2012; **153**:3828–38.
43. Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner M. Plastics derived endocrine disruptors (BPA, DEHP and DBP) induce epigenetic transgenerational inheritance of obesity, reproductive disease and sperm epimutations. *PLoS One* 2013; **8**:1–18, e55387.
44. Chamorro-Garcia R, Diaz-Castillo C, Shoucri B, Kaech M, Leavitt H, Shioda RT, Blumberg B. Ancestral perinatal obesogen exposure results in a transgenerational thrifty phenotype in mice. *Nat Commun* 2017; **8**:2012.
45. Skinner MK, Manikkam M, Tracey R, Guerrero-Bosagna C, Haque MM, Nilsson E. Ancestral dichlorodiphenyltrichloroethane (DDT) exposure promotes epigenetic transgenerational inheritance of obesity. *BMC Med* 2013; **11**:228, 1–16.
46. Tracey R, Manikkam M, Guerrero-Bosagna C, Skinner M. Hydrocarbons (jet fuel JP-8) induce epigenetic transgenerational inheritance of obesity, reproductive disease and sperm epimutations. *Reprod Toxicol* 2013; **36**:104–16.
47. Abbott CW, Rohac DJ, Bottom RT, Patadia S, Huffman KJ. Prenatal ethanol exposure and neocortical development: a transgenerational model of FASD. *Cereb Cortex* 2017; **1**–14, bxx168.
48. Lombo M, Fernandez-Diez C, Gonzalez-Rojo S, Navarro C, Robles V, Herraez MP. Transgenerational inheritance of heart disorders caused by paternal bisphenol A exposure. *Environ Pollut* 2015; **206**:667–78.
49. Carvan MJI, Kalluvila TA, Klingler RH, Larson JK, Pickens M, Mora-Zamorano FX, Connaughton VP, Sadler-Riggleman I, Beck D, Skinner MK. Mercury-induced epigenetic transgenerational inheritance of abnormal neurobehavior is correlated with sperm epimutations in zebrafish. *PLoS One* 2017; **12**:e0176155–26.
50. Knecht AL, Truong L, Marvel SW, Reif DM, Garcia A, Lu C, Simonich MT, Teeguarden JG, Tanguay RL. Transgenerational inheritance of neurobehavioral and physiological deficits from developmental exposure to benzo[a]pyrene in zebrafish. *Toxicol Appl Pharmacol* 2017; **329**:148–57.
51. Razoux F, Russig H, Mueggler T, Baltes C, Dikaoui K, Rudin M, Mansuy IM. Transgenerational disruption of functional 5-HT_{1A}R-induced connectivity in the adult mouse brain by traumatic stress in early life. *Mol Psychiatry* 2017; **22**:519–26.
52. Franklin TB, Linder N, Russig H, Thony B, Mansuy IM. Influence of early stress on social abilities and serotonergic functions across generations in mice. *PLoS One* 2011; **6**:e21842.
53. He N, Kong QQ, Wang JZ, Ning SF, Miao YL, Yuan HJ, Gong S, Cui XZ, Li CY, Tan JH. Parental life events cause behavioral difference among offspring: adult pre-gestational restraint stress reduces anxiety across generations. *Sci Rep* 2016; **6**:39497.
54. Saavedra-Rodriguez L, Feig LA. Chronic social instability induces anxiety and defective social interactions across generations. *Biol Psychiatry* 2013; **73**:44–53.
55. Kiss D, Ambeskovic M, Montina T, Metz GA. Stress transgenerationally programs metabolic pathways linked to altered mental health. *Cell Mol Life Sci* 2016; **73**:4547–57.
56. Bygren LO, Tinghog P, Carstensen J, Edvinsson S, Kaati G, Pembrey ME, Sjöström M. Change in paternal grandmothers' early food supply influenced cardiovascular mortality of the female grandchildren. *BMC Genet* 2014; **15**:12.
57. Dunn GA, Bale TL. Maternal high-fat diet effects on third-generation female body size via the paternal lineage. *Endocrinology* 2011; **152**:2228–36.
58. Masuyama H, Mitsui T, Nobumoto E, Hiramatsu Y. The effects of high-fat diet exposure in utero on the obesogenic and diabetogenic traits through epigenetic changes in adiponectin and leptin gene expression for multiple generations in female mice. *Endocrinology* 2015; **156**:2482–91.
59. Nguyen NM, de Oliveira Andrade F, Jin L, Zhang X, Macon M, Cruz MI, Benitez C, Wehrenberg B, Yin C, Wang X, Xuan J, de Assis S, Hilakivi-Clarke L. Maternal intake of high n-6 polyunsaturated fatty acid diet during pregnancy causes transgenerational increase in mammary cancer risk in mice. *Breast Cancer Res* 2017; **19**:77.
60. Skinner MK, Anway MD. Epigenetic transgenerational actions of vinclozolin on the development of disease and cancer. *Crit Rev Oncog* 2007; **13**:75–82.
61. Pavlinkova G, Margaryan H, Zatecka E, Valaskova E, Elzeinova F, Kubatova A, Bohuslavova R, Peknicova J. Transgenerational inheritance of susceptibility to diabetes-induced male subfertility. *Sci Rep* 2017; **7**:4940.
62. Wei Y, Yang CR, Wei YP, Zhao ZA, Hou Y, Schatten H, Sun QY. Paternally induced transgenerational inheritance of susceptibility to diabetes in mammals. *Proc Natl Acad Sci USA* 2014; **111**:1873–8.
63. de Castro Barbosa T, Ingerslev LR, Alm PS, Versteijhe S, Massart J, Rasmussen M, Donkin I, Sjögren R, Mudry JM, Vetterli L, Gupta S, Krook A, Zierath JR, Barres R. High-fat diet reprograms the epigenome of rat spermatozoa and transgenerationally affects metabolism of the offspring. *Mol Metab* 2016; **5**:184–97.
64. Quadrana L, Colot V. Plant transgenerational epigenetics. *Annu Rev Genet* 2016; **50**:467–91.
65. Greer EL, Maures TJ, Ucar D, Hauswirth AG, Mancini E, Lim JP, Benayoun BA, Shi Y, Brunet A. Transgenerational epigenetic inheritance of longevity in *Caenorhabditis elegans*. *Nature* 2011; **479**:365–71.
66. Buescher JL, Musselman LP, Wilson CA, Lang T, Keleher M, Baranski TJ, Duncan JG. Evidence for transgenerational metabolic programming in *Drosophila*. *Dis Model Mech* 2013; **6**:1123–32.
67. Dew-Budd K, Jarnigan J, Reed LK. Genetic and sex-specific transgenerational effects of a high fat diet in *Drosophila melanogaster*. *PLoS One* 2016; **11**:e0160857.
68. Xia B, de Belle JS. Transgenerational programming of longevity and reproduction by post-eclosion dietary manipulation in *Drosophila*. *Aging Aging* 2016; **8**:1115–34.
69. Xia B, Gerstin E, Schones DE, Huang W, Steven de Belle J. Transgenerational programming of longevity through E(z)-mediated histone H3K27 trimethylation in *Drosophila*. *Aging (Aging)* 2016; **8**:2988–3008.
70. Vandegehuchte MB, Lemiere F, Vanhaecke L, Vanden Berghe W, Janssen CR. Direct and transgenerational impact on *Daphnia magna* of chemicals with a known effect on DNA methylation. *Comp Biochem Physiol C Toxicol Pharmacol* 2010; **151**:278–85.

71. Baker TR, Peterson RE, Heideman W. Using zebrafish as a model system for studying the transgenerational effects of dioxin. *Toxicol Sci* 2014;**138**:403–11.
72. Bhandari RK, vom Saal FS, Tillitt DE. Transgenerational effects from early developmental exposures to bisphenol A or 17alpha-ethinylestradiol in medaka, *Oryzias latipes*. *Sci Rep* 2015;**5**:9303.
73. Leroux S, Gourichon D, Letierrier C, Labrune Y, Coustham V, Riviere S, Zerjal T, Coville JL, Morisson M, Minvielle F, Pitel F. Embryonic environment and transgenerational effects in quail. *Genet Sel Evol* 2017;**49**:14.
74. Brun JM, Bernadet MD, Cornuez A, Leroux S, Bodin L, Basso B, Davail S, Jaglin M, Lessire M, Martin X, Sellier N, Morisson M, Pitel F. Influence of grand-mother diet on offspring performances through the male line in Muscovy duck. *BMC Genet* 2015;**16**:145.
75. Skinner MK, Manikkam M, Guerrero-Bosagna C. Epigenetic transgenerational actions of endocrine disruptors. *Reprod Toxicol* 2011;**31**:337–43.
76. Braunschweig M, Jagannathan V, Gutzwiller A, Bee G. Investigations on transgenerational epigenetic response down the male line in F2 pigs. *PLoS One* 2012;**7**:e30583.
77. Veenendaal MV, Painter RC, de Rooij SR, Bossuyt PM, van der Post JA, Gluckman PD, Hanson MA, Roseboom TJ. Transgenerational effects of prenatal exposure to the 1944–45 Dutch famine. *BJOG* 2013;**120**:548–53.
78. Anway MD, Leathers C, Skinner MK. Endocrine disruptor vinclozolin induced epigenetic transgenerational adult-onset disease. *Endocrinology* 2006;**147**:5515–23.
79. Anway MD, Skinner MK. Transgenerational effects of the endocrine disruptor vinclozolin on the prostate transcriptome and adult onset disease. *Prostate* 2008;**68**:517–29.
80. Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner MK. Dioxin (TCDD) induces epigenetic transgenerational inheritance of adult onset disease and sperm epimutations. *PLoS One* 2012;**7**:e46249–15.
81. Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner M. Pesticide and insect repellent mixture (Permethrin and DEET) induces epigenetic transgenerational inheritance of disease and sperm epimutations. *Reprod Toxicol* 2012;**34**:708–19.
82. Manikkam M, Guerrero-Bosagna C, Tracey R, Haque MM, Skinner MK. Transgenerational actions of environmental compounds on reproductive disease and identification of epigenetic biomarkers of ancestral exposures. *PLoS One* 2012;**7**:1–12, e31901.
83. Nilsson EE, Anway MD, Stanfield J, Skinner MK. Transgenerational epigenetic effects of the endocrine disruptor vinclozolin on pregnancies and female adult onset disease. *Reproduction* 2008;**135**:713–21.
84. Nilsson EE, Skinner MK. Environmentally induced epigenetic transgenerational inheritance of reproductive disease. *Biol Reprod* 2015;**93**:145.
85. Skinner MK, Anway MD, Savenkova MI, Gore AC, Crews D. Transgenerational epigenetic programming of the brain transcriptome and anxiety behavior. *PLoS One* 2008;**3**:1–11, e3745.
86. Manikkam M, Haque MM, Guerrero-Bosagna C, Nilsson E, Skinner MK. Pesticide methoxychlor promotes the epigenetic transgenerational inheritance of adult onset disease through the female germline. *PLoS One* 2014;**9**:1–19, e102091.
87. Nilsson E, Larsen G, Manikkam M, Guerrero-Bosagna C, Savenkova M, Skinner M. Environmentally induced epigenetic transgenerational inheritance of ovarian disease. *PLoS One* 2012;**7**:1–18, e36129.
88. Bateson P, Gluckman P, Hanson M. The biology of developmental plasticity and the predictive adaptive response hypothesis. *J Physiol* 2014;**592**:2357–68.
89. Skinner MK. Environmental epigenetics and a unified theory of the molecular aspects of evolution: a neo-lamarckian concept that facilitates neo-Darwinian evolution. *Genome Biol Evol* 2015;**7**:1296–302.
90. Hackett JA, Surani MA. Beyond DNA: programming and inheritance of parental methylomes. *Cell* 2013;**153**:737–9.
91. Smith ZD, Chan MM, Mikkelsen TS, Gu H, Gnirke A, Regev A, Meissner A. A unique regulatory phase of DNA methylation in the early mammalian embryo. *Nature* 2012;**484**:339–44.
92. Constancia M, Pickard B, Kelsey G, Reik W. Imprinting mechanisms. *Genome Res* 1998;**8**:881–900.
93. Skinner MK, Manikkam M, Guerrero-Bosagna C. Epigenetic transgenerational actions of environmental factors in disease etiology. *Trends Endocrinol Metab* 2010;**21**:214–22.
94. Jirtle RL, Skinner MK. Environmental epigenomics and disease susceptibility. *Nat Rev Genet* 2007;**8**:253–62.
95. Guerrero-Bosagna C, Savenkova M, Haque MM, Nilsson E, Skinner MK. Environmentally induced epigenetic transgenerational inheritance of altered sertoli cell transcriptome and epigenome: molecular etiology of male infertility. *PLoS One* 2013;**8**:1–12, e59922.
96. Skinner MK, Manikkam M, Haque MM, Zhang B, Savenkova M. Epigenetic transgenerational inheritance of somatic transcriptomes and epigenetic control regions. *Genome Biol* 2012;**13**:R91.
97. Guerrero-Bosagna C, Settles M, Lucker B, Skinner M. Epigenetic transgenerational actions of vinclozolin on promoter regions of the sperm epigenome. *PLoS One* 2010;**5**:1–17, e13100.
98. Skinner MK, Guerrero-Bosagna C. Role of CpG deserts in the epigenetic transgenerational inheritance of differential DNA methylation regions. *BMC Genomics* 2014;**15**:692.
99. Haque MM, Holder LB, Skinner MK. Genome-wide locations of potential epimutations associated with environmentally induced epigenetic transgenerational inheritance of disease using a sequential machine learning prediction approach. *PLoS One* 2015;**10**:1–25, e0142274.
100. Beck D, Sadler-Riggelman I, Skinner MK. Generational comparisons (F1 versus F3) of vinclozolin induced epigenetic transgenerational inheritance of sperm differential DNA methylation regions (epimutations) using MeDIP-Seq. *Environ Epigenetics* 2017;**3**:1–12, dvx016.
101. Rodgers AB, Morgan CP, Leu NA, Bale TL. Transgenerational epigenetic programming via sperm microRNA recapitulates effects of paternal stress. *Proc Natl Acad Sci USA* 2015;**112**:13699–704.
102. Schuster A, Skinner MK, Yan W. Ancestral vinclozolin exposure alters the epigenetic transgenerational inheritance of sperm small noncoding RNAs. *Environ Epigenet* 2016;**2**:1–10, dvw001.
103. Gapp K, Jawaid A, Sarkies P, Bohacek J, Pelczar P, Prados J, Farinelli L, Miska E, Mansuy IM. Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nat Neurosci* 2014;**17**:667–9.
104. Skinner MK, Ben Maamar M, Sadler-Riggelman I, Beck D, Nilsson E, McBirney M, Klukovich R, Xie Y, Tang C, Yan W. Alterations in sperm DNA methylation, non-coding RNA and histone retention associate with DDT-induced epigenetic transgenerational inheritance of disease. *Epigenet Chromatin* 2018;**4**:8.

105. Bao J, Bedford MT. Epigenetic regulation of the histone-to-protamine transition during spermiogenesis. *Reproduction* 2016;**151**:R55–70.
106. Rathke C, Baarends WM, Awe S, Renkawitz-Pohl R. Chromatin dynamics during spermiogenesis. *Biochim Biophys Acta* 2014;**1839**:155–68.
107. Ihara M, Meyer-Ficca ML, Leu NA, Rao S, Li F, Gregory BD, Zalenskaya IA, Schultz RM, Meyer RG. Paternal poly (ADP-ribose) metabolism modulates retention of inheritable sperm histones and early embryonic gene expression. *PLoS Genet* 2014;**10**:e1004317.
108. Ben Maamar M, Sadler-Riggelman I, Beck D, Skinner MK. Epigenetic transgenerational inheritance of altered sperm histone retention sites. *Sci Rep* 2018;**8**:5308.
109. Hammoud SS, Nix DA, Hammoud AO, Gibson M, Cairns BR, Carrell DT. Genome-wide analysis identifies changes in histone retention and epigenetic modifications at developmental and imprinted gene loci in the sperm of infertile men. *Hum Reprod* 2011;**26**:2558–69.
110. Siklenka K, Erkek S, Godmann M, Lambrot R, McGraw S, Laffleur C, Cohen T, Xia J, Suderman M, Hallett M, Trasler J, Peters AH, Kimmins S. Disruption of histone methylation in developing sperm impairs offspring health transgenerationally. *Science* 2015;**350**: 1–14, aab2006.
111. Kelly WG. Transgenerational epigenetics in the germline cycle of *Caenorhabditis elegans*. *Epigenetics Chromatin* 2014;**7**:6.
112. Ruden DM, Lu X. Hsp90 affecting chromatin remodeling might explain transgenerational epigenetic inheritance in *Drosophila*. *Cg* 2008;**9**:500–8.
113. Ben Maamar M, Sadler-Riggelman I, Beck D, McBirney M, Nilsson E, Klukovich R, Xie Y, Tang C, Yan W, Skinner MK. Alterations in sperm DNA Methylation, Non-Coding RNA expression, and histone retention mediate Vinclozolin induced epigenetic transgenerational inheritance of disease. *Environmental Epigenetics* 2018;**4**:1–19, dvy010.
114. Kung JT, Colognori D, Lee JT. Long noncoding RNAs: past, present, and future. *Genetics* 2013;**193**:651–69.
115. Saha P, Verma S, Pathak RU, Mishra RK. Long noncoding RNAs in Mammalian development and diseases. *Adv Exp Med Biol* 2017;**1008**:155–98.
116. Liu KS, Li TP, Ton H, Mao XD, Chen YJ. Advances of long non-coding RNAs-mediated regulation in reproduction. *Chin Med J* 2018;**131**:226–34.
117. Haque MM, Nilsson EE, Holder LB, Skinner MK. Genomic clustering of differential DNA methylated regions (epimutations) associated with the epigenetic transgenerational inheritance of disease and phenotypic variation. *BMC Genomics* 2016;**17**:418.
118. Pastuszak AW, Lamb DJ. The genetics of male fertility—from basic science to clinical evaluation. *J Androl* 2012;**33**:1075–84.
119. Matzuk MM, Lamb DJ. The biology of infertility: research advances and clinical challenges. *Nat Med* 2008;**14**:1197–213.
120. Skinner MK. Differential DNA methylation analysis optimally requires purified cell populations. *Fertil Steril* 2016;**106**: 551.
121. Hanson MA, Skinner MK. Developmental origins of epigenetic transgenerational inheritance. *Environ Epigenet* 2016;**2**: 1–9, dvw002.
122. Skinner MK. A new kind of inheritance. *Sci Am* 2014;**311**: 44–51.
123. Bruner-Tran KL, Osteen KG. Developmental exposure to TCDD reduces fertility and negatively affects pregnancy outcomes across multiple generations. *Reprod Toxicol* 2011;**31**:344–50.
124. Doyle TJ, Bowman JL, Windell VL, McLean DJ, Kim KH. Transgenerational effects of di-(2-ethylhexyl) phthalate on testicular germ cell associations and spermatogonial stem cells in mice. *Biol Reprod* 2013;**88**:112.
125. McBirney M, King SE, Michelle Pappalardo M, Houser E, Unkefer M, Nilsson E, Sadler-Riggelman I, Beck D, Winchester P, Skinner MK. Atrazine induced epigenetic transgenerational inheritance of disease, lean phenotype and sperm epimutation pathology biomarkers. *PLoS One* 2017;**12**:e0184306–37.
126. Padmanabhan N, Jia D, Geary-Joo C, Wu X, Ferguson-Smith AC, Fung E, Bieda MC, Snyder FF, Gravel RA, Cross JC, Watson ED. Mutation in folate metabolism causes epigenetic instability and transgenerational effects on development. *Cell* 2013;**155**:81–93.
127. Zheng X, Chen L, Li M, Lou Q, Xia H, Wang P, Li T, Liu H, Luo L. Transgenerational variations in DNA methylation induced by drought stress in two rice varieties with distinguished difference to drought resistance. *PLoS One* 2013;**8**:e80253.
128. Suter L, Widmer A. Environmental heat and salt stress induce transgenerational phenotypic changes in *Arabidopsis thaliana*. *PLoS One* 2013;**8**:e60364.
129. Rehan VK, Liu J, Sakurai R, Torday JS. Perinatal nicotine-induced transgenerational asthma. *Am J Physiol Lung Cell Mol Physiol* 2013;**305**:L501–7.
130. Govorko D, Bekdash RA, Zhang C, Sarkar DK. Male germline transmits fetal alcohol adverse effect on hypothalamic proopiomelanocortin gene across generations. *Biol Psychiatry* 2012;**72**:378–88.
131. Norouzitallab P, Baruah K, Vandegehuchte M, Van Stappen G, Catania F, Vanden Bussche J, Vanhaecke L, Sorgeloos P, Bossier P. Environmental heat stress induces epigenetic transgenerational inheritance of robustness in parthenogenetic *Artemia* model. *FASEB J* 2014;**28**:3552–63.
132. Waddington CH. Gene assimilation of an acquired character. *Evolution* 1953;**7**:118–26.

Spring 2020 – Systems Biology of Reproduction
 Lecture Outline – Epigenetics and Transgenerational Reproductive Disease
 Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday
 March 10, 2020
 Week 9

Epigenetics and Transgenerational Reproductive Disease

- Environmental Induced Pathology
- Transgenerational Disease Phenotype
- Compound Specificity
- Epigenetic Mechanism
- Epimutations and Exposure Specificity
- Transgenerational Transcriptome
- Transgenerational Testis Disease
- Transgenerational Ovary Disease
- Broader Impact

REQUIRED READING

Nilsson EE, Sadler-Riggleman I, Skinner MK. Environmentally induced epigenetic transgenerational inheritance of disease. Environ Epigenet. 2018 Jul 17;4(2):1-13, dvy016.

Spring 2020 – Systems Biology of Reproduction
 Discussion Outline – Epigenetics and Transgenerational Reproductive Disease
 Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday
 March 12, 2020
 Week 9

Epigenetics and Transgenerational Reproductive Disease

Primary Papers:

1. Kubsad, et al. (2019) Scientific Reports 9 :6372
2. Sadler-Riggleman, et al. (2019) Environmental Epigenetics 5(3) dvz013
3. Ben Maamar, et al. (2020) Developmental Biology 458:106-119

Discussion

Student 22: Reference 1 above

- What transgenerational reproductive phenotypes were observed?
- What is glyphosate and effects on sperm?
- What is generational toxicology?


Student 23: Reference 2 above

- What environmental epigenetics were used?
- What was the Sertoli cell effects observed?
- What basic information on testis disease was obtained?

Student 24: Reference 3 above

- What is parent-of-origin allelic epigenetic inheritance?
- What transgenerational reproductive disease was observed?
- What transgenerational disease was transmitted through the only paternal allele?

GENETIC DETERMINISM




ENVIRONMENTAL IMPACT ON BIOLOGY

- Regional Disease Frequencies
- Low Frequency of Genetic Component of Disease
- Increases In Disease Frequencies
- Identical Twins and Variable Disease Frequency
- Environmental Exposures and Disease
- Evolution and Rapid Induction

GROWING SICKNESS

Although people are living longer, they are also living with more chronic conditions, as seen here in data for the developed world.


Number of conditions ■ 0 ■ 1-2 ■ 3-4 ■ 5-6 ■ 7-8 ■ ≥9




ENVIRONMENTAL IMPACT ON BIOLOGY




- Regional Disease Frequencies
- Low Frequency of Genetic Component of Disease
- Increases In Disease Frequencies
- Identical Twins and Variable Disease Frequency
- Environmental Exposures and Disease
- Evolution and Rapid Induction

GENETIC DETERMINISM



EPIGENETIC SOLUTIONS TO GENETIC DETERMINISM FAILURES







Model Endocrine Disruptor: Vinclozolin

- Vinclozolin is a systemic fungicide (e.g. Wine Industry)
- Vinclozolin and its metabolites are **anti-androgenic**
- Late embryonic/early postnatal exposure causes abnormal reproductive tract development and gonadal function

Vinclozolin Effects on Spermatogenic Cell Apoptosis



Germ Cell Apoptosis




* P<0.05


Vinclozolin

Complete Male Infertility (10%)

F3 Generation Males










Sperm Concentration P60-P150



* P<0.05

Vinclozolin





TRANSGENERATIONAL DISEASE ETIOLOGY

MALE	FEMALE
<ul style="list-style-type: none"> Spermatogenic Defect (>90%) Male infertility (complete ~10%, severe 20%) Prostate disease (~50%) 	<ul style="list-style-type: none"> Kidney disease (~30-40%) Increase in mammary tumor formation (~10-20%) Behavior (Mate Preference, Anxiety, & Stress) (>90%) Pubertal Abnormalities (~25%) Obesity (~10-50%) Lean (~10-20%) Immune (~20%) Pre-eclampsia-like during late pregnancy (~10%) Premature Ovarian Failure POF (>90%) Ovarian Polycystic Ovarian Disease (>90%) Female Premature Pubertal Onset (>90%) Parturition (birth) defects

Compound Specificity


Other Endocrine Disruptors And Environmental Compounds

Environmental Compound Specificity

(Exposure Groups)	F1	F3
A. Vinclozolin [agricultural fungicide]	Yes	Yes
B. Flutamide [anti-androgenic pharmaceutical]	Yes	No
C. TCDD/Dioxin (industrial pollutant)	Yes	Yes
D. Plastics Compounds [Bisphenol-A BPA, Phthalate-DEHP & DBP]	Yes	Yes
E. Jet Fuel [JP8] (Hydrocarbon Mixture)	Yes	Yes
F. Pesticide & Insect Repellent [Permethrin & DEET]	No	Yes
G. DDT (pesticide)	Yes	Yes
H. Methoxychlor (pesticide, replace DDT)	Yes	Yes
I. Mercury (Industrial pollutant)	Yes	Yes
J. Atrazine (agricultural herbicide)	No	Yes



Assessment of Glyphosate Induced Epigenetic Transgenerational Inheritance of Pathologies and Sperm Epimutations: Generational Toxicology.

Kubsad D, Nilsson EE, King SE, Sadler-Riggleman I, Beck D, Skinner MK.
Sci Rep. 2019 Apr 23;9(1):6372.



Transgenerational Disease Etiology

- Spermatogenic Defect (>90%)
- Male infertility (complete ~10%, severe 20%)
- Kidney disease (~30-40%)
- Prostate disease (~50%)
- Increase in mammary tumor formation (~10-20%)
- Behavior (Mate Preference, Anxiety & Stress) (>90%)
- Pre-eclampsia-like during late pregnancy (~10%)
- Premature Ovarian Failure POF (>90%)
- Ovarian Polycystic Ovarian Disease (>90%)
- Female Premature Pubertal Onset (>90%)
- Obesity (~10-50%)



ENVIRONMENTALLY INDUCED EPIGENETIC TRANSGENERATIONAL INHERITANCE

Environmental Toxicants

Vinclozolin (Agricultural Fungicide)	Permethrin & DEET (Insect Repellants)
Methoxychlor (Agricultural Pesticide)	DDT (Pesticide)
Dioxin/TCDD (Industrial Contaminant)	Tributyltin (Industrial Toxicant & Biocide)
Plastic Compounds (BPA & Phthalates)	Hydrocarbons (Jet Fuel)
Atrazine (Herbicide)	Glyphosate (Pesticide / Herbicide)

Other Types Exposures

Nutrition (High Fat or Caloric Restriction)	Smoking & Alcohol
Temperature & Drought (Plant Health & Flowering)	Stress Trauma (Behavioral)

Plants

Flies

Worms

Fish

Birds



Rodents

Pigs


Humans

Summary

- Transient Embryonic Exposure Effects Adult
 - Sex Determination Period
 - Fetal Basis of Disease
- Spermatogenic Fertility Defect & Other Diseases
- Transgenerational Phenotype
 - WHAT MECHANISM?




EXPOSURE SPECIFIC TRANSGENERATIONAL (F3) SPERM PROMOTER EPIGENOME ALTERATIONS




Plos One 2014

Chromatin Structure – Composition and Function during Spermiogenesis

Khochbin, S. and Ward, W.S



Environmentally Induced New Transgenerational Histone Retention




Epigenetic & Chromatin 2018

Ben Maamar, et al. (2018) Sci Rep. 28.8(1)5308


Developmental Origins of Transgenerational Sperm Histone Retention Following Ancestral Exposures

Ben Maamar M, Beck D, Nilsson E, McCarrey JR, and Skinner MK
Developmental Biology (submitted)




Developmental Origins of Sperm Epimutations (DHRs)




Ancestral vinclozolin exposure alters the epigenetic transgenerational inheritance of sperm small noncoding RNAs (CC)
 Andrew Schuster, Michael K. Skinner, Wei Yan
 Environmental Epigenetics (2016) 2 (1): 1-15




Small RNAs in the Transgenerational Inheritance of Epigenetic Information.
 Duempelmann L, Skribbe M, Bühler M.
 Trends Genet. 2020 Jan 14. doi: 10.1016/j.tig.2019.12.001. [Epub ahead of print]




Developmental origins of transgenerational sperm DNA methylation epimutations following ancestral DDT exposure.
 Ben Maamar M, Nilsson E, Sadler-Riggleman I, Beck D, McCarrey JR, Skinner MK.
 Dev Biol. 2019 Jan 15;445(2):280-293.



Transgenerational sperm DNA methylation epimutation developmental origins following ancestral vinclozolin exposure.
 Skinner MK, Nilsson E, Sadler-Riggleman I, Beck D, Ben Maamar M, McCarrey JR.
 Epigenetics. 2019 Jul;14(7):721-739.




Developmental Origins of Sperm Epimutations (DMRs)




Collaborator- John McCarrey, UTSA

Ben Maamar, et al. (2019) Dev Biol 15;445(2):280-293
 Skinner, et al (2019) Epigenetics 14(7):721-739

Female / Male Environmental Exposure



Gestating Female Environmental Exposure




Multigenerational Exposures


Transgenerational Inheritance

Environmentally induced transgenerational epigenetic inheritance: Schematic of environmental exposure and affected generations.

Generational comparisons (F1 versus F3) of vinclozolin induced epigenetic transgenerational inheritance of sperm differential DNA methylation regions (epimutations) using MeDIP-Seq
 Environ Epigenet. 2017 Jul;3(3). pii: dx016.
 Beck D, Sadler-Riggleman I, Skinner MK.



Unravelling the complex mechanisms of transgenerational epigenetic inheritance
 Curr Opin Chem Biol. 2016 Aug;33:101-7.
 Blake GE, Watson ED.




F3 Generation Sperm Epigenome Mapping (Epimutations) (Epigenetic Biomarkers for Ancestral Exposures)

Transgenerational Sperm Epigenome Alterations (>200 differential DNA methylation sites)



Potential DNA Sequence Motif Identified (EDM1) & CpG% (Susceptibility epigenetic transgenerational mark)

Genome Activity Alterations? (transcriptome)


E16 Testis Transcriptome




Tissue Specific Transgenerational Transcriptomes (F3)



Epigenetic Control Regions





Epigenetic Control Region




Transgenerational Transcriptomes (F3 Generation)





- Variety Somatic Tissues and Cell Types
- Cell and Tissue Specific Altered Transcriptomes (>10X more genes than epimutations)
- Gene Clusters associate with Epimutations (Epigenetic Control Regions)
- Epimutation transmission to disease?**





Epigenetic Transgenerational Inheritance of Ovarian Disease







Environmental toxicant induced epigenetic transgenerational inheritance of ovarian pathology and granulosa cell epigenome and transcriptome alterations: ancestral origins of polycystic ovarian syndrome and primary ovarian insufficiency. Nilsen E, Klukovich R, Sadler-Riggleman I, Beck D, Xie Y, Yan W, Skinner MK. *Epigenetics*. 2018;13(8):875-895.











ROLE OF GERM LINE IN EPIGENETIC TRANSGENERATIONAL INHERITANCE



Summary

- Transient Embryonic Exposure Effects Generational Phenotype
 - Sex Determination Period/Germline
 - Fetal/Early Life Exposure Basis of Adult Disease
 - Disease (Testis/Prostate/Kidney/Immune/Cancer)
 - Brain/Behavior (Sexual Selection/Anxiety/Stress)
 - Somatic Cell versus Germline

EPIGENETIC AND GENETIC CASCADE OF EVENTS INVOLVED IN DEVELOPMENT






Summary

- Transient Embryonic Exposure Effects Generational Phenotype
 - Sex Determination Period/Germline
 - Somatic Cell versus Germline
 - Fetal/Early Life Exposure Basis of Disease
 - Disease (Testis/Prostate/Kidney/Immune/Cancer)
 - Brain/Behavior (Sexual Selection/Anxiety/Stress)
- Epigenetic Transgenerational Inheritance
 - Toxicology Environmental Exposures
 - Permanent Re-Program (Imprint) Germ-Line
 - Disease Etiology
 - Evolutionary Biology

Evolutionary Biology

- Mechanism-Adaptation (ie DNA mutations)
- Problem- Timing
- Environment- Influence (?How)
- Determinant- Sexual Selection

Collaboration- David Crews, UT Austin



Role of Epigenetics in the Speciation of Darwin Finches and Evolutionary Biology

Goal-



Demonstrate the role of environment and epigenetics as critical molecular factors in evolutionary biology.



Objectives-

-Use Darwin Finches to determine the genetic and epigenetic differences between variety species (ie For, Ful, Spa, Par, Cra).

-Use Chromosomal Genomic Hybridization (CGH) to associate copy number variation between species (genetic differences).

-Use Methylated DNA Immunoprecipitation followed by tiling array (MeDIP-Chip) to associate epimutations between species (epigenetic differences).





Transgenerational Phenotype

	DNA Mutation	Epigenetic Mutation
Frequency -	<0.01% (Hot Spot 1-5%)	High (30-100%)
Reproducible-	Random Event	Highly Reproducible
Genetics-	Mendelian (decline frequency generationally)	Non-Mendelian

The Big Blue[®] procedure

- Treat Big Blue rodent**
 - The proprietary mouse or rat of the Big Blue mutagenesis assay system has a lambda phage vector integrated into its genome. This vector contains a target site for mutagenesis.
 - Deliver the test compound by any route of administration to the mouse or rat.
- Isolate tissue and extract DNA**
 - Isolate any tissue of interest and extract the DNA using a quick and simple protocol.
- Add Transpack packaging extract**
 - Add the Transpack lambda DNA packaging extract to automatically excise the lambda vector target and package it into a lambda head.
- Plate to yield plaques**
 - Infect host DNA with packaged DNA and plate on indicator agar plates to yield plaques.
 - Each plaque represents a target gene.
 - Count blue mutant plaques versus the total number of plaques to determine mutant frequency.
 - Further characterize mutation if desired.


Mutant (blue) plaque among non-mutant (clear) plaques

DNA Sequence Mutation Rate

Control F1 Male	Sperm	2.08± 0.60
	Kidney	2.34± 0.69
Vinclozolin F1 Male	Sperm	1.81± 0.36
	Kidney	1.88± 0.36

•Big Blue Rat mutation frequencies in treated and control F1 generation animal sperm and kidney DNA samples. No statistical difference with a P>0.4 for kidney and P>0.5 for sperm


Tertiary Epimutations - A Novel Aspect of Epigenetic Transgenerational Inheritance Promoting Genome Instability
 PLoS One. 2016 Dec 19;11(12):e0168038.
 McCarrey JR, Lehe JD, Raju SS, Wang Y, Nilsson EE, Skinner MK.





Transgenerational Phenotype


	DNA Mutation	Epigenetic Mutation
Frequency -	<0.01% (Hot Spot 1-5%)	High (30-100%)
Reproducible-	Random Event	Highly Reproducible
Genetics-	Mendelian (decline frequency generationally)	Non-Mendelian

Unified Theory of Evolution: A Neo-Lamarckian Concept that Facilitates Neo-Darwinian Evolution






Exposure	Pathology	Reference
Toxicants		
Vinclozolin	Testis, Prostate, Kidney Disease, Tumors, Immune Gender Specific Changes in Anxiety-like Behavior Immune and Reproductive	Anway, et al., 2005 [3]; 2006 [32] Skinner, et al., 2008 [33] Nilsson, et al., 2008 [34]
Methoxychlor	Testis, Kidney, Ovary, Obesity	Anway, et al., 2005 [3], Manikkam, 2014 [35]
Permethrin/DEET	Prostate, Kidney Disease	Manikkam, et al., 2012 [36]
Dioxin	Prostate, Kidney, Fertility, Pregnancy	Manikkam, et al., 2012 [37] Bruner-Tran, 2011 [38]
BPA/Phthalates	Prostate, Kidney, Obesity	Manikkam et al., 2013 [39]
Hydrocarbon Mixture (Jet Fuel)	Prostate, Kidney, Obesity, Immune and Reproduction	Tracey et al., 2013 [40]
Vinclozolin, Permethrin/DEET, Plastics, Dioxin, Jet Fuel	Polycystic Ovaries, Reduced Primordial Follicle Pool	Nilsson et al., 2012 [41]
DDT	Obesity, Kidney, Testis	Skinner, et al., 2013 [5]
Phthalate	Testis and Spermatogonial Stem Cell	Doyle, et al., 2013 [42]
Tributyltin	Obesity and Adipose Cell	Chamorro-Garcia, et al., 2013 [43]
BPA	Social Behavior, Implantation, Litter Size, Sperm	Wolstenholme, et al., 2012 [44]; Salian, et al., 2009 [45]
Others		
Caloric Restriction	Cardiovascular Mortality	Bygren, et al., 2014 [46]
High Fat Diet	Growth and Insulin Sensitivity	Dunn and Bale, 2011 [6]
Folate	Congenital Malformations	Padmanabhan, et al., 2013 [47]
Drought	DNA Methylation Changes	Zheng, et al., 2013 [7]
Heat/Salt	Flowering and Salt Tolerance	Suter and Widmer, 2013 [48]
Prediabetes	Glucose Tolerance and Insulin Sensitivity	Wei, et al., 2014 [49]
Smoking	Abnormal Pulmonary Function	Rehan, 2013 [50]
Alcohol	Endocrine and Neuronal Function	Govorko, 2012 [51]
Heat Stress	Increased Hsp70 Production and Tolerance to Heat Stress	Norouzitallab, et al., 2014 [8]




Male germline transmits fetal alcohol adverse effect on hypothalamic proopiomelanocortin gene across generations.

Govorko D, et al. (2012) *Biol Psychiatry*. 1:72(5):378-88.




Progressive, transgenerational changes in offspring phenotype and epigenotype following nutritional transition.

Burdge GC, et al. (2011) *PLoS One*. 6(11):e28282.



Transgenerational epigenetic inheritance of longevity in Caenorhabditis elegans.

Greer EL, et al. (2011) *Nature*. 19:479(7373):365-71.



Transgenerational epigenetic inheritance in plants.



Hauser MT, et al. (2011) *Biochim Biophys Acta*. 2011 Aug;1809(8):459-68.

Table 1
Impact of variations in the epigenetic status on plant growth and development

Plant species	Induction	Epigenetic status	Effect	Reference
Oryza sativa DWARF1		DNA hypomethylation	Growth	[205,210]
Wheat Triticum aestivum	Glutenin gene	DNA methylation	Induced growth	[211]
Triticale		Hypomethylation	Transgene silencing	[212]
Tobacco Nicotiana glauca	Transgenesis	Loss of expression of transgene, hypermethylation of 5'-DNA	Transgene silencing	[213]
Arabidopsis Petunia	Transgenesis	Loss of expression of transgene, hypermethylation of 5'-DNA	Hypermethyl sensitivity	[214]
Arabidopsis AGAMOUS	met1, ddm1 background	DNA hypermethylation	Floral organ development, patterning	[216,217]
Silene latifolia	5-azacytidine	DNA hypomethylation	Sex change to androecium/gynoecium	[218]
Arabidopsis SUPERMAN/clark kent	met1, ddm1 background	DNA hypermethylation	Floral organ development, patterning	[219,220]
Arabidopsis FRI (phosphoribosyltransferase isomerase)	Natural	Ecotypes with hypermethylated multiple FRI genes, result in the silencing of the unlinked FRI genes	Floral development	[221-224]
Linaria vulgaris LYCLODEA	Natural	Cytosine DNA methylation of glabrenin genes	Floral development	[36]
Arabidopsis FLOWERING WAGENINGEN (FWA)	Natural	Loss of methylation, ectopic expression of FWA	Late flowering	[225]
Arabidopsis BALKFRI (constitutive expressor of FR genes 1)	Natural	dm1 background	Overexpression of CFR1	[226]
Arabidopsis Flowering Receptor Locus (FLC)	Natural	H3K9 and H3K27 dimethylation at FLC locus after vernalization	Flower induction after vernalization	[227,228]
Rax Linum sativum	5-azacytidine	DNA hypomethylation, chromatin remodeling	Number of leaves, flowering age, stem height	[229,230]
Rax Linum sativum	Nutrient conditions, heat	DNA methylation, presence of insertion element	Biomass, growth	[136]
Tomato	Natural	SBP-Box gene	Fruit ripening	[231]
Arabidopsis	Natural	Non-LTR retrotransposon Sadhu is differentially expressed in natural accessions as Col (active), Col and Ler (inactive)	Fruit ripening	[232]
Arabidopsis BONSAI	dm1 background	DNA hypomethylation and silencing of APC13/BNS gene by nearby a long interspersed	Growth	[233]
Melon Cucumis melo	Natural transposon insertion	Spreading of DNA methylation to WIP1 promoter	Sex determination, expression of WIP1 leads to carpel abortion	[234]
Wild potato Solanum tuberosum	Species of hybrid origin	Phenotypic instability of flower development	Flower development	[235]

Environmental heat and salt stress induce transgenerational phenotypic changes in Arabidopsis thaliana.

Suter L, Widmer A. (2013) PLoS One. 9;8(4):e60364.



Exposure to the environmental endocrine disruptor TCDD and human reproductive dysfunction: Translating lessons from murine models

Reprod Toxicol. 2017 Mar;68:59-71.

Bruner-Tran KL, Gnecco J, Ding T, Glone DR, Pensabene V, Osteen KG.

Pregnant Dam (F0 generation)



F1 fetus (contains the F2 germ cells)

Cumulative overview of reproductive characteristics in control and toxicant exposed female mice.^a

Mouse History	Pregnancy Rate	Term Delivery ^b	Adenomyosis	Methylation of PR ^c
Control	95-100%	100%	0%	0%
F1	46	37	70	66
F3	50	60	56	33

^a Data compiled from references [10,101,102].


^b Delivery was considered term when occurring E19-E21 (plug date is considered E0.5).

^c Whole uteri collected from non-pregnant mice at estrus.

Transgenerational Effects of Di-(2-Ethylhexyl) Phthalate (DEHP) on Stress Hormones and Behavior.

Quinnies KM, Doyle TJ, Kim KH, Rissman EF.

Endocrinology. 2015 Sep;156(9):3077-83.



Mean (\pm SEM) corticosterone levels (ng/mL) measured in plasma from 150-mg/kg DEHP-female and control F3 mice.

(A) Under baseline conditions DEHP-lineage mice tended to have lower corticosterone levels than control females (#P = .10). (B) After 15 minutes of restraint stress, DEHP-lineage females had lower corticosterone than control females; *, P < .05 (control-lineage females stressed, n = 10; control-lineage females baseline, n = 9; DEHP-lineage females stressed, n = 8; DEHP-lineage females baseline, n = 10; control-lineage males stressed, n = 6; control-lineage males baseline, n = 6; DEHP-lineage males stressed, n = 8; DEHP-lineage males baseline, n = 7).


The history of Distilbene® (Diethylstilbestrol) told to grandchildren--the transgenerational effect.

Fénichel P, Brucker-Davis F, Chevalier N.

Ann Endocrinol (Paris). 2015 Jul;76(3):253-9.




Table 1
In utero exposure to Distilbene®: historical.

1936	DES Synthesis. Dodd and Larson, [22]
1946	Distilbene® advocated for spontaneous miscarriage and premature labour. 5 to 125 mg/day from 6WA to 35WA
1953	Randomised double-blind study n = 840 not superior to placebo. Dieckman, [3]
1970	7 cases of clear cell adenocarcinoma (CCA) of the vagina and cervix in very young women exposed to DES in utero. Herbst [4]
1972	US ban by the FDA
1975	First case of CCA described in France
1976	Disappears from medical references as treatment for miscarriage in France
1977	Excluded for pregnancy
1977	Uterine defects related to DES. Kauffmann, [23]
1981	Pregnancy complications. Paplernik, [24]
1990	Genital abnormalities described in men
2006	RR v 3 breast cancer after 50 years of age in second generation. Palmer, [1]
2002	Third generation hypospadias. Klip, [2]
2009	Epigenetic mechanisms of foetal programming after exposure to DES hypermethylation of HOX 10 gene in uterine cancer. Bromer, [25]



Fetal Basis of Adult Onset Disease

Embryo	Most Sensitive
Postnatal	Sensitive
Pubertal	Less Sensitive
Adult	Insensitive
Aged Adult	Most Disease



Schedule/Lecture Outline –

January	14 & 16	Week 1	Systems Biology Introduction
	21 & 23	Week 2	Molecular/ Cellular/ Reproduction Systems
	28 & 30	Week 3	Sex Determination Systems
February	4 & 6	Week 4	Male Reproductive Tract Development & Function
	11 & 13	Week 5	Female Reproductive Tract Development & Function
	18 & 20	Week 6	Gonadal Developmental Systems Biology
	25 & 27	Week 7	Testis Systems Biology
March	3 & 5	Week 8	Ovary Systems Biology
	10 & 12	Week 9	Epigenetics and Transgenerational Gonadal Disease
	16 – 20	Week 10	Spring Break
	24 & 26	Week 11	Gametogenesis/ Stem Cells/ Cloning
	31 & 2	Week 12	Hypothalamus-Pituitary Development & Function
April	7 & 9	Week 13	Reproductive Endocrinology Systems
	14 & 16	Week 14	Fertilization & Implantation Systems
	21 & 23	Week 15	Fetal Development & Birth Systems
	28 & 30	Week 16	Assisted Reproduction/Contraception
May	5 & 7	Week 17	Exam or Grant Review