

Spring 2026 – Systems Biology of Reproduction
Lecture Outline – Hypothalamus-Pituitary Development & Function
Eric Nilsson – Biol 475/575
10:35-11:50 am, Tuesday & Thursday
March 31, 2026
Week 12

Hypothalamus-Pituitary Development & Function

Cell Biology

Structure / Lobes and Development
Cell Populations and Hormones
Regulators and Mutations

Hormones

Growth Hormone / Receptors / GHRH
Prolactin / Development
Opiomelanocortin

Gonadotropins

GnRH / Pulsitive Secretion
GnRH Actions / Signaling
LH/FSH Pulsitive Secretion/Menstrual Cycle

Regulation of Development

Cyclisity / Estrous Cycle / Circadian Systems

Required Reading

de Kretser, et al. (2018) Hypothalamic Pituitary Testis Axis. In: Encyclopedia of Reproduction (Second Edition). Volume 1, Pages 180-183.

Padmanabhan, et al. (2018) Hypothalamus–Pituitary–Ovary Axis. In: Encyclopedia of Reproduction (Second Edition). Volume 2, Pages 121-129.

REFERENCES

Sousa-Vidal ÉK, Henrique G, da Silva REC, Serrano-Nascimento C. Intrauterine exposure to di(2-ethylhexyl) phthalate (DEHP) disrupts the function of the hypothalamus-pituitary-thyroid axis of the F1 rats during adult life. Front Endocrinol (Lausanne). 2023 Jan 13;13:995491.

- Ignatiuk V, Izvol'skaia M, Sharova V, Zakharova L. Disruptions in Hypothalamic-Pituitary-Gonadal Axis Development and Their IgG Modulation after Prenatal Systemic Inflammation in Male Rats. *Int J Mol Sci*. 2023 Feb 1;24(3):2726.
- Calcaterra V, Magenes VC, Tagi VM, et al. Association between Vitamin D Levels, Puberty Timing, and Age at Menarche. *Children (Basel)*. 2023 Jul 19;10(7):1243.
- Wang D, Tang Y, Wang Z. Role of sphingolipid metabolites in the homeostasis of steroid hormones and the maintenance of testicular functions. *Front Endocrinol (Lausanne)*. 2023 Mar 17;14:1170023.
- Aasif A, Alam R, Ahsan H, Khan MM, Khan A, Khan S. The role of kisspeptin in the pathogenesis of a polycystic ovary syndrome. *Endocr Regul*. 2023 Dec 21;57(1):292-303.
- Harding L, Schultz IR, Young G, Swanson P. Salmonid Pituitary Cells as a Test System for Identifying Endocrine-Disrupting Compounds. *Environ Toxicol Chem*. 2023 Aug;42(8):1730-1742.
- Birzniece V, Ho KKY. MECHANISMS IN ENDOCRINOLOGY: Paracrine and endocrine control of the growth hormone axis by estrogen. *Eur J Endocrinol*. 2021 May 6;184(6):R269-R278.
- Zmijewska A, Czelejewska W, Drzewiecka EM, Franczak A. Effect of kisspeptin (KISS) and RFamide-related peptide-3 (RFRP-3) on the synthesis and secretion of FSH in vitro by pituitary cells in pigs. *Theriogenology*. 2021 Sep 1;171:72-84.
- Rotgers E, Nicol B, Rodriguez K, Rattan S, Flaws JA, Yao HH-C. Constitutive expression of Steroidogenic factor-1 (NR5A1) disrupts ovarian functions, fertility, and metabolic homeostasis in female mice. *FASEB J*. 2021 Aug;35(8):e21770.
- Akin L, Rizzoti K, Gregory LC, Corredor B, et al. Pathogenic variants in RNPC3 are associated with hypopituitarism and primary ovarian insufficiency. *Genet Med*. 2022 Feb;24(2):384-397.
- Scagliotti V, Esse RCF, Willis TL, et al. Dynamic Expression of Imprinted Genes in the Developing and Postnatal Pituitary Gland. *Genes (Basel)*. 2021 Mar 30;12(4):509.
- Matsumoto R, Takahashi Y. Human pituitary development and application of iPSCs for pituitary disease. *Cell Mol Life Sci*. 2021 Mar;78(5):2069-2079.
- Laporte E, Vennekens A, Vankelecom H. Pituitary Remodeling Throughout Life: Are Resident Stem Cells Involved? *Front Endocrinol (Lausanne)*. 2021 Jan 29;11:604519.
- Ozaki H, Suga H, Arima H. Hypothalamic-pituitary organoid generation through the recapitulation of organogenesis. *Dev Growth Differ*. 2021 Feb;63(2):154-165.
- Cimino I, Kim H, Tung YCL, et al. Activation of the hypothalamic-pituitary-adrenal axis by exogenous and endogenous GDF15. *Proc Natl Acad Sci U S A*. 2021 Jul 6;118(27):e2106868118.
- Salvio G, Martino M, Giancola G, et al. Hypothalamic-Pituitary Diseases and Erectile Dysfunction. *J Clin Med*. 2021 Jun 9;10(12):2551.
- Kałużna M, Budny B, Rabijewski M, et al. Defects in GnRH Neuron Migration/Development and Hypothalamic-Pituitary Signaling Impact Clinical Variability of Kallmann Syndrome. *Genes (Basel)*. 2021 Jun 5;12(6):868.
- Bosch E, Alviggi C, Lispi M, et al. Reduced FSH and LH action: implications for medically assisted reproduction. *Hum Reprod*. 2021 May 17;36(6):1469-1480.

- Allensworth-James M, Banik J, Odle A, et al. Control of the Anterior Pituitary Cell Lineage Regulator POU1F1 by the Stem Cell Determinant Musashi. *Endocrinology*. 2021 Mar 1;162(3):bqaa245.
- Camilletti MA, Mayer JM, Vishnopolka SA, Perez-Millan MI. From Pituitary Stem Cell Differentiation to Regenerative Medicine. *Front Endocrinol (Lausanne)*. 2021 Jan 19;11:614999.
- Toews JNC, Hammond GL, Viau V. Liver at the nexus of rat postnatal HPA axis maturation and sexual dimorphism. *J Endocrinol*. 2021 Jan;248(1):R1-R17.
- Tenuta M, Carlomagno F, Cangiano B, et al. Somatotrophic-Testicular Axis: A crosstalk between GH/IGF-I and gonadal hormones during development, transition, and adult age. *Andrology*. 2021 Jan;9(1):168-184.
- Ni Y, Chen Q, Cai J, Xiao L, Zhang J. Three lactation-related hormones: Regulation of hypothalamus-pituitary axis and function on lactation. *Mol Cell Endocrinol*. 2021 Jan 15;520:111084.
- Araki T, Tone Y, Yamamoto M, et al. Two Distinctive POMC Promoters Modify Gene Expression in Cushing Disease. *J Clin Endocrinol Metab*. 2021 Aug 18;106(9):e3346-e3363.
- Ogawa S, Parhar IS. Heterogeneity in GnRH and kisspeptin neurons and their significance in vertebrate reproductive biology. *Front Neuroendocrinol*. 2022 Jan;64:100963.
- Maione L, Bouvattier C, Kaiser UB. Central precocious puberty: Recent advances in understanding the aetiology and in the clinical approach. *Clin Endocrinol (Oxf)*. 2021 Oct;95(4):542-555.
- Tan S, Zhou Y, Zhao H, et al. Comprehensive transcriptome analysis of hypothalamus reveals genes associated with disorders of sex development in pigs. *J Steroid Biochem Mol Biol*. 2021 Jun;210:105875.
- Wen Y-L, Guo X-F, Ma L, Zhang X-S, et al. The expression and mutation of BMPR1B and its association with litter size in small-tail Han sheep (*Ovis aries*). *Arch Anim Breed*. 2021 May 28;64(1):211-221.
- Kim J-H, Park I, Shin HR, Rhee J, et al. The hypothalamic-pituitary-gonadal axis controls muscle stem cell senescence through autophagosome clearance. *J Cachexia Sarcopenia Muscle*. 2021 Feb;12(1):177-191.
- Sen A, Hoffmann HM. Role of core circadian clock genes in hormone release and target tissue sensitivity in the reproductive axis. *Mol Cell Endocrinol*. 2020 Feb 5;501:110655.
- Gregory LC, Dattani MT. The molecular basis of congenital hypopituitarism and related disorders. *J Clin Endocrinol Metab*. 2019 Nov 8. [Epub ahead of print]
- Yang N, Li T, Cheng J, Tuo Q, Shen J. Role of apelin/APJ system in hypothalamic-pituitary axis. *Clin Chim Acta*. 2019 Dec;499:149-153.
- Bao AM, Swaab DF. The human hypothalamus in mood disorders: The HPA axis in the center. *IBRO Rep*. 2018 Dec 14;6:45-53.
- Dardente H, Wood S, Ebling F, Sáenz de Miera C. An integrative view of mammalian seasonal neuroendocrinology. *J Neuroendocrinol*. 2019 May;31(5):e12729.
- Feldt-Rasmussen U, Klose M, Benvenga S. Interactions between hypothalamic pituitary thyroid axis and other pituitary dysfunctions. *Endocrine*. 2018 Dec;62(3):519-527.

- Ellsworth BS, Stallings CE. Molecular Mechanisms Governing Embryonic Differentiation of Pituitary Somatotropes. *Trends Endocrinol Metab.* 2018 Jul;29(7):510-523.
- Acevedo-Rodriguez A, Kauffman AS, Cherrington BD, Borges CS, Roepke TA, Laconi M. Emerging insights into hypothalamic-pituitary-gonadal axis regulation and interaction with stress signalling. *J Neuroendocrinol.* 2018 Oct;30(10):e12590.
- Ahmad MH, Fatima M, Mondal AC. Role of Hypothalamic-Pituitary-Adrenal Axis, Hypothalamic-Pituitary-Gonadal Axis and Insulin Signaling in the Pathophysiology of Alzheimer's Disease. *Neuropsychobiology.* 2019;77(4):197-205.
- Newton CL, Riekert C, Millar RP. Gonadotropin-releasing hormone analog therapeutics. *Minerva Ginecol.* 2018 Oct;70(5):497-515.
- Edwards W, Raetzman LT. Complex integration of intrinsic and peripheral signaling is required for pituitary gland development. *Biol Reprod.* 2018 Sep 1;99(3):504-513.
- Bhattacharya I1,2, Sen Sharma S2, Majumdar SS. Pubertal orchestration of hormones and testis in primates. *Mol Reprod Dev.* 2019 Nov;86(11):1505-1530.
- Ongaro L, Schang G, Ho CC, Zhou X, Bernard DJ. TGF- β Superfamily Regulation of Follicle-Stimulating Hormone Synthesis by Gonadotrope Cells: Is There a Role for Bone Morphogenetic Proteins? *Endocrinology.* 2019 Mar 1;160(3):675-683.
- Yvinec R, Crépieux P, Reiter E, Poupon A, Clément F. Advances in computational modeling approaches of pituitary gonadotropin signaling. *Expert Opin Drug Discov.* 2018 Sep;13(9):799-813.
- Chourpiliadi C, Paparodis R. Physiology, Pituitary Issues During Pregnancy. *StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020-. 2019 Dec 13.*
- Bagath M, Krishnan G, Devaraj C, Rashamol VP, Pragna P, Lees AM, Sejian V. The impact of heat stress on the immune system in dairy cattle: A review. *Res Vet Sci.* 2019 Oct;126:94-102.
- Kota AS1, Ejaz S1. Precocious Puberty. *StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020-. 2019 Nov 18.*
- Sansone A, Kliesch S, Isidori AM, Schlatt S. AMH and INSL3 in testicular and extragonadal pathophysiology: what do we know? *Andrology.* 2019 Mar;7(2):131-138.
- Le Tissier P, Fiordelisio Coll T, Mollard P. The Processes of Anterior Pituitary Hormone Pulse Generation. *Endocrinology.* 2018 Oct 1;159(10):3524-3535.
- Mancini A, Howard SR, Cabrera CP, et al. EAP1 regulation of GnRH promoter activity is important for human pubertal timing. *Hum Mol Genet.* 2019 Apr 15;28(8):1357-1368.
- Yelamanchi SD, Tyagi A, Mohanty V, et al. Proteomic Analysis of the Human Anterior Pituitary Gland. *OMICS.* 2018 Dec;22(12):759-769.
- Alexandre PA, Naval-Sanchez M, et al. Systems Biology Reveals NR2F6 and TGFB1 as Key Regulators of Feed Efficiency in Beef Cattle. *Front Genet.* 2019 Mar 22;10:230.
- Zhao Y, Singh C, Prober DA, Wayne NL. (2016) Morphological and Physiological Interactions Between GnRH3 and Hypocretin/Orexin Neuronal Systems in Zebrafish (*Danio rerio*). *Endocrinology.*;157(10):4012-4020.
- León-Olea M, Martyniuk CJ, Orlando EF, Ottinger MA, Rosenfeld C, Wolstenholme J, Trudeau VL. (2014) Current concepts in neuroendocrine disruption. *Gen Comp Endocrinol.* 203:158-173.

- Lovejoy DA, Barsyte-Lovejoy D. (2013) Systems approaches to genomic and epigenetic inter-regulation of peptide hormones in stress and reproduction. *Prog Biophys Mol Biol.* 113(3):375-86.
- Tonsfeldt KJ, Chappell PE. (2012) Clocks on top: the role of the circadian clock in the hypothalamic and pituitary regulation of endocrine physiology. *Mol Cell Endocrinol.* 349(1):3-12.
- Chengalvala MV, Meade EH Jr, Cottom JE, Hoffman WH, Shanno LK, Wu MM, Kopf GS, Shen ES. (2006) Regulation of female fertility and identification of future contraceptive targets. *Curr Pharm Des.* 12(30):3915-28.
- Hsieh M, Conti M. (2005) G-protein-coupled receptor signaling and the EGF network in endocrine systems. *Trends Endocrinol Metab.* 16(7):320-6.
- Cook RW, Thompson TB, Jardetzky TS, Woodruff TK. (2004) Molecular biology of inhibin action. *Semin Reprod Med.* 22(3):269-76.
- Tena-Sempere M, Barreiro ML. (2002) Leptin in male reproduction: the testis paradigm. *Mol Cell Endocrinol.* 2002 Feb 25;188(1-2):9-13.
- Amory JK, Bremner WJ. (1998) The use of testosterone as a male contraceptive. *Baillieres Clin Endocrinol Metab.* 12(3):471-84.
- Walker DM, Gore AC. (2017) Epigenetic impacts of endocrine disruptors in the brain. *Front Neuroendocrinol.* Jan;44:1-26.
- Maggi R, Zasso J, Conti L. (2015) Neurodevelopmental origin and adult neurogenesis of the neuroendocrine hypothalamus. *Front Cell Neurosci.* 6;8:440.
- Sookoian S, Gianotti TF, Burgueño AL, Pirola CJ. (2013) Fetal metabolic programming and epigenetic modifications: a systems biology approach. *Pediatr Res.* 73(4 Pt 2):531-42.
- Stevenson TJ, Hahn TP, MacDougall-Shackleton SA, Ball GF. (2012) Gonadotropin-releasing hormone plasticity: a comparative perspective. *Front Neuroendocrinol.* 33(3):287-300.
- Horseman ND, Gregerson KA. (2013) Prolactin actions. *J Mol Endocrinol.* 19;52(1):R95-106.
- Stevenson TJ, Hahn TP, MacDougall-Shackleton SA, Ball GF. (2012) Gonadotropin-releasing hormone plasticity: a comparative perspective. *Front Neuroendocrinol.* 33(3):287-300.
- Lovejoy DA, Michalec OM, Hogg DW, Wosnick DI. (2017) Role of elasmobranchs and holocephalans in understanding peptide evolution in the vertebrates: Lessons learned from gonadotropin releasing hormone (GnRH) and corticotropin releasing factor (CRF) phylogenies. *Gen Comp Endocrinol.* 2017 Sep 19. pii: S0016-6480(17)30489-6. doi: 10.1016/j.ygcen.2017.09.013. [Epub ahead of print]
- Jadhao AG, Pinelli C, D'Aniello B, Tsutsui K. (2017) Gonadotropin-inhibitory hormone (GnIH) in the amphibian brain and its relationship with the gonadotropin releasing hormone (GnRH) system: An overview. *Gen Comp Endocrinol.* 2017 240:69-76.
- van der Veen DR, Riede SJ, Heideman PD, Hau M, van der Vinne V, Hut RA. (2017) Flexible clock systems: adjusting the temporal programme. *Philos Trans R Soc Lond B Biol Sci.* 19;372(1734).
- Scarpa JR, Jiang P2, Losic B, Readhead B, Gao VD, Dudley JT, Vitaterna MH, Turek FW, Kasarskis A. (2016) Systems Genetic Analyses Highlight a TGF β -FOXO3

- Dependent Striatal Astrocyte Network Conserved across Species and Associated with Stress, Sleep, and Huntington's Disease. *PLoS Genet.* 8;12(7):e1006137.
- Seon S, Jeon D, Kim H, Chung Y, Choi N, Yang H. (2017) Testosterone Regulates NUCB2 mRNA Expression in Male Mouse Hypothalamus and Pituitary Gland. *Dev Reprod.* 21(1):71-78.
- Nikmard F, Hosseini E, Bakhtiyari M, Ashrafi M, Amidi F, Aflatoonian R. (2017) Effects of melatonin on oocyte maturation in PCOS mouse model. *Anim Sci J.* 2017 Apr;88(4):586-592.
- Tsutsui K, Osugi T, Son YL, Ubuka T. (2017) Review: Structure, function and evolution of GnIH. *Gen Comp Endocrinol.* 2017 Jul 25. pii: S0016-6480(17)30314-3. doi: 10.1016/j.ygcen.2017.07.024. [Epub ahead of print]
- Kühnen P, Handke D, Waterland RA, et al. (2016) Interindividual Variation in DNA Methylation at a Putative POMC Metastable Epiallele Is Associated with Obesity. *Cell Metab.* 2016 Sep 13;24(3):502-509.
- Calligaris D, Feldman DR, Norton I, et al (2015) MALDI mass spectrometry imaging analysis of pituitary adenomas for near-real-time tumor delineation. *Proc Natl Acad Sci U S A.* 112(32):9978-83.
- Zordan MA, Sandrelli F. (2015) Circadian Clock Dysfunction and Psychiatric Disease: Could Fruit Flies have a Say? *Front Neurol.* 20;6:80.
- Zhang F, Tanasa B, Merkurjev D, et al. (2015) Enhancer-bound LDB1 regulates a corticotrope promoter-pausing repression program. *Proc Natl Acad Sci U S A.* 112(5):1380-5.
- Perry JR, Day F, Elks CE, Sulem P, et al. (2014) Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature.* 2;514(7520):92-7.
- León-Olea M, Martyniuk CJ, Orlando EF, et al. (2014) Current concepts in neuroendocrine disruption. *Gen Comp Endocrinol.* 203:158-73.
- Ernst DK, Lynn SE, Bentley GE. (2016) Differential response of GnIH in the brain and gonads following acute stress in a songbird. *Gen Comp Endocrinol.* 1;227:51-7.
- Haddad-Tóvolli R, Paul FA, Zhang Y, et al. (2015) Differential requirements for Gli2 and Gli3 in the regional specification of the mouse hypothalamus. *Front Neuroanat.* 25;9:34.
- Merkle FT, Maroof A, Wataya T, et al. (2015) Generation of neuropeptidergic hypothalamic neurons from human pluripotent stem cells. *Development.* 142(4):633-43.
- Ikegami K, Atsumi Y, Yorinaga E, Ono H, (2015) Low temperature-induced circulating triiodothyronine accelerates seasonal testicular regression. *Endocrinology.* 156(2):647-59.
- Liu H, Yang Y, Zhang L, Liang R, (2014) Basic fibroblast growth factor promotes stem Leydig cell development and inhibits LH-stimulated androgen production by regulating microRNA expression. *J Steroid Biochem Mol Biol.* 144 Pt B:483-91.
- Hoshino S, Kurotani R, Miyano Y, (2014) Macrophage colony-stimulating factor induces prolactin expression in rat pituitary gland. *Zoolog Sci.* 31(6):390-7.
- Choi SG, Wang Q, Jia J, et al. (2014) Growth differentiation factor 9 (GDF9) forms an incoherent feed-forward loop modulating follicle-stimulating hormone β -subunit (FSH β) gene expression. *J Biol Chem.* 6;289(23):16164-75.

- Choi YS, Lee HJ, Ku CR, et al. (2014) FoxO1 is a negative regulator of FSH β gene expression in basal and GnRH-stimulated conditions in female. *Endocrinology*. 155(6):2277-86.
- Murray PG, Higham CE, Clayton PE. (2015) 60 YEARS OF NEUROENDOCRINOLOGY: The hypothalamo-GH axis: the past 60 years. *J Endocrinol*. 226(2):T123-40.
- Plant TM. (2015) 60 YEARS OF NEUROENDOCRINOLOGY: The hypothalamo-pituitary-gonadal axis. *J Endocrinol*. 226(2):T41-54.
- Yavropoulou MP, Maladaki A, Yovos JG. (2015) The role of Notch and Hedgehog signaling pathways in pituitary development and pathogenesis of pituitary adenomas. *Hormones (Athens)*. 14(1):5-18.
- Sellix MT. (2015) Circadian clock function in the mammalian ovary. *J Biol Rhythms*. 30(1):7-19.
- Gamble KL, Berry R, Frank SJ, Young ME. (2014) Circadian clock control of endocrine factors. *Nat Rev Endocrinol*. 10(8):466-75.
- Finch CE. (2014) The menopause and aging, a comparative perspective. *J Steroid Biochem Mol Biol*. 142:132-41.
- Grattan DR. (2015) 60 YEARS OF NEUROENDOCRINOLOGY: The hypothalamo-prolactin axis. *J Endocrinol*. 226(2):T101-22.
- Bernard V, Young J, Chanson P, Binart N. (2015) New insights in prolactin: pathological implications. *Nat Rev Endocrinol*. 11(5):265-75.
- Crowley WR. (2015) Neuroendocrine regulation of lactation and milk production. *Compr Physiol*. 5(1):255-91.
- Cawley NX, Li Z, Loh YP. (2016) 60 YEARS OF POMC: Biosynthesis, trafficking and secretion of pro-opiomelanocortin-derived peptides. *J Mol Endocrinol*. 2016 Feb 15. pii: JME-15-0323. [Epub ahead of print]
- Keller-Wood M. (2015) Hypothalamic-Pituitary--Adrenal Axis-Feedback Control. *Compr Physiol*. 1;5(3):1161-82.
- Dores RM, Londraville RL, Prokop J, Davis P, Dewey N, Lesinski N. (2014) Molecular evolution of GPCRs: Melanocortin/melanocortin receptors. *J Mol Endocrinol*. 52(3):T29-42.
- Huo X, Chen D, He Y, Zhu W, Zhou W, Zhang J. (2015) Bisphenol-A and Female Infertility: A Possible Role of Gene-Environment Interactions. *Int J Environ Res Public Health*. 7;12(9):11101-16.
- Clarke IJ. (2015) Hypothalamus as an endocrine organ. *Compr Physiol*. 5(1):217-53.
- León-Olea M, Martyniuk CJ, Orlando EF, et al. (2014) Current concepts in neuroendocrine disruption. *Gen Comp Endocrinol*. 1;203:158-73.
- Halupczok J, Kluba-Szyszk A, Bidzińska-Speichert B, Knychalski B. (2015) Ovarian Hyperstimulation Caused by Gonadotroph Pituitary Adenoma--Review. *Adv Clin Exp Med*. 24(4):695-703.
- Jin JM, Yang WX. (2014) Molecular regulation of hypothalamus-pituitary-gonads axis in males. *Gene*. 551(1):15-25.
- Motta G, Allasia S, Ghigo E, Lanfranco F. (2016) Ghrelin Actions on Somatotropic and Gonadotropic Function in Humans. *Prog Mol Biol Transl Sci*. 138:3-25.

- Halupczok J, Kluba-Szyska A, Bidzińska-Speichert B, Knychalski B. (2015) Ovarian Hyperstimulation Caused by Gonadotroph Pituitary Adenoma--Review. *Adv Clin Exp Med.* 24(4):695-703.
- Ding Y, Yu J, Qu P, Ma P, Yu Z. (2015) The negative effects of chronic exposure to isoflurane on spermatogenesis via breaking the hypothalamus-pituitary-gonadal equilibrium. *Inhal Toxicol.* 27(12):621-8.
- Kauffman AS, Thackray VG, Ryan GE, et al. (2015) A Novel Letrozole Model Recapitulates Both the Reproductive and Metabolic Phenotypes of Polycystic Ovary Syndrome in Female Mice. *Biol Reprod.* 93(3):69.
- Su S, Sun X, Zhou X, Fang F, Li Y. (2015) Effects of GnRH immunization on the reproductive axis and thymulin. *J Endocrinol.* 226(2):93-102.
- Gómez JM, Penalba R, Cardoso N, et al. (2015) Exposure to a low dose of bisphenol A impairs pituitary-ovarian axis in prepubertal rats: effects on early folliculogenesis. *Environ Toxicol Pharmacol.* 39(1):9-15.
- Han X, Gu L, Xia C, et al. (2015) Effect of immunization against GnRH on hypothalamic and testicular function in rams. *Theriogenology.* 83(4):642-9.
- Kuiri-Hänninen T, Sankilampi U, Dunkel L. (2014) Activation of the hypothalamic-pituitary-gonadal axis in infancy: minipuberty. *Horm Res Paediatr.* 82(2):73-80.
- Cai S, Huang S, Hao W. (2015) New hypothesis and treatment targets of depression: an integrated view of key findings. *Neurosci Bull.* 31(1):61-74.
- Schauer C, Tong T, Petitjean H, et al. (2015) Hypothalamic gonadotropin-releasing hormone (GnRH) receptor neurons fire in synchrony with the female reproductive cycle. *J Neurophysiol.* 114(2):1008-21.
- Barile VL, Terzano GM, Pacelli C, Todini L, Malfatti A, Barbato O. (2015) LH peak and ovulation after two different estrus synchronization treatments in buffalo cows in the daylight-lengthening period. *Theriogenology.* 84(2):286-93.
- Liu X, Shi H. (2015) Regulation of Estrogen Receptor α Expression in the Hypothalamus by Sex Steroids: Implication in the Regulation of Energy Homeostasis. *Int J Endocrinol.* 2015:949085.
- Bales KL, Saltzman W. (2016) Fathering in rodents: Neurobiological substrates and consequences for offspring. *Horm Behav.* 77:249-59.
- Houlahan KE, Prokopec SD, Moffat ID, et al. (2015) Transcriptional profiling of rat hypothalamus response to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicology.* 328:93-101.
- Zhan X, Long Y. (2016) Exploration of Molecular Network Variations in Different Subtypes of Human Non-functional Pituitary Adenomas. *Front Endocrinol (Lausanne).* 10:7:13.
- Grizzi F, Borroni EM, Vacchini A, et al. (2015) Pituitary Adenoma and the Chemokine Network: A Systemic View. *Front Endocrinol (Lausanne).* 11:6:141.
- Zmora N, Stubblefield JD, Wong TT, Levavi-Sivan B, Millar RP, Zohar Y. (2015) Kisspeptin Antagonists Reveal Kisspeptin 1 and Kisspeptin 2 Differential Regulation of Reproduction in the Teleost, *Morone saxatilis*. *Biol Reprod.* 93(3):76.
- Yao K, Ge W. (2015) Differential regulation of kit ligand A (*kitlga*) expression in the zebrafish ovarian follicle cells--evidence for the existence of a cyclic adenosine 3', 5' monophosphate-mediated binary regulatory system during folliculogenesis. *Mol Cell Endocrinol.* 15:402:21-31.

- Vasilev V, Daly AF, Thiry A, et al. (2014) McCune-Albright syndrome: a detailed pathological and genetic analysis of disease effects in an adult patient. *J Clin Endocrinol Metab.* 99(10):E2029-38.
- Choi SG, Wang Q, Jia J, Pincas H, et al. (2014) Growth differentiation factor 9 (GDF9) forms an incoherent feed-forward loop modulating follicle-stimulating hormone β -subunit (FSH β) gene expression. *J Biol Chem.* 289(23):16164-75.
- León-Olea M, Martyniuk CJ, et al. (2014) Current concepts in neuroendocrine disruption. *Gen Comp Endocrinol.* pii: S0016-6480(14)00048-3.
- Kommadath A, Te Pas MF, Smits MA. (2013) Gene coexpression network analysis identifies genes and biological processes shared among anterior pituitary and brain areas that affect estrous behavior in dairy cows. *J Dairy Sci.* 96(4):2583-95.
- Navarro VM, Tena-Sempere M. (2011) Neuroendocrine control by kisspeptins: role in metabolic regulation of fertility. *Nat Rev Endocrinol.* 2011 Sep 13;8(1):40-53.
- Budry L, Lafont C, et al. (2011) Related pituitary cell lineages develop into interdigitated 3D cell networks. *Proc Natl Acad Sci U S A.* 108(30):12515-20.
- Bailey M, Silver R. (2014) Sex differences in circadian timing systems: implications for disease. *Front Neuroendocrinol.* 35(1):111-39.
- Tasker JG, Oliet SH, et al. (2012) Glial regulation of neuronal function: from synapse to systems physiology. *J Neuroendocrinol.* 24(4):566-76.
- Bentley GE, Tsutsui K, Kriegsfeld LJ. (2010) Recent studies of gonadotropin-inhibitory hormone (GnIH) in the mammalian hypothalamus, pituitary and gonads. *Brain Res.* 10;1364:62-71.
- McGuire NL, Bentley GE. (2010) Neuropeptides in the gonads: from evolution to pharmacology. *Front Pharmacol.* 9;1:114.
- Zhang D1, Xiong H, et al. (2009) Defining global neuroendocrine gene expression patterns associated with reproductive seasonality in fish. *PLoS One.* 5;4(6):e5816.
- Chimento A, Sirianni R, Casaburi I, Pezzi V. (2014) Role of Estrogen Receptors and G Protein-Coupled Estrogen Receptor in Regulation of Hypothalamus-Pituitary-Testis Axis and Spermatogenesis. *Front Endocrinol (Lausanne).* 16;5:1. eCollection 2014.
- Stevenson EL, Corella KM, Chung WC. (2013) Ontogenesis of gonadotropin-releasing hormone neurons: a model for hypothalamic neuroendocrine cell development. *Front Endocrinol (Lausanne).* 16;4:89
- Colledge WH, Doran J, Mei H. (2013) Model systems for studying kisspeptin signalling: mice and cells. *Adv Exp Med Biol.* 784:481-503.
- Atkin SD, Owen BM, et al. (2013) Nuclear receptor LRH-1 induces the reproductive neuropeptide kisspeptin in the hypothalamus. *Mol Endocrinol.* 27(4):598-605.
- Wen S, Ai W, Alim Z, Boehm U. (2010) Embryonic gonadotropin-releasing hormone signaling is necessary for maturation of the male reproductive axis. *Proc Natl Acad Sci U S A.* 14;107(37):16372-7.
- Colledge WH, Mei H, d'Anglemont de Tassigny X. (2010) Mouse models to study the central regulation of puberty. *Mol Cell Endocrinol.* 5;324(1-2):12-20.
- Bliss SP, Miller A, et al. (2009) ERK signaling in the pituitary is required for female but not male fertility. *Mol Endocrinol.* 23(7):1092-101.
- Ivell R, Heng K, Anand-Ivell R. (2014) Insulin-Like Factor 3 and the HPG Axis in the Male. *Front Endocrinol (Lausanne).* 27;5:6. eCollection 2014.

- Kumar S, Kaur G. (2013) Intermittent fasting dietary restriction regimen negatively influences reproduction in young rats: a study of hypothalamo-hypophysial-gonadal axis. *PLoS One*. 8(1):e52416.
- Brayman MJ, Pepa PA, Mellon PL. (2012) Androgen receptor repression of gonadotropin-releasing hormone gene transcription via enhancer 1. *Mol Cell Endocrinol*. 5;363(1-2):92-9.
- Perrett RM, McArdle CA. (2013) Molecular Mechanisms of Gonadotropin-Releasing Hormone Signaling: Integrating Cyclic Nucleotides into the Network. *Front Endocrinol (Lausanne)*. 20;4:180. eCollection 2013.
- Terasawa E, Kurian JR, et al. (2010) Recent discoveries on the control of gonadotrophin-releasing hormone neurones in nonhuman primates. *J Neuroendocrinol*. 22(7):630-8.
- Miller BH, Takahashi JS. (2014) Central Circadian Control of Female Reproductive Function. *Front Endocrinol (Lausanne)*. 22;4:195. eCollection 2013.
- Chimento A, Sirianni R, et al. (2014) Role of Estrogen Receptors and G Protein-Coupled Estrogen Receptor in Regulation of Hypothalamus-Pituitary-Testis Axis and Spermatogenesis. *Front Endocrinol (Lausanne)*. 16;5:1. eCollection 2014.
- Sellix MT. (2013) Clocks underneath: the role of peripheral clocks in the timing of female reproductive physiology. *Front Endocrinol (Lausanne)*. 23;4:91. eCollection 2013.
- Stevenson EL, Corella KM, Chung WC. (2013) Ontogenesis of gonadotropin-releasing hormone neurons: a model for hypothalamic neuroendocrine cell development. *Front Endocrinol (Lausanne)*. 16;4:89. eCollection 2013.
- Ratnasabapathy R, Dhillon WS. (2013) The effects of kisspeptin in human reproductive function - therapeutic implications. *Curr Drug Targets*. 14(3):365-71.
- Aluru N, Vijayan MM. (2009) Stress transcriptomics in fish: a role for genomic cortisol signaling. *Gen Comp Endocrinol*. 164(2-3):142-50.

MALE ENDOCRINOLOGY

Hypothalamic Pituitary Testis Axis

David Morritz de Kretser, Hudson Institute and Monash University, Melbourne, VIC, Australia

© 2018 Elsevier Inc. All rights reserved.

Understanding testicular function and the pathological states that result in male infertility or androgen deprivation requires knowledge of the hormonal control of the testis. The regulation of reproduction is under the control of the hypothalamic-pituitary-gonadal axis. As early as 1930, the importance of the pituitary gland in controlling reproduction as well as many other functions was recognized (Smith, 1930).

Clear evidence for the involvement of this system also emerges from the seasonal influence on reproductive processes in females and males from many species but there is less evidence for a role of season in the control of human reproduction (Popa and Fielding, 1931; Harris, 1948; Green and Harris, 1951). These observations indicate that the changing day length is detected via the eyes and result in a change in reproductive behavior. In addition, the effect of psychological stress and marked weight loss associated with anorexia are known to influence the regularity of menstrual cycles. Further, as early as 1930 the need for replacement therapy following the removal of the pituitary was recognized (Smith, 1930).

The neural pathways involved result in stimulating the hypothalamus to send signals that act to alter the secretion of hormones from the pituitary gland that influence testicular and ovarian function (Popa and Fielding, 1931; Harris, 1948; Green and Harris, 1951).

As with all endocrine glands in the body, the endocrine control of the testis is exerted via the hypothalamic-pituitary testicular axis. The pituitary gland is formed by two components, one of epithelial origin from the developing pharynx which forms the anterior pituitary or adenohypophysis and a neural down growth from the hypothalamic region of the brain that forms the neurohypophysis (Figs. 1–4); (Popa and Fielding, 1931). The latter is responsible for the secretion of oxytocin and vasopressin.

Those signals are transmitted by peptide hormones that utilize a portal system of blood vessels to carry them from the hypothalamus to the anterior part of the pituitary gland (Fig. 4). In contrast, the posterior pituitary is composed of nerve fibers arising from nerve cells which release their products into blood vessels thus entering the circulation and transporting those entities to their sites of action (Green and Harris, 1951). Harris, one of the pioneers in this field articulated the basis of our current understanding of the physiology of the hypothalamo-hypophyseal system. He used electrical stimulation to demonstrate the neural link to the pituitary gland (Green and Harris, 1951).

The mechanisms involved are the secretion of gonadotropin releasing hormone (GnRH), a decapeptide secreted by hypothalamic neurones into the hypothalamo-hypophyseal portal vascular system that conveys the secreted GnRH to stimulate the gonadotrophic hormones, follicle stimulating hormone (FSH) and luteinizing hormone (LH) secretion by the gonadotrophin secreting cells in the anterior pituitary gland (Guillemin, 1964; Gersh, 1938). In turn, FSH and LH circulate in the blood stream to stimulate spermatogenesis and testosterone secretion by the testis in the male, and ovulation and estradiol secretion by the ovary in the female.

The pulsatile nature of LH secretion clearly supports the pulsatile nature of the GnRH stimulus but the pulsatility is less evident from the pattern of FSH secretion since the longer half-life of FSH obscures the pulsatility. The pulse frequency and amplitude is set




Fig. 1 A diagram of a median section of the pituitary gland.

FLOOR OF THIRD VENTRICLE: FROM BELOW




Fig. 2 A diagram of the region of the third ventricle from below.

MEDIAN SECTION: BODY OF SPHENOID & HYPOPHYSIS




Fig. 3 A diagram of a median section through the pituitary and the body of the sphenoid bone.

CORONAL SECTION: HYPOPHYSIS CEREBRI




Fig. 4 A diagram of a coronal section of the pituitary gland and adjacent structures.

by signals from a collection of nerve cells in the hypothalamus called the arcuate nucleus. Thus external stimuli, such as the changes that occur due to seasonal alterations in photoperiod, are detected by the eyes and relayed to the hypothalamus. Therein, these visual stimuli are transduced into neural signals that are transmitted to pathways that affect the GnRH pulsatility. The latter, in turn, influence the secretion of FSH and LH by the pituitary. The early onset of pulsatile GnRH secretion results in precocious puberty whereas the failure to initiate pulsatile GnRH results in a delay or a failure to commence puberty in both males and females.

Modifications in the peptide structure of GnRH has resulted in some analogues having a higher affinity and high potency whereas other structural changes can result in agents that can bind to the receptor but fail to induce downstream signaling and can thus act as contraceptive agents.

As with most hormonal systems, the target tissue responds to the hormone and also sends feedback signals that modulate the bioactivity concerned. Thus the steroid hormones produced by the ovary and testis, estradiol and testosterone, exert a negative feedback action on the secretion of FSH and LH.

FSH also stimulates the ovary and the testis to secrete inhibin, a protein hormone that acts as its feedback regulator (Robertson et al., 1987). It is composed of two subunits termed α and β and specifically suppresses FSH and the DNA sequences for the encoding genes were cloned for bovine inhibin (Forage et al., 1986). A second protein that specifically suppresses FSH was also isolated (Robertson et al., 1987) and given the name FSH suppressing protein. The same protein was also isolated independently and given the name, follistatin, the name by which it is now known (Ying et al., 1987). Other agents such as neuropeptide Y enhance GnRH binding to its receptors and augment LH pulses. Immuno-neutralization of NPY attenuates the LH surge at ovulation. Similarly, the actions of estradiol on GnRH are complex, suppressing the magnitude in the first 36 h of exposure and then augmenting the response after 48 h.

Since the release of GnRH by the hypothalamus is pulsatile, the subsequent FSH and LH release by the pituitary is also pulsatile. The pulsatility is more evident for LH which has a shorter half-life than FSH. Through GnRH, which enables neural inputs into the regulation of the gonadotrophins, external factors such as photoperiod, nutrition, stress, infection, and inflammation can affect their secretory patterns. The arcuate nucleus, a collection of nerve cells or neurones in the hypothalamus, is responsible for setting the pulse frequency and amplitude of GnRH secretion. Thus the early onset of pulsatile secretion initiates precocious puberty and a delay in the onset results in delayed puberty. Some mutations in the GnRH receptor can negate the pulses of GnRH from influencing FSH and LH secretion and thus acting as a cause of the failure to initiate puberty.

The pathway controlling GnRH pulse frequency involves kisspeptin acting on the GPR-54 receptor on GnRH secreting neurones. In general kisspeptin and glutamate increase GnRH release and endogenous opioid peptides inhibit GnRH secretion. The actions of estrogenic steroids such as estradiol are complex with increasing levels being initially inhibitory and later stimulatory, in part dependent on the pulse frequency. As indicated above, the onset of FSH and LH secretion determines the timing of puberty in that an early onset results in precocious puberty and a failure or delay in the onset results in delayed puberty. In some instance, mutations in the genes encoding the GnRH receptor are the cause of the failure of FSH and LH secretion or result in low levels of these gonadotrophins causing delayed puberty or infertility. Mutations in the gene encoding this receptor are known to cause hypogonadotropic hypogonadism which presents in males and females as the failure of pubertal onset. NPY is a 36 amino acid protein member of the pancreatic polypeptide family and enhances GnRH binding to gonadotrophs in the pituitary gland and augments the LH response to GnRH. In the testis, NPY expressing nerve fibers are confined to the capsule of the testis and capsular blood vessels (Allen et al., 1989).

Given that FSH and LH are the agents that circulate via the blood to the ovary and the testis, they are critical components that regulate spermatogenesis in males and oogenesis and ovulation in females. In turn, the ovarian and testicular steroid hormones, estradiol and testosterone, act predominantly as negative regulators of FSH and LH. In addition, FSH stimulates the gonads to produce a hormone, inhibin that acts as a specific inhibitor of FSH secretion. There are two forms of inhibin which are dimers of two subunits, alpha (α) and beta (β_A) and beta (β_B). Inhibin A ($\alpha\beta_A$) and Inhibin B ($\alpha\beta_B$) both suppress FSH secretion. Castration of males or females results in a rapid decrease in the serum inhibin levels indicating that the gonads are the source of these proteins. In the male, the Sertoli cells of the testes are the site of production of inhibin. There is yet another protein that regulates FSH and that is called follistatin which is structurally unrelated to the activins and inhibins (Meinhardt et al., 1998). In contrast to inhibin, the levels of follistatin did not change after castration, indicating that it was also produced at other sites.

There are also proteins that exert a stimulatory influence on FSH levels. These are dimers of the inhibin β_A and β_B units which can form proteins called activin A ($\beta_A\beta_A$) and activin B ($\beta_B\beta_B$) and stimulate FSH secretion (Vale et al., 1986; Ling et al., 1986).

Follistatin acts by binding the activins with high affinity and in vivo, the follistatin-activin complex is targeted to a lysosomal degradation pathway. Follistatin also binds to heparin sulfate proteoglycans that form part of the basement membranes of many tissues throughout the body. In addition to binding and inactivating the activins, follistatin also binds another protein called myostatin which, as its name suggests, is a "repressor" of muscle mass. Thus mutations of myostatin in cattle, which render myostatin inactive, have very large muscles as seen in the Belgian Blue breed of cattle.

The amino acid sequence of the activins and follistatin from rodents and humans is very highly conserved such that they are virtually identical. This degree of conservation across such a wide range of animal species suggests that these proteins have important functions.

Although, the activins are involved in the control of FSH, they are also regulators of our body's response to infection. Within 30 min of a challenge using lipopolysaccharide (LPS), a component of the Salmonella bacterium, activin A increases significantly (Jones et al., 2007). Other data available suggests that activin A is a major regulator of our body's inflammatory response. For instance, the levels of serum activin A and activin B are markedly elevated in patients admitted to intensive care units with acute

respiratory failure caused by a variety of agents (de Kretser et al., 2013) should they remain elevated, they are indicators of a very poor outcome not just in the acute phase of the infection but also for up to a year after that hospital admission.

The outcome of inflammation in many organs is fibrosis or scarring and activin is responsible for the fibrosis and follistatin can be used to attenuate the degree of fibrosis.

While this section focuses on the male, these proteins have important regulatory functions in the female, covered in another section of this Encyclopedia of Reproduction.

Acknowledgment

The figures are based on illustrations from *Regional Anatomy Illustrated*, Churchill Livingstone 1983, a company that does not exist any longer. It was formed from a merger of E&S Livingstone (Edinburgh, Scotland) and J&A Churchill (London). This company, originally owned by Harcourt and Pearson, is now integrated into Elsevier's Health Science Division.

References

- Allen, L. G., Wilson, F. J., & MacDonald, G. J. (1989). Neuropeptide Y containing nerve fibres in rat gonads: Sex difference and development. *Biology of Reproduction*, 1989(40), 371–378.
- de Kretser, et al. (2013). Serum activin A and B levels predict outcome in patients with acute respiratory failure: A prospective cohort study. *Critical Care*, 17, R263.
- Forage, R. G., Ring, J. M., Brown, R. W., McInerney, B. V., Cobon, G. S., Gregson, R. P., Robertson, D. M., Morgan, F. S., Hearn, M. T., Findlay, J. K., & de Kretser, D. M. (1986). Cloning and sequence analysis of cDNA species coding for the two subunits of inhibin from bovine follicular fluid. *Proceedings of the National Academy of Sciences of the United States of America*, 83, 3091–3095.
- Gersh, I. (1938). Relation of histological structure to the active substances extracted from the posterior lobe of the hypophysis. *Research Publications - Association for Research in Nervous and Mental Disease*, 17, 433.
- Green, J. D., & Harris, G. W. (1951). The neurovascular link between the neurohypophysis and adenohypophysis. *The Journal of Endocrinology*, 5, 136–144.
- Guillemin, R. (1964). Hypothalamic factors releasing pituitary hormones. *Recent Progress in Hormone Research*, 20, 89–130.
- Harris, G. W. (1948). Neural control of the pituitary gland. *Physiological Reviews*, 28, 139–145.
- Jones, K. L., Mansell, A., Patella, S., Scott, B. J., Hedger, M. P., de Kretser, D. M., & Phillips, D. J. (2007). Activin is a critical component of the inflammatory response and its binding protein follistatin reduces mortality in endotoxemia. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 16239–16244.
- Ling, N., Ying, S. Y., Ueno, N., Shimasaki, S., Esch, F., Hotta, M., & Guillemin, R. (1986). Pituitary FSH is released by a heterodimer of β subunits from the two forms of inhibin. *Nature*, 321, 779–782.
- Meinhardt, A., O'Bryan, M. K., McFarlane, J. R., Lovelad, K. L., Mallidis, C., Foulds, L. M., Phillips, D. J., & de Kretser, D. M. (1998). Localisation of follistatin in the testis. *Journal of Reproduction and Fertility*, 112, 233–241.
- Popa, G., & Fielding, U. (1931). A portal circulation from pituitary to hypothalamic region. *Journal of Anatomy*, 65, 88–90.
- Robertson, D. M., Klein, R., de Vos, F., McLachlan, R. I., Wettenhall, R. E., Hearn, M. T., Burger, H. G., & de Kretser, D. M. (1987). The isolation of polypeptides with FSH suppressing activity from bFF which are structurally different to inhibin. *Biochemical and Biophysical Research Communications*, 149, 744–749.
- Smith, P. E. (1930). Hypophysectomy and replacement therapy. *American Journal of Anatomy*, 45, 205–207.
- Vale, W., Rivier, J., Vaughan, J., McClintock, R., Corrigan, A., Woo, W., Karr, D., & Spies, J. (1986). Purification and characterisation of an FSH releasing protein from porcine ovarian fluid. *Nature*, 321, 776–779.
- Ying, S. Y., Becker, A., Swanson, G., Tan, P., Ling, N., Esch, F., Ueno, N., Shimasaki, R., & Guillemin, R. (1987). Follistatin specifically inhibits pituitary follicle stimulating hormone release in vitro. *Biochemical and Biophysical Research Communications*, 149, 133–139.

Further Reading

- Robertson, D., Foulds, L., Leversha, L., Morgan, F. J., Hearn, M. T., Burger, H. G., Wettenhall, R. E., & de Kretser, D. M. (1985). Isolation of inhibin from bovine follicular fluid. *Biochemical and Biophysical Research Communications*, 126, 220–226.

Hypothalamus–Pituitary–Ovary Axis

Vasantha Padmanabhan and Muraly Puttabyatappa, University of Michigan, Ann Arbor, MI, United States
Rodolfo C Cardoso, Texas A&M University, College Station, TX, United States

© 2018 Elsevier Inc. All rights reserved.

Introduction

In female mammals, reproductive processes such as puberty, ovarian cyclicity and pregnancy require precise coordination of the hypothalamic–pituitary–ovarian (HPO) axis. The major components of the HPO axis include the hypothalamic gonadotropin-releasing hormone (GnRH) neurons, the pituitary gonadotropes and ovarian structures, such as follicles and the corpus luteum (Fig. 1). The gonadotropes respond to GnRH by releasing the gonadotropins follicle-stimulating hormone (FSH) and luteinizing hormone (LH), which in turn stimulate folliculogenesis and steroid and peptidergic hormone secretion by the ovaries. The hypothalamic, pituitary, and ovarian hormones integrate to orchestrate the different events within the reproductive cycle so they occur in a timely manner to ensure successful reproduction (see earlier review (Beshay and Carr, 2013)). In this chapter, we present an overview of the key components and hormones of the HPO axis, discuss the complex endocrine interactions that exist between the different levels of this axis, and summarize recent findings on the main factors and mechanisms that regulate the HPO axis in the female mammals.

The Hypothalamus

The hypothalamus encompasses the lower part of the lateral wall and the floor of the third ventricle, containing several localized clusters of neurons, called nucleus, that control multiple processes such as metabolism, reproduction, and behavior. The median eminence located at the base of the hypothalamus extends to the pituitary and contains neurosecretory neurons that control hormone synthesis and secretion from the pituitary.




Fig. 1 Schematic showing the components of the hypothalamic–pituitary–ovarian axis.

GnRH and Its Mode of Release

The GnRH neuron

The decapeptide, GnRH—the primary hypothalamic regulator of reproductive function, is synthesized in specialized neurons as part of a larger precursor molecule (prohormone) containing 92 amino acids. GnRH neurons originate in the olfactory placode and migrate to the hypothalamus where several hundreds of these cells can be found diffusely distributed in the anterior portion of the hypothalamus, the preoptic area (POA), and more caudally in the medial basal hypothalamus (MBH). Most GnRH neurons have two dendritic projections, which may extend remarkable distances (2–3 mm) from the cell body. The main target of hypothalamic GnRH projections is the primary plexus of the pituitary portal system in the median eminence.

The hypothalamic–pituitary portal vasculature

Upon reaching the median eminence, GnRH axons end on capillary loops that give rise to long portal veins that descend along the pituitary stalk to terminate in the pituitary sinusoids. The portal system facilitates the rapid exchange between the hypothalamus and the pituitary, requiring only minute amounts of neurohormones to stimulate release of hormones by the pituitary. The superior hypophyseal arteries form the primary capillary plexus that supplies blood to the median eminence. From this capillary system, the blood is drained in hypophyseal portal veins to the secondary plexus, a network of fenestrated sinusoid capillaries that provide blood to the anterior portion of the pituitary. This anatomical arrangement allows rapid and undiluted transport of GnRH to the pituitary gland.

GnRH biosynthesis

To date, three isoforms of GnRH (GnRH-I, GnRH-II, and GnRH-III) have been identified in humans and many others have been detected in fish, amphibians, and protochordates. GnRH-I is the classic hypothalamic peptide responsible for regulating the synthesis and secretion of gonadotropins. GnRH-II, first reported in brain tissue and since been found in other peripheral tissues, such as the uterus and ovaries is also capable of stimulating LH release. The role of GnRH-III, first identified in the lamprey, in humans remains unclear (Schneider et al., 2006).

Upon synthesis, GnRH-I gets transported to the median eminence of the hypothalamus and then released in the portal circulation in a pulsatile manner. The lifespan of GnRH-I is very short with a half-life of 2–4 min. Because of its short half-life and rapid dilution in the peripheral circulation, GnRH-I concentrations are very difficult to measure in the peripheral circulation.

GnRH pulsatility

The cellular network that elicits the pulsatile increases in GnRH secretion is referred to as the “GnRH pulse generator” and includes the synchronously firing GnRH neurons. While there is supporting evidence that GnRH pulsatility is intrinsic to GnRH neurons themselves and that extensive intercellular mechanisms orchestrate synchrony within the network, more recent observations strongly indicate that other neurons in the MBH target the GnRH neuronal network directly to drive the pulsatile release of GnRH (Fig. 2). Kisspeptin, the most potent stimulator of GnRH secretion yet discovered, appears to be an integral component of the GnRH pulse generator. Of particular interest are the neurons in the arcuate nucleus (ARC) that co-express the neuropeptides kisspeptin, neurokinin B, and dynorphin, termed KNDy neurons. In contrast to the endogenous opioid dynorphin that has inhibitory actions on GnRH release, neurokinin B is generally stimulatory to GnRH secretion. Based on the action of these neuropeptides on GnRH release and the neuroanatomical evidences that KNDy neurons communicate with GnRH neurons (particularly at the level of the median eminence), a model for GnRH pulse generation has been established. This




Fig. 2 Schematic depicting the control of the GnRH neurons in the hypothalamus. KNDy neurons in the arcuate nucleus project to GnRH cell bodies and axonal projections to modulate GnRH release into the median eminence (ME).

model proposes that GnRH pulse generation is initiated in the KNDy neuronal network in the ARC by a reciprocating interplay of stimulatory neurokinin B signals and inhibitory dynorphin inputs and the output of the pulse generator is relayed to GnRH neurons via the release of kisspeptin from axonal terminals originating from KNDy neurons (Fig. 2; Lehman et al., 2010).

The Pituitary Gland

The pituitary, also known as the master endocrine gland, is a small gland located beneath the third ventricle of the brain and above the sphenoidal sinus in a bone cavity called sella turcica. The pituitary gland contains two major lobes: anterior and posterior.

The Anterior Pituitary

Development

The anterior pituitary, also known as adenohypophysis, is a classic endocrine gland in that it is composed of secretory cells of epithelial origin supported by connective tissue rich in blood and lymphatic capillaries. The anterior pituitary is derived from the ectoderm, more specifically from the Rathke's pouch, part of the developing hard palate in the fetus. The Rathke's pouch eventually loses its connection with the pharynx, giving rise to the anterior pituitary. The pouch becomes committed to organ development several days prior to the expression of markers of individual cell types within the mature anterior pituitary gland. Between the time of organ commitment and maturation, a series of cell type-specific differentiation and proliferative events occur.

Cell types

The five cell types present in the mature anterior pituitary gland are defined based on the trophic factors that they synthesize and secrete. Corticotropes secrete adrenocorticotrophin, thyrotropes produce thyroid-stimulating hormone, somatotropes secrete growth hormone, lactotropes produce prolactin and gonadotropes secrete the gonadotropins, LH and FSH.

The Posterior Pituitary

The posterior pituitary, also known as neurohypophysis, is a grouping of axonal projections that secretes oxytocin responsible for increasing uterine contractions and vasopressin, an antidiuretic hormone.

Gonadotropes

Gonadotropes are specialized basophilic cells in the anterior pituitary that express cell surface receptors for GnRH and produce LH and FSH.

The GnRH receptor

The GnRH-I receptor is a G-protein coupled receptor that utilizes inositol triphosphate and diacylglycerol as second messenger molecules to stimulate protein kinase, release calcium ions, and cyclic adenosine monophosphate activity. While the pituitary is the classic target tissue for GnRH actions, the GnRH-I receptor is expressed in many other tissues, including ovarian follicles, pancreas, liver, heart, and the placenta. Importantly, the continuous activation of the GnRH receptor results in its desensitization thus enabling the use of GnRH agonists for treatment of reproductive disorders such as precocious puberty and endometriosis, and to prevent GnRH actions prior to ovarian stimulation cycles for in vitro fertilization. Therefore, only pulsatile GnRH administration can restore normal gonadotropin secretion in individuals lacking endogenous GnRH secretion (e.g., Kallmann syndrome).

Luteinizing hormone

LH is a glycoprotein heterodimer consisting of two subunits: α - and β -subunits. The α -subunit is common between LH, FSH, and TSH and consists of 92 amino acids. The β -subunit is distinct and hormone-specific, which allows the different biological actions of each hormone. The LH β -subunit has 120 amino acids and one to two sialic acid residues, giving it its shorter half-life of approximately 20 min. LH is secreted in a pulsatile manner except preceding ovulation when a massive release of LH (LH surge) occurs to trigger ovulation. Since LH secretion is primarily regulated by GnRH, peripheral levels of LH serve as good surrogate markers for assessing pattern of GnRH release.

Follicle-stimulating hormone

FSH is also a glycoprotein heterodimer consisting of two subunits: α - and β -subunits. The FSH β -subunit contains 118 amino acids and 5 sialic acid residues. The sialic acid content of FSH and LH is responsible for the half-life of these hormones, with greater number of sialic acid residues yielding longer half-life. FSH has a half-life of approximately 3–4 h, thus providing sustained support to facilitate growth of ovarian follicles.

Ovaries

The ovaries are the female gonads and are primarily responsible for the generation of gametes and production of hormones that regulate reproductive function. In most mammals, a pair of ovaries is present on either side of the body and is structurally

made of inner medulla and the outer cortex that is encapsulated by a connective tissue rich membrane called tunica albuginea. While the medulla contains vascular structures such as blood and lymphatic vessels, the cortex contains the follicles, which are the functional units of the ovary.

Follicular Development

An ovarian follicle contains the female gamete—oocyte, and the somatic cells, granulosa and theca cells. The number and type of somatic cells varies depending on the stage of follicular development. During the embryonic development, the primordial germ cells (PGCs) originate from the endoderm of the yolk sac and migrate into the genital ridge. These PGCs mitotically divide to form the oogonia, which later undergoes meiosis to form the oocyte. The primordial follicles are formed when the naked oocytes are surrounded by the pregranulosa cells (Fig. 3). These primordial follicles form the ovarian follicular pool or reserve and majority of these follicles remain in a quiescent state. A few follicles emerge from the reserve at specific intervals, a process termed as activation, and either undergo atresia (degeneration) or develop into primary follicles with the transformation of granulosa cells from spindle to cuboidal shaped. Follicular growth continues with granulosa cells dividing to form several layers around the oocyte leading to the development of the preantral follicle. During this developmental period, the follicle also acquires the theca cells that surround the granulosa cells. It has been suggested that follicular development up until the preantral stage occurs in a gonadotropin-independent manner, however, the role of gonadotropins cannot be ruled out because FSH receptors are present in early follicular stages.

Subsequent follicular differentiation occurs under the influence of gonadotropins with preantral follicles progressing to the antral stage with development of a fluid-filled cavity, termed the antrum. Selection of a dominant follicle from the antral follicular pool that is destined for ovulation (Fig. 3) occurs under the influence of FSH. Further maturation into Graafian or preovulatory follicle occurs predominantly under LH stimulation. Upon ovulation, the release of egg from the follicle, subsequent transformation of the follicular remnant into the endocrine structure corpus luteum is driven by LH.

Ovarian Hormones

The ovarian hormones produced under the influence of LH and FSH include steroidal and non-steroidal polypeptide hormones. Androgens and estrogens are produced by theca and granulosa cells, respectively. Under the influence of LH, theca cells take up cholesterol and convert it into 17-hydroxy-pregnenolone through the Δ^5 pathway, which is then acted on by 17-hydroxylase to synthesize the androgen androstenedione. Majority of androstenedione is aromatized into the major estrogen, estradiol, by the enzyme aromatase (CYP19) in granulosa cells, a process that occurs under the influence of FSH. In the corpus luteum, synthesis of progesterone is stimulated by LH through the Δ^4 pathway (Miller, 2008).

The non-steroidal hormones, among others, include those belonging to the transforming growth factor beta (TGF β) family namely inhibin, activin, anti-Mullerian hormone (AMH), TGF α , bone morphogenetic proteins, and growth differentiation factors. Other factors produced by the ovary include insulin-like growth factors (IGF), kit ligand, fibroblast growth factors, epidermal growth factors, and prostaglandins.

Inhibins

Inhibins are protein hormones consisting of two subunits α and β and are secreted mainly by the granulosa cells under the influence of FSH but are also produced by the pituitary gonadotropes. Two forms of inhibin have been identified, Inhibin-A and -B. The α subunits are identical, while the β -subunit differs between both forms of inhibin (denoted as β_A and β_B , respectively). Inhibin-B is predominantly secreted during the follicular phase, while the A form is secreted during the luteal phase. Inhibins function to suppress FSH secretion by the anterior pituitary.

Activins

Activins are secreted by granulosa cells and are homo- or hetero-dimers of the inhibin β subunits, with activin A and B being the homodimer of two β_A and β_B subunits, respectively and AB being the heterodimer. Activin is also produced in the pituitary and




Fig. 3 Schematic representation of the different stages in the ovarian follicular development.

stimulates FSH release. Other β forms have been identified, raising the possibility of additional homo- and hetero-dimeric forms although their role in regulation of gonadotropin secretion is still being ascertained (Wijayarathna and De Kretser, 2016).

Follistatin

Follistatin is a monomeric glycoprotein produced by the ovary as well as anterior pituitary folliculostellate cells. It is a high affinity activin binding protein therefore its main role in female reproduction is to sequester activin and inhibit FSH secretion.

Anti-Mullerian hormone

AMH is produced by granulosa cells. AMH production increases from primordial to preantral follicle stage, being highest in preantral and small antral follicles but declining thereafter. AMH is implicated in inhibition of follicular recruitment from the resting pool. It acts on antral follicles to decrease FSH sensitivity, reduce aromatase activity and allow the selection of the dominant follicle (Visser and Themmen, 2014). AMH is increasingly being used as a biomarker of the relative size of the ovarian reserve (Pankhurst, 2017). Recently, receptors for AMH have also been identified in the mouse and human hypothalamic neurons suggesting a potential direct role in the neuroendocrine system (Cimino et al., 2016).

All intra-ovarian factors act in a paracrine manner to influence follicular growth and maturation (Gougeon, 2010).

Regulation of the HPO Axis

The cyclic functioning of the HPO axis involves a complex interaction between the hypothalamus, pituitary and ovary that include feed-forward, feedback and paracrine loops (Fig. 4).




Fig. 4 Hypothalamic–pituitary–ovarian axis depicting feed-forward, feedback, and paracrine loops.

The Feed-Forward Regulation in HPO Axis

A pulsatile release of GnRH by the hypothalamic GnRH pulse generator leads to feed-forward regulation of the anterior pituitary gonadotropes that promotes synthesis, storage and secretion of gonadotropins (Fig. 4). Gonadotropin secretion depends on the frequency of pulsatile GnRH secretion with faster frequency favoring LH- β subunit gene transcription and LH release, and slower frequency favoring FSH- β gene transcription and FSH release.

The gonadotropins released under the influence of GnRH have feed-forward actions at the ovarian level (Fig. 4) where they bring about morphological and hormonal changes. These include regulation of follicular differentiation by FSH, induction of ovulation and formation of corpus luteum by the surge release of LH, stimulation of theca cell androgen production by LH, conversion of androgen to estrogen as well as synthesis of inhibin and activin by FSH, and the stimulation of progesterone production in the corpus luteum by LH.

The Feedback Regulation of the HPO Axis

The feedback regulation of HPO axis, which may be negative or positive, involves control of hormonal release from the hypothalamus, pituitary or ovary by the end products of their stimulation (Fig. 4). Such feedback ensures that follicular growth, maturation, ovulation, corpus luteum formation and preparation of the uterine lining occur in a controlled and timely manner so as to facilitate successful conception or renewal of the reproductive cycle in the event of unsuccessful conception.

Negative feedback regulation

Estradiol, progesterone, and inhibin are the main ovarian hormones that participate in the negative feedback regulation of the neuroendocrine axis to decrease gonadotropin release (Fig. 4). At the level of the hypothalamus, the exact afferent neuronal pathways and mechanisms mediating the gonadal steroid-dependent negative feedback process remain unresolved as the majority of GnRH neurons do not express receptors for gonadal steroids. Current evidence suggests that kisspeptin neurons located in the POA and ARC play a key role conveying the feedback regulatory effects of gonadal steroids on GnRH neurosecretory activity. Kisspeptin neurons located in the ARC (KNDy neurons) express receptors for gonadal steroids and have been postulated to mediate the negative feedback actions of estradiol on GnRH secretion (Lehman et al., 2010). There is considerable evidence that the inhibitory effects of progesterone on GnRH pulsatile secretion is mediated through dynorphin from KNDy neurons. This premise is supported by the observations that KNDy neurons express the progesterone receptor, local administration of an antagonist to the dynorphin receptor in the MBH increases LH pulse frequency during the luteal phase in sheep, ovariectomy decreases the expression of dynorphin in the ARC and local administration of a progesterone receptor antagonist to the ARC disrupts the negative feedback actions of progesterone on GnRH/LH secretion (Lehman et al., 2010).

The inhibitory effects of estradiol on gonadotropin secretion, apart from its action on GnRH release, also appear to be partially mediated via direct pituitary effects. This involves repression of FSH- β subunit expression by estradiol. Inhibin from the ovary also exerts negative feedback on the pituitary FSH secretion and one mechanism proposed involves competing with activin in binding to its receptors, thereby preventing activin's role in stimulating FSH secretion.

Positive feedback regulation

The ovarian hormone estradiol is the main player that exerts positive feedback action in the HPO axis (Fig. 4) and research findings suggest that there are important differences in the neuronal systems mediating its positive actions on GnRH surge release between different animal models. While in rodents it appears that the anteroventral periventricular area is the main hypothalamic area involved, in sheep and primates the estradiol positive feedback is largely mediated in the MBH. Despite these differences, in most female mammals increased expression of kisspeptin during the late follicular phase promotes a robust release of GnRH that in turn stimulates the surge release of gonadotropins. In humans, the anterior pituitary appears to be an important site of estradiol positive feedback regulation of LH secretion.

Paracrine Regulation

In addition to the positive and negative feedback regulation discussed earlier, there are several paracrine/autocrine regulatory loops at all levels of the HPO axis. For instance, GnRH has been shown to regulate its own secretion (Padmanabhan et al., 1995). Several neuropeptides produced by hypothalamic neurons, such as kisspeptin, also participate in paracrine regulation of GnRH release. More recent studies suggest neurosteroids produced within the hypothalamus may also participate in feedback regulation of GnRH release (Kenealy et al., 2013). At the pituitary level, activin, inhibin and follistatin produced by the pituitary modulate FSH synthesis through paracrine actions. Activins promote FSH secretion by increasing the number of GnRH receptors on gonadotropes (Fortin et al., 2015) while inhibin and follistatin oppose activin actions.

At the level of the ovary, there are several paracrine systems operational. Some examples of such paracrine regulation include: (1) modulation of activin action by its binding protein, follistatin, (2) promotion of theca androgen production by inhibin, (3) down-regulation of FSH receptors in antral follicles by AMH, which helps reduce FSH sensitivity thus allowing emergence of dominant follicles, (4) induction of inhibin and gonadotropin receptors by estradiol, and (5) promotion of FSH-induced estrogen production in granulosa cells and LH-induced androgen production in theca cells by inhibins. Another important example of a paracrine




Fig. 5 Hypothalamic, pituitary, and ovarian hormonal, ovarian follicular and corpus luteum and uterine epithelial changes during the menstrual cycle. The mean concentrations of each of the hormonal changes are shown with LH pulsatile secretion shown as insets during the early and later follicular, periovulatory and luteal phases. *E2*, estradiol; *P4*, progesterone; *inh.A*, inhibin A; *inh.B*, inhibin B. Adapted with permission from Mahutte, N. G. and Ouhilal, S. (2007). In: Hurd, W. W. and Falcone, T. (eds.) *Clinical reproductive medicine and surgery*. St. Louis, MO: Elsevier.

feedback loop within the ovary involves androgen and inhibin with inhibin acting on theca cells to regulate androgen production and androgen acting back on granulosa cells to regulate inhibin production.

Other Regulators of HPO Axis

In addition to regulation of the HPO axis by its components and their secretions, nutrition, metabolic status and seasonality, among other factors, also regulate the HPO axis. Reproductive processes in female mammals can be energetically costly, thus requiring a coordinated regulation by the nutritional and metabolic systems. Therefore, the HPO axis must receive and integrate information about peripheral metabolic cues in order to regulate reproductive function. Neuropeptides and leptin are the major cues of the body's metabolic status.

Although leptin, a metabolic hormone produced primarily by adipocytes, does not affect GnRH secretion directly as its receptor is not present on GnRH neurons, the effects of leptin are mediated through intermediate neuronal pathways that encompass hypothalamic neuropeptide Y (NPY)/agouti-related protein (AgRP) neurons and proopiomelanocortin (POMC) neurons. While NPY has inhibitory action, melanocyte-stimulating hormone alpha (α MSH), one of the products of the *POMC* gene, has stimulatory effect on GnRH neurons (Evans and Anderson, 2017). Undernutrition upregulates NPY expression in the ARC reducing the GnRH pulsatility while increased leptin concentrations in circulation not only downregulates NPY but also increases α MSH increasing the GnRH/LH pulsatility (Evans and Anderson, 2017).

Photoperiod is a major factor determining annual changes in reproductive function in mammalian species that breed seasonally such as sheep (short day breeders) and mares (long day breeders). While the neuronal mechanisms involved in photoperiod control of GnRH/LH pulses remain uncertain it may involve hypothalamic dopaminergic neurons, *KNDy* neurons in the ARC, and neurons expressing RF-amide related peptide 3, which is also referred to gonadotropin inhibitory hormone 3 (Clarke, 2011). Another important factor modulating GnRH release in female mammals is stress. Stress influences the GnRH pulse generator reducing the frequency of GnRH/LH pulses through increased cortisol secretion (Whirledge and Cidlowski, 2010).

Apart from actions at the GnRH neurons, metabolic hormones such as insulin, leptin and adiponectin enhance gonadotropin synthesis and release via direct action on the gonadotropes. Stress in addition to reducing GnRH pulsatility also reduces pituitary responsiveness to GnRH, and amplifies the negative feedback while inhibiting the positive feedback effects of estradiol (Whirledge and Cidlowski, 2010).

At the level of the ovary, the major metabolic hormones modulating ovarian function are insulin and adiponectin. Insulin is referred to as a co-gonadotropin in the ovary as in addition to acting alone can potentiate gonadotropin action. Insulin can also activate IGF receptors in the ovary. The ovary also expresses adiponectin receptors and adiponectin promotes ovulatory and luteinization processes induced by the LH surge by increasing steroidogenesis, prostaglandin synthesis and angiogenesis (Palin et al., 2012). While glucocorticoids have an anti-inflammatory role in the ovary and aid in the formation of the corpus luteum, stress-induced increase in glucocorticoids has negative effects by suppressing the LH-induced steroidogenesis (Whirledge and Cidlowski, 2010).

Conclusions

Precise coordination of components of the HPO axis via endocrine and paracrine feedback mechanisms is essential to maintain adequate reproductive cyclic function. At the beginning of reproductive cycle, increasing levels of LH and FSH facilitate growth of multiple follicles to early antral stage (Fig. 5). The estradiol and inhibin B produced by the growing follicles exert negative feedback effects at the hypothalamic–pituitary level leading to a reduction in FSH secretion. The increased estradiol also promotes uterine epithelial growth to prepare for receiving the conceptus. The follicle that develops increased sensitivity to FSH in the face of reduced FSH support differentiates into the dominant follicle. The dominant follicle contributes to a further increase in estradiol secretion leading to positive feedback response at the hypothalamic–pituitary level and surge release of gonadotropins triggering ovulation. Upon release of the oocyte, the follicle remnant develops into a progesterone and inhibin A producing organ called corpus luteum. The increase in progesterone and inhibin secreted by the corpus luteum exerts negative feedback at the hypothalamic–pituitary level to reduce LH and FSH secretion. In the event of conception, luteal function is maintained by factors produced by the developing conceptus. If conception fails, the corpus luteum undergoes regression leading to a decline in progesterone and inhibin A secretion that result in removal of negative feedback effects at the hypothalamic–pituitary level leading to an increase in LH and FSH secretion and reinitiation of the cycle. All in all, the HPO axis integrates peripheral signals and via feed-forward, feedback and paracrine mechanisms it enables reproductive success to ensure the survival of the species.

References

- Beshay, V. E., & Carr, B. R. (2013). Hypothalamic-pituitary-ovarian axis and control of the menstrual cycle. In *Clinical reproductive medicine and surgery*. New York: Springer.
- Cimino, I., Casoni, F., Liu, X., Messina, A., Parkash, J., Jamin, S. P., Cateau-Jonard, S., Collier, F., Baroncini, M., DeWailly, D., Pigny, P., Prescott, M., Campbell, R., Herbison, A. E., Prevot, V., & Giacobini, P. (2016). Novel role for anti-Müllerian hormone in the regulation of GnRH neuron excitability and hormone secretion. *Nature Communications*, 7, 10055.
- Clarke, I. J. (2011). Control of GnRH secretion: one step back. *Frontiers in Neuroendocrinology*, 32, 367–375.
- Evans, M. C., & Anderson, G. M. (2017). Neuroendocrine integration of nutritional signals on reproduction. *Journal of Molecular Endocrinology*, 58, R107–R128.
- Fortin, J., Ongaro, L., Li, Y., Tran, S., Lamba, P., Wang, Y., Zhou, X., & Bernard, D. J. (2015). Minireview: activin signaling in gonadotropes: what does the FOX say ... to the SMAD? *Molecular Endocrinology*, 29, 963–977.
- Gougeon, A. (2010). Human ovarian follicular development: from activation of resting follicles to preovulatory maturation. *Annales D'Endocrinologie*, 71, 132–143.
- Kenealy, B. P., Kapoor, A., Guerriero, K. A., Keen, K. L., Garcia, J. P., Kurian, J. R., Ziegler, T. E., & Terasawa, E. (2013). Neuroestradiol in the hypothalamus contributes to the regulation of gonadotropin releasing hormone release. *The Journal of Neuroscience*, 33, 19051–19059.
- Lehman, M. N., Coolen, L. M., & Goodman, R. L. (2010). Minireview: kisspeptin/neurokinin B/dynorphin (*KNDy*) cells of the arcuate nucleus: a central node in the control of gonadotropin-releasing hormone secretion. *Endocrinology*, 151, 3479–3489.
- Miller, W. L. (2008). Steroidogenic enzymes. *Endocrine Development*, 13, 1–18.
- Padmanabhan, V., Evans, N. P., Dahl, G. E., McFadden, K. L., Mauger, D. T., & Karsch, F. J. (1995). Evidence for short or ultrashort loop negative feedback of gonadotropin-releasing hormone secretion. *Neuroendocrinology*, 62, 248–258.

- Palin, M. F., Bordignon, V. V., & Murphy, B. D. (2012). Adiponectin and the control of female reproductive functions. *Vitamins and Hormones*, *90*, 239–287.
- Pankhurst, M. W. (2017). A putative role for anti-Müllerian hormone (AMH) in optimising ovarian reserve expenditure. *The Journal of Endocrinology*, *233*, R1–R13.
- Schneider, F., Tomek, W., & Gründker, C. (2006). Gonadotropin-releasing hormone (GnRH) and its natural analogues: a review. *Theriogenology*, *66*, 691–709.
- Visser, J. A., & Themmen, A. P. (2014). Role of anti-Müllerian hormone and bone morphogenetic proteins in the regulation of FSH sensitivity. *Molecular and Cellular Endocrinology*, *382*, 460–465.
- Whirledge, S., & Cidlowski, J. A. (2010). Glucocorticoids, stress, and fertility. *Minerva Endocrinologica*, *35*, 109–125.
- Wijayarathna, R., & De Kretser, D. M. (2016). Activins in reproductive biology and beyond. *Human Reproduction Update*, *22*, 342–357.

"Systems Biology of Reproduction"

Spring 2024 (Even Years) – Course Syllabus
 Biol 475/575 Undergraduate/Graduate (3 Credit)
 SLN: (475) – 06763, (575) – 06764
 Time - Tuesday and Thursday 10:35 am-11:50 am
 Course Lectures in person and recorded on Canvas/Panopto and Discussion Sessions live in person and on WSU Zoom for all campuses (Hybrid Course)
 Room – CUE 418
 Course Director – Michael Skinner, Abelson Hall 507, 335-1524, skinner@wsu.edu
 Co-Instructor – Eric Nilsson, Abelson Hall 507, 225-1835, nilsson@wsu.edu
Learning Objective -
 Current literature based course on the Systems Biology of Reproduction. Learning Systems approaches to the biology of reproduction from a molecular to physiological level of understanding.

Schedule/Lecture Outline –

January	9 & 11	Week 1	Systems Biology Introduction
	16 & 18	Week 2	Molecular/ Cellular/ Reproduction Systems
	23 & 25	Week 3	Sex Determination Systems
Jan /Feb	30 & 1	Week 4	Male Reproductive Tract Development & Function
February	6 & 8	Week 5	Female Reproductive Tract Development & Function
	13 & 15	Week 6	Gonadal Developmental Systems Biology
	20 & 22	Week 7	Testis Systems Biology
	27 & 29	Week 8	Ovary Systems Biology
March	5 & 7	Week 9	Epigenetics and Transgenerational Gonadal Disease
	11 – 15	Week 10	Spring Break
	19 & 21	Week 11	Gametogenesis/ Stem Cells/ Cloning
	26 & 28	Week 12	Hypothalamus- Pituitary Development & Function
April	2 & 4	Week 13	Reproductive Endocrinology Systems
	9 & 11	Week 14	Fertilization & Implantation Systems
	16 & 18	Week 15	Fetal Development & Birth Systems
	23 & 25	Week 16	Assisted Reproduction/Contraception
Apr/May	30 & 2	Week 17	Exam or Grant Review

Spring 2024 – Systems Biology of Reproduction
Lecture Outline – Hypothalamus-Pituitary Development & Function
 Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday
 March 26, 2024
 Week 12

Hypothalamus-Pituitary Development & Function

Cell Biology

Structure / Lobes and Development
 Cell Populations and Hormones
 Regulators and Mutations

Hormones

Growth Hormone / Receptors / GHRH
 Prolactin / Development
 Opiomelanocortin

Gonadotropins

GnRH / Pulsitive Secretion
 GnRH Actions / Signaling
 LH/FSH Pulsitive Secretion/Menstrual Cycle

Regulation of Development

Cyclisity / Estrous Cycle / Circadian Systems

Required Reading

de Kretser, et al. (2018) Hypothalamic Pituitary Testis Axis. In: Encyclopedia of Reproduction (Second Edition), Volume 1, Pages 180-183.

Padmanabhan, et al. (2018) Hypothalamus–Pituitary–Ovary Axis. In: Encyclopedia of Reproduction (Second Edition), Volume 2, Pages 121-129.

Spring 2024 – Systems Biology of Reproduction
Discussion Outline – Hypothalamus-Pituitary Development & Function
 Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday
 March 28, 2024
 Week 12

Hypothalamus-Pituitary Development & Function

Primary Papers:

1. Belchetz et al. (1978) Science 202:631
2. Ozaki et al. (2021) Develop Growth Differ 63:154-165
3. Bhattacharya et al. (2019) Mol Reprod Dev 86 :1505-30

Discussion

Student 6: Reference 1 above

- What unique endocrine parameter was identified in the hypothalamic regulation of pituitary function?
- What physiological advantage does this have?
- How does this information fit into the understanding of Brain-Pituitary-Gonadal axis?


Student 7: Reference 2 above



- What was the experimental design and objectives of the study?
- How did the in vivo development maintain the organoid?
- What insights into organogenesis of the hypothalamus-pituitary were obtained?

Student 8: Reference 3 above

- How did the hypothalamus and pituitary induce puberty?
- What was the hypothalamus-pituitary and testis hormone axis established?
- What molecular mechanism in the hypothalamus and pituitary were involved?

Hypogonadal – pituitary – testis axis





Hypothalamus and Pituitary Cell Biology




Figure 1 | **Anatomy of the pituitary gland.** Sagittal section through the midline shows the pituitary gland within the diaphragma sella (a fold that encases the pituitary), situated immediately posterior to the sphenoid sinus.




Figure 22-1. Diagrammatic outline of the hypothalamus and the various parts of the organ in the adult.




Figure 22-2. Ultrastructure of the anterior pituitary, showing the characteristics of 5 types of secretory cells. Note that in this illustration, 3 morphologically different subtypes of gonadotropes (A, B, and C) have been identified. SFLL, follicle-stimulating cell (Sertoli) cell, with processes, from Sledge HW, Pincus JJ. Anterior pituitary. In: *Handbook of Endocrinology*. Case GH, Kaplan HW (eds). CRC Press, 1985. Copyright © CRC Press, Inc., Boca Raton, FL.





Figure 2. Sagittal MRI scan showing anterior pituitary hypoplasia. APH—anterior pituitary hypoplasia. PP—posterior pituitary, which is ectopic on this image. OC—optic chiasm, normal here. The infundibulum connecting the pituitary to the hypothalamus is not shown.



Hypothalamic-derived cells
GT1-1, GT1-3, GT1-7
Postmigratory, HIGH GnRH

Olfactory placode-derived cells
NLT, Gn10, Gn11
Migratory, LOW GnRH

TRENDS in Endocrinology & Metabolism

Figure 1. GnRH neuronal migration in the mouse and targeting of immortalized GnRH neuronal cell lines at specific windows across development. Immortalized NLT and GnRH neuronal cells were derived from a tumor in the olfactory region to cribriform plate, whereas the GT GnRH neuronal cells were derived from a tumor in the forebrain. The photomicrograph is from E15 mouse brain stained for GnRH and peripherin, shows the normal migratory route of GnRH neurons from the olfactory placode region across the cribriform plate to the forebrain and is adapted from [1]. Abbreviations: BF, basal forebrain; CP, cribriform plate; E, embryonal day; GnRH, gonadotropin-releasing hormone; OB, olfactory bulb; OP/VNO, olfactory placode/vomerolateral organ.




Figure 4. Phorbol ester signaling in GT1 GnRH neurons promotes neurite outgrowth. 12-O-tetradecanoylphorbol-13 acetate (TPA) signals to PKC, and in GT GnRH neurons, PKC promotes neurite outgrowth, translocates to the growth cone and activates MAPK. It also localizes to focal adhesions to stimulate FAK and p130Cas phosphorylation. PKC is proposed to be another activator of neurite outgrowth. PKC, by contrast, is a negative regulator of neurite outgrowth and initially localizes to the nucleus, then to membrane ruffles. The PKC pathway also promotes neurite outgrowth through alterations in N-cadherin and translocation of -catenin to the growth cone. Based on results in [21, 22 and 23]. Abbreviations: -cat, -catenin; FAK, focal adhesion kinase; GnRH, gonadotropin-releasing hormone; MAPK, mitogen-activated protein kinase; PKC, protein kinase C; TPA, 12-O-tetradecanoylphorbol-13 acetate.









Table 22-1. Pituitary hormones in mammals.*

Name and Source	Principal Actions
Anterior lobe	
Thyroid-stimulating hormone (TSH, thyrotropin)	Stimulates thyroid secretion and growth of thyroid gland.
Adrenocorticotrophic hormone (ACTH, corticotropin)	Stimulates secretion and growth of zona fasciculata and zona reticularis of adrenal cortex.
Growth hormone (GH, somatotropin, STH)	Accelerates body growth; stimulates secretion of IGF-I.
Follicle-stimulating hormone (FSH)	Stimulates ovarian follicle growth in female and spermatogenesis in male.
Luteinizing hormone (LH, interstitial cell-stimulating hormone, ICSH)	Stimulates ovulation and luteinization of ovarian follicles in female and testosterone secretion in male.
Prolactin (lactotropic hormone, LTH, luteotropin, lactogenic hormone, mamotropin)	Stimulates secretion of milk and maternal behavior. Maintains corpus luteum in female rodents but apparently not in other species.
β -Lipotropin (β -LPH)	?
γ -Melanocyte-stimulating hormone (γ -MSH)	May maintain adrenal sensitivity.
Intermediate lobe	
α - and β -melanocyte-stimulating hormones (α - and β -MSH; referred to collectively as melanotropin or eumelanin)	Expands melanophores in fish, amphibians, and reptiles; stimulates melanin synthesis in melanocytes in humans.
γ -Lipotropin (γ -LPH), corticotropin-like intermediate lobe peptide (CLIP), other fragments of pro-opiomelanocortin	?
Posterior lobe	
Vasopressin (antidiuretic hormone, ADH)	Promotes water retention.
Oxytocin	Causes milk ejection.

*In addition, a variety of gastrointestinal and other polypeptides are found in one or more lobes of the pituitary gland. These include CCK, gastrin, renin, angiotensin II, and calcitonin gene-related peptide (CGRP).

Table 1. Anterior pituitary cell types and hormone regulation

	Somatotropes	Lactotropes	Thyrotropes	Corticotropes	Gonadotropes
Hormone product	Growth hormone (GH)	Prolactin (PRL)	Thyroid-stimulating hormone (TSH)	Adrenocorticotrophic hormone (ACTH)	Gonadotropins: luteinizing hormone (LH) and follicle-stimulating hormone (FSH)
Site of action	Liver, kidney, most tissues	Mammary	Thyroid	Adrenal	Ovary, testis
Positive regulator	Growth hormone-releasing hormone	Estrogen, thyrotropin-releasing hormone	Thyrotropin-releasing hormone	Corticotropin-releasing hormone	Gonadotropin-releasing hormone
Negative regulator	Somatostatin, insulin-like growth factor	Dopamine	Thyroid hormone	Corticosteroids	Gonad steroids, inhibits
Hypopituitarism phenotype	Dwarfism	Failure to lactate	Thyroid hypoplasia, dwarfism, cretinism, hypothyroidism	Adrenal hypoplasia	Sexual immaturity
Hyperpituitarism phenotype	Gigantism, acromegaly		Thyroid hyperplasia, hyperthyroidism	Cushing disease	Precocious puberty



Hypothalamus and Pituitary Development










Fig. 1. Development of the hypothalamic-pituitary axis. The midline embryological primordia of the pituitary in the anterior neural ridge and of the endocrine hypothalamus in the neural plate are indicated. Cell types later arise in a temporally and spatially specific fashion: E12.5 corticotrope (C), E12.5 rostral tip thyrotropes (Tr), and subsequently, from E15.5 to E17.5, from somatotropes (S), lactotropes (L), thyrotropes (T), gonadotropes (G), and melanotropes (M).

Related pituitary cell lineages develop into interdigitated 3D cell networks.

Budry L, et al. (2011) Proc Natl Acad Sci U S A. 26;108(30):12515-20.









FIG. 5. Analysis of *Pitx1/Hesx1: Pitx1/TLE1* double-transgenic embryos. (A) Loss of anterior pituitary and morphological similarity to pituitary defects in Ames *df/df* mice, except when the *Pitx1/Hesx1* transgene encodes a mutated eh1 domain, which cannot interact with TLE1. H&E staining. (B) *In situ* hybridization showing loss of growth hormone (*GH*), *Pit-1*, and ventral *Prop-1*, while expression of proopiomelanocortin (*POMC*) appears unaffected and *Lhx3* and *Prop-1* continue to be expressed in the ectoderm of RP. Asterisk indicates the dorsal pouch ectoderm.



Molecular Mechanisms Governing Embryonic Differentiation of Pituitary Somatotropes.
Ellsworth BS, Stallings CE.
Trends Endocrinol Metab. 2018 Jul;29(7):510-523.





FIGURE 3. Proposed model for pituitary cell lineage and regulation of proliferation by secreted factors. The five endocrine cell types (gray) in the anterior pituitary are labeled with the hormone product that they secrete. Steps in precursor-cell differentiation that might be regulated by transcription factors (gray boxes) are also indicated. Possible regulation of cell proliferation by secreted factors (gray boxes) is also indicated. The points where SV40 T-Antigen transgenes are thought to block differentiation are shown in gray^{53,54}. The block in differentiation results in pituitary tumors that have been immortalized for further study. Abbreviations: ACTH, adrenocorticotropic hormone; FSH, follicle-stimulating hormone; GH, growth hormone; LH, luteinizing hormone; PRL, prolactin; T4, thyroid hormone; TSH, thyroid-stimulating hormone.




Table 1

Transcription factors associated with human pituitary disorders.

Gene	Class	Locus	Inheritance	Phenotype	Animal model	Pituitary phenotypes
<i>HESX1</i>	Paired HD	3p21	Recessive	CPHD with SOD	Knockout	Absence of pituitary or multiple oral ectoderm invagination and cellular proliferation.
<i>PITX2</i>	Bicoid-like HD	4q25	Dominant	IGHD	Knockout	RP forms but fails to proliferate and differentiate at E12.5; lacks all cell types except corticotropes.
<i>LHX3</i>	LIM HD	9q34	Recessive	CPHD with rigid cervical spine	Knockout	Hypoplastic pituitary; RP forms but is unable to proceed; lacks all cell types except corticotropes.
<i>LHX4</i>	LIM HD	1q25	Dominant	CPHD with defects in sella turcica and cerebellar tonsil	Knockout	Hypoplastic anterior pituitary with reduction of all cell types; increased apoptosis.
<i>GLI2</i>	Kruppel	2q14	Dominant	CPHD with variable HPE features	Knockout you-too	Variable loss of pituitary; deletion of both <i>Gl1</i> and <i>Gl2</i> causes complete loss of pituitary.
<i>PROP1</i>	Paired HD	5q35	Recessive	CPHD	Knockout	Transdifferentiation into a lens.
<i>PIT1</i>	POU HD	3q11	Recessive	CPHD	Ames dwarf	Lack somatotropes, thyrotropes and lactotropes, reduced LH and FSH
<i>TBX19</i>	T-box	1q23	Recessive	ACTH deficiency	Shell dwarf mutations	Loss of somatotropes, thyrotropes and lactotropes and increased gonadotropes. Loss of somatotropes, thyrotropes and lactotropes and expanded corticotropes.
<i>SF1</i>	NR	9q33	Dominant	Adrenal failure, 46, XY gonadal dysgenesis	Knockout	Reduced corticotropes and melanotropes, melanotropes transdifferentiate into gonadotropes and <i>Pit1</i> -independent thyrotropes.
<i>SIX6</i>	SIX HD	14q23	Dominant	Bilateral anophthalmia and pituitary anomalies	Knockout	Impaired pituitary FSH and LH expression.
<i>SOX3</i>	HMG-box	Xq27	Recessive	IGHD with X-linked mental retardation	Knockout	Hypoplastic pituitary.

Abbreviations: HD, homeodomain; HMG, high mobility group; HPE holoprosencephaly, NR nuclear receptor.


Table 2





Signal pathways and other transcription factors critical for pituitary development and function.

Gene	Expression	Animal model	Pituitary phenotypes
Signaling molecules/receptors			
<i>BMP4</i>	VD	Knockout and <i>Pitx1-Noggin Tg</i>	RP fails to form, embryonic lethal.
<i>FGF10</i>	VD	knockout	RP arrested at E10 with loss of all cell types except corticotropes.
<i>FGFR2-IIIb</i>	RP	knockout	Anterior pituitary agenesis owing to increased apoptosis.
<i>FGF3, Zebrafish</i>	VD	Knockout	Anterior pituitary agenesis owing to increased apoptosis.
<i>Wnt4</i>	RP	Knockout	Increased apoptosis leading to a complete loss of pituitary.
<i>Wnt5</i>	VD	Knockout	Hypoplastic pituitary with marked reduction of <i>Pit1</i> lineages.
<i>SHH</i>	VD, oral ectoderm except RP	<i>Pitx1-HIP Tg</i>	Pituitary dysmorphogenesis, cell differentiation occurs normally.
<i>GHRH</i>	Hypothalamus	Knockout	Pituitary hypoplasia with loss of expression of ventral transcription factors
<i>GHRHR</i>	Anterior pit	<i>litle</i> , point mutation	<i>sonic-you (syc)</i> , zebrafish)
<i>GHRH2</i>	Hypothalamus	<i>lpg</i> , deletion	Hypoplastic pituitary with reduced <i>pomc</i> - and <i>pr</i> -positive cells and absence of <i>gh</i> and <i>tsh</i> -positive cells.
<i>D29</i>	Pituitary	Knockout	IGHD, hypoplastic pituitary with reduced production of GH.
<i>ACVR2</i>	Pituitary	Knockout	IGHD, postnatal dwarf, reduced proliferation of caudomedial somatotropes.
Transcription factors/cofactors			
<i>Tf1/Nkx2.1</i>	VD	Knockout	Decreased LHβ, FSHβ, ACTH, PRL in pituitary.
<i>Brr2</i>	Hypothalamus	Knockout	Prolacinomas.
<i>Gsh1</i>	Hypothalamus	Knockout	Reduced expression of <i>FSHβ</i> .
<i>Nhh2</i>	Hypothalamus, Pituitary	Knockout	Absence of pituitary, owing to ablation of FGFs expression domains in VD.
<i>Isl1</i>	RP	Knockout	Loss of posterior pituitary, owing to defects in survival of hypothalamus neurons.
<i>Pit1</i>	RP	Knockout	Hypoplastic anterior pituitary; reduced production of GHRH, GH, PRL, LH.
<i>Pax6</i>	Dorsal region of RP	Knockout	Hypogonadal; reduced production of FSH; adult-onset obesity. Deletion of both <i>Nhh1</i> and <i>Nhh2</i> results in significant reduction of GHRH-1 neurons.
<i>NeuroD1</i>	RP	Knockout	RP forms but remains primitive, thin pouch wall, embryonic lethal.
<i>Aes</i>	Dorsal region of RP	Knockout	Decreased expression of <i>LHβ</i> , <i>FSHβ</i> , <i>TSHβ</i> ; increased expression of <i>ACTH</i> .
<i>Tcf4</i>	ND	Knockout	Dorsal expansion of ventral cell types at the expense of dorsal cell types.
<i>Egr1</i>	Pit	Knockout	Delayed corticotropes differentiation.
<i>Otx1</i>	Postnatal pituitary	Knockout	Pituitary dysmorphogenesis.
			Hyperplastic pituitary, prolonged <i>Prop1</i> expression.
			No <i>LHβ</i> expression, reduced number of somatotropes.
			Transient dwarfism, delayed production of <i>LHβ</i> , <i>FSHβ</i> and GH.

Abbreviations: ND, not determined; RP, Rathke's pouch; Tg, transgenic; VD, ventral diencephalon.

Embryonic day: Signaling molecules: Transcription factors:










Fig. 1. Areas of expression (green or black) and action radius (indicated by red arrows) of bone morphogenetic protein 4 (BMP-4), BMP-2, fibroblast growth factor 8 (FGF-8) and sonic hedgehog (Shh) in Rathke's pouch (RP) and surrounding areas on embryonic day 8.5 (E8.5) (a) and E10.5 (b) in the mouse.










Fig. 2. Areas of expression (green) and action radius (indicated by red arrows) of bone morphogenetic protein 2 (BMP-2), fibroblast growth factor 8 (FGF-8) and chordin in embryonic pituitary (P) and surrounding areas on embryonic day 11.5 (E11.5) and E12.5 in the mouse. Areas in which corticotropes (C), gonadotropes (G), lactotropes (L), melanotropes (M), somatotropes (S), thyrotropes (T) and rostral thyrotropes (rT) first appear during embryonic development are indicated.

TGF- β Superfamily Regulation of Follicle-Stimulating Hormone Synthesis by Gonadotrope Cells: Is There a Role for Bone Morphogenetic Proteins?
 Ongaro L, Schang G, Ho CC, Zhou X, Bernard DJ.
 Endocrinology. 2019 Mar 1;160(3):675-683.





Dynamic Expression of Imprinted Genes in the Developing and Postnatal Pituitary Gland.
 Scagliotti V, Esse RCF, Willis TL, et al.
 Genes (Basel). 2021 Mar 30;12(4):509.



Hypothalamus and Pituitary Hormones




Table 1. Anterior pituitary cell types and hormone regulation.

Somatotropes	Lactotropes	Thyrotropes	Corticotropes	Gonadotropes
Hormone product Growth hormone (GH)	Prolactin (PRL)	Thyroid-stimulating hormone (TSH)	Adrenocorticotropic hormone (ACTH)	Gonadotropins: luteinizing hormone (LH) and follicle-stimulating hormone (FSH)
Site of action Liver, kidney, most tissues	Mammary	Thyroid	Adrenal	Ovary, testis
Positive regulator Growth-hormone-releasing hormone	Estrogen, thyrotropin-releasing hormone	Thyrotropin-releasing hormone	Corticotropin-releasing hormone	Gonadotropin-releasing hormone
Negative regulator Somatostatin, insulin-like growth factor	Dopamine	Thyroid hormone	Corticosteroids	Gonad steroids, inhibitors
Hypopituitarism phenotype Dwarfism	Failure to lactate	Thyroid hypoplasia, dwarfism, cretinism, hypothyroidism	Adrenal hypoplasia	Sexual immaturity
Hyperpituitarism phenotype Gigantism, acromegaly	Galactorrhoea, infertility	Thyroid hyperplasia, hyperthyroidism	Cushing disease	Precocious puberty





Figure 22-4. Structure of the principal human growth hormone (continuous chain). The black bars indicate disulfide bridges. The 29 residues alongside the chain identify residues that differ in human chorionic somatomammotropin (hCS; see Chapter 23). All the other residues in hCS are the same, and hCS also has 191 amino acid residues. (Reproduced, with permission, from Parfoniis, JA [editor], *Pepptide Hormones*. University Park Press, 1976.)



Nature Reviews | Cancer



Figure 22-6. Primary structure of human IGF-1 and 3. Identical variants: a 21-amino-acid residue extension at the C-terminal, a histopropyl substitution at Ser-30, and a histopropyl substitution at Ser-32. Single-letter codes are used for amino acid residues. (Reproduced, with permission, from Saw VH, Hall K. Human-like growth factors and their binding proteins. *Physiol Rev* 1990;70:281.)

THE PITUITARY GLAND 7 305




Figure 22-10. Current view of actions mediated by growth hormone (GH) and IGF-1. (Courtesy of R Daak and N Goussard.)








Figure 22-11. Feedback control of growth hormone secretion. The dashed arrows indicate inhibitory effects and the solid arrows stimulatory effects. Note that IGF-1 stimulates the secretion of somatostatin (SS) from the hypothalamus and acts directly on the pituitary to inhibit growth hormone (GH) secretion. (Compare with Figs 15-14, 20-21, 22-22, and 23-36.)

Table 22-3. Stimuli that affect growth hormone secretion (Hormones).

Stimuli that increase secretion	Stimuli that decrease secretion
<ul style="list-style-type: none"> Deficiency of energy substrate Hypoglycemia 2-Diacylglycerol Exercise Fasting Increase in circulating levels of certain amino acids Protein meal Influx of arginine and some other amino acids Starvation Stressful stimuli Pyrogen Lipine anoxemia Various psychological stresses Going to sleep Dopa and n-dihydroxy amphetamines that penetrate the brain Apomorphine and other dopamine receptor agonists Estrogens and androgens 	<ul style="list-style-type: none"> Estrogens and androgens REM sleep Glucose Cortisol FFA Methoxyprogesterone Growth hormone



Macrophage colony-stimulating factor induces prolactin expression in rat pituitary gland.

Hoshino S, Kurotani R, Miyano Y, et al.
 Zoolog Sci. 2014 Jun;31(6):390-7.

Abstract

We investigated the role of macrophage colony-stimulating factor (M-CSF) in the pituitary gland to understand the effect of M-CSF on pituitary hormones and the relationship between the endocrine and immune systems. When we attempted to establish pituitary cell lines from a thyrotropic pituitary tumor (TtT), a macrophage cell line, TtT/M-87, was established. We evaluated M-CSF-like activity in conditioned media (CM) from seven pituitary cell lines using TtT/M-87 cells. TtT/M-87 proliferation significantly increased in the presence of CM from TtT/GF cells, a pituitary folliculostellate (FS) cell line. M-CSF mRNA was detected in TtT/GF and MtT/E cells by reverse transcriptase-polymerase chain reaction (RT-PCR), and its expression in TtT/GF cells was increased in a lipopolysaccharide (LPS) dose-dependent manner. M-CSF mRNA expression was also increased in rat anterior pituitary glands by LPS. M-CSF receptor (M-CSFR) mRNA was only detected in TtT/M-87 cells and increased in the LPS-stimulated rat pituitary glands. In rat pituitary glands, M-CSF and M-CSFR were found to be localized in FS cells and prolactin (PRL)-secreting cells, respectively, by immunohistochemistry. The PRL concentration in rat sera was significantly increased at 24 h after M-CSF administration, and mRNA levels significantly increased in primary culture cells of rat anterior pituitary glands. In addition, TNF- α mRNA was increased in the primary culture cells by M-CSF. **These results revealed that M-CSF was secreted from FS cells and M-CSF regulated PRL expression in rat pituitary glands.**




Fig. 7. Neurochemical mechanisms and their site of action, including the HPA axis. ACTH: adrenocorticotropic hormone; CRF: corticotropin-releasing factor; LL: locus caeruleus; NE: norepinephrine.






Figure 22-3. Schematic representation of the prepro-opiomelanocortin molecule formed in pituitary cells, neurons, and other tissues. The numbers in parentheses identify the amino acid sequences in each of the polypeptide fragments. For convenience, the amino acid sequences are numbered from the N terminal of ACTH and read toward the C-terminal portion of the parent molecule, whereas the amino acid sequences in the other portion of the molecule read to the left to -131, the N terminal of the parent molecule. The locations of Lys-Arg and other pairs of basic amino acids residues are also indicated; these are the sites of proteolytic cleavage in the formation of the smaller fragments of the parent molecule. AL, anterior lobe; IL, intermediate lobe.



Interindividual Variation in DNA Methylation at a Putative POMC Metastable Epiallele Is Associated with Obesity.
 Cell Metab. 2016 Sep 13;24(3):502-509.
 Kühnen P, Hanke D, Waterland RA, et al.





In Brief


Kühnen et al. explore the epigenetic regulation of body weight/BMI and identify a POMC DNA hypermethylation variant in obese individuals. This putative metastable epiallele is present after birth and leads to an increased individual risk of developing obesity later in life.

Highlights

- A POMC hypermethylation variant leads to increased risk for obesity development
- Maternal C1 metabolism at conception affects POMC methylation in the offspring
- Paternal POMC methylation correlates with methylation in the offspring
- This POMC region fulfills criteria for a putative metastable epiallele




Insulin-Like Factor 3 and the HPG Axis in the Male.
 Ivell R, Heng K, Anand-Ivell R. Front Endocrinol (Lausanne). 2014 Jan 27;5:6. eCollection 2014.





INSL3 and HPG axis. Scheme to show the relationship between the INSL3/RXFP2 system and testosterone as endpoint effectors of the HPG axis within the testis. Arrows are directed only to cells where there are known to be specific cognate receptors.

Complex integration of intrinsic and peripheral signaling is required for pituitary gland development.
 Edwards W, Raetzman LT,
 Biol Reprod. 2018 Sep 1;99(3):504-513.



Pubertal orchestration of hormones and testis in primates.
 Bhattacharya I, Sen Sharma S, Majumdar SS,
 Mol Reprod Dev. 2019 Nov;86(11):1505-1530.



VERTEBRATE HYPOTHALAMIC-HYPOPHYSIC-GONADAL AXIS

TABLE I
GnRH Molecular Forms

GnRH	1	2	3	4	5	6	7	8	9	10
Mammal	pGlu	His	Trp	Set	Tyr	Gly	Leu	Arg	Pro	Gly-NH ₂
Guinea pig	—	Tyr	—	—	—	—	Val	—	—	—
Chicken II	—	—	—	His	—	—	Trp	—	—	—
Chicken I	—	—	—	—	—	—	—	—	—	—
Frog	—	—	—	—	—	—	—	—	—	—
Salmon	—	—	—	—	—	—	—	—	—	—
Dogfish	—	—	—	—	His	—	—	—	—	—
Catfish	—	—	—	—	His	—	—	—	—	—
Scabreim	—	—	—	—	—	—	—	—	—	—
Medaka	—	—	—	—	—	—	—	—	—	—
Herring	—	—	—	—	—	—	—	—	—	—
Lamprey I	—	—	Tyr	—	—	Leu	Glu	Trp	Lys	—
Lamprey II	—	—	—	—	—	—	—	—	—	—
Tunicate I	—	—	—	—	—	—	—	—	—	—
Tunicate II	—	—	—	—	—	—	—	—	—	—




Fig. 3. Schematic of transcripts regulated by GnRH in LT2 cells. Reprinted from [196], with permission of The American Society for Biochemistry and Molecular Biology.




FIG. 5. Episodic secretion of biologically active LH as revealed by deconvolution analysis of pulsatile plasma bioactive LH profiles in normal women studied at three stages of the menstrual cycle. The stage of the menstrual cycle is noted in each subpanel. Plasma bioactive LH concentrations were measured in the rat interstitial cell testosterone bioassay in blood sampled at 10- or 15-minute intervals for 6 to 8 hours. (Left panels) Serial plasma bioactive LH concentrations and the computer-estimated recombinant fit (continuous curve) and individual half-lives (\pm SD) of LH estimated in each data series. (Right panels) Calculated bioactive LH secretory rates, the units of which are mIU/mL/min (LH/Lmin). Note that punctuated bursts of LH secretion without tonic release account for the pulsatile LH profiles. (Adapted from ref. 54, with permission.)




FIG. 4. Illustrative profiles of deconvolution-resolved LH secretory bursts in two women during each of three studied phases of the menstrual cycle. Upper panels: Serial LH concentrations measured by radioimmunoassay (mIU/ml). Intra-sample mean (\pm SD) and the fitted recombinant curve based on the calculated secretion and clearance parameters. (Lower panels) Resolved, discrete LH secretory bursts. (Shown with permission.)








FIG. 4. Frequency and amplitude of LH pulses on selected days of the normal human menstrual cycle. The days of the cycle are indicated relative to the day of the LH surge (day 0). Significant LH pulses are indicated by a dot. Luteal phase progesterone levels are 1, 5, 6, and 12 ng/ml on days +1, +2, +5, and +11, respectively. During the follicular phase (days -11 to -2), low-amplitude LH pulses occur almost hourly. On the day of the LH peak, the frequency of the LH pulses remains circadian, but the amplitude is increased. During the luteal phase, the LH pulse frequency is reduced. (From ref. 104.)






FIG. 12. The establishment of two ovulatory menstrual cycles in response to an unvarying pulsatile GnRH replacement regimen in a rhesus monkey subjected to a radiofrequency lesion in the MBH at time 0. Note that the lesion abolished the positive feedback action of administered estradiol/benzate (EB). Diastereotomy at the end of the experiment, to open a putative negative feedback loop, did not increase gonadotropin secretion, indicating the completeness of the lesion and the absence of reorganization or regeneration of the hypothalamic apparatus responsible for the control of gonadotropin secretion. Small vertical lines below some baseline points indicate concentrations below the limits of sensitivity of the assays. (from ref. 52)






Fig. 4. Spectrum of GnRH abnormalities in 4 women with hypothalamic amenorrhea. ∇ = Detected LH pulsations. Note the predominance of LH pulsations during sleeping hours (open bars) [reproduced from 22].




Regulation Hypothalamus and Pituitary Development


Emerging insights into hypothalamic-pituitary-gonadal axis regulation and interaction with stress signalling.
 Acevedo-Rodriguez A, Kauffman AS, Cherrington BD, Borges CS, Roepke TA, Laconi M.
 J Neuroendocrinol. 2018 Oct;30(10):e12590.



The Processes of Anterior Pituitary Hormone Pulse Generation.
 Le Tissier P, Fiordeliso Coll T, Mollard P.
 Endocrinology. 2018 Oct 1;159(10):3524-3535.



The human hypothalamus in mood disorders: The HPA axis in the center.
 Bao AM, Swaab DF.
 IBRO Rep. 2018 Dec 14;6:45-53.





Gene coexpression network analysis identifies genes and biological processes shared among anterior pituitary and brain areas that affect estrous behavior in dairy cows.
 Kommadath A, Te Pas MF, Smits MA. (2013) J Dairy Sci. 2013 Apr;96(4):2583-95.

Table 1. Summary of steps taken in the analysis and their results

Steps	Results
Process data of 14 pairs of dye-swapped microarrays per tissue collected from 14 cows at start of estrous cycle	Gene expression values (M-values) of 23,496 probes per array obtained
Select good quality probes based on probe reannotation and average the M-values of probes representing the same gene	16,620 good-quality probes per array representing 13,234 genes obtained
Select the top 50% most variable genes per tissue and identify genes shared by each tissue within a tissue pair (assuming that genes affecting estrous behavior expression would have a variable expression across the experimental cows showing differing levels of estrous behavior)	Approximately 4,000 to 5,000 genes per tissue pair obtained: AM-DH, HC-DH, VH-DH, AM-HC, and AP-DH ¹
Perform coexpression network analysis on gene expression data of shared genes within each tissue pair and identify consensus modules	Gene coexpression networks constructed for tissues within each pair and consensus modules identified
Identify consensus modules within tissue pairs whose module eigenvalues correlate with estrous behavior scores	Consensus modules that correlated with estrous behavior identified: AM-DH: 1 of 3, HC-DH: 5 of 10, VH-DH: 0 of 2, AM-HC: 3 of 8, and AP-DH: 10 of 23
Test for enriched gene ontology and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway terms within estrous behavior-correlated consensus modules	Significant enriched terms obtained (summary of the significant terms per module are reported in Figure 1)
Identify hub genes within the estrous behavior correlated consensus modules	Hub genes per module obtained (the top 3 hub genes per module are reported in Table 3)

¹Tissues are abbreviated as follows: anterior pituitary (AP), dorsal hypothalamus (DH), ventral hypothalamus (VH), amygdala (AM), and hippocampus (HC).








GENOMIC REGULATION OF BOVINE ESTROUS BEHAVIOR





Table 4. Estrous behavior-associated genes and processes in dairy cows known to correspond to processes within the growth, amplification, preparation, permission, and synchronization (GAPPS) modules

GAPPS module	Characteristics	Corresponding genes and processes in cows (with reference in parentheses)
Growth	Increase in the input/output connections for behavior-directing hypothalamic neurons	Synaptic plasticity: Immune related genes: <i>CTLA4, IL1RL1, MARCO</i> (Kommadath et al., 2011) Neurotransmitter receptors: <i>CHRM1, CHRM3, CHRNA5</i> (Kommadath et al., 2011) Ribosomal genes: <i>RPL14, RPL18, RPL24, RPS11, RPS18</i> (this study) Others: <i>NEFL, NDRG2, THY1, GAP43</i> (this study)
Amplification	Amplification of estrogen effect by progesterone mediated by progesterone receptor	<i>PCR</i> gene upregulated in the anterior pituitary at d 0 (Kommadath, 2012)
Preparation	Preparation for mating	Female sexual receptivity: <i>OXT, AVP, HTR2A, DRD2, CABRA6</i> (this study and Kommadath et al., 2011) Anxiolytic effect: <i>OXT, FTH, KCNN2</i> (this study and Kommadath et al., 2011) Altered feeding motivation and mood: <i>POMC, MCHR1, MOBP, LTA4H</i> (Kommadath et al., 2011)
Permission	Permissive actions by hypothalamic neurons for the mating behavior to occur	Arousal, activation of protein kinases and release of Ca^{2+} : <i>CHRM1, CHRM3, CHRNA5, PLCB2, ITPKA</i> (Kommadath et al., 2011)
Synchronization	Synchronize mating behavior with ovulation	Prostaglandin regulators: <i>PTGDS, PTGIS, PTGFB</i> (Kommadath et al., 2011)


¹This table is adapted from the PhD thesis of the first author (Kommadath, 2012).








Somatotropic-Testicular Axis: A crosstalk between GH/IGF-I and gonadal hormones during development, transition, and adult age.
 Tenuta M, Carlomagno F, Cangiano B, et al.
Andrology. 2021 Jan;9(1):168-184.




Association between Vitamin D Levels, Puberty Timing, and Age at Menarche.
 Calcaterra V, Magenes VC, Tagi VM, Grazi R, Bianchi A, Cena H, Zuccotti G, Fabiano V.
Children (Basel). 2023 Jul 19;10(7):1243.




Potential mechanisms linking vitamin D and pubertal timing. VDR = vitamin D receptor.


Growth differentiation factor 9 (GDF9) forms an incoherent feed-forward loop modulating follicle-stimulating hormone β -subunit (FSH β) gene expression.
 Choi SG, Wang Q, Jia J, Pincas H, Turgeon JL, Sealfon SC.
J Biol Chem. 2014 Jun 6;289(23):16164-75.




Current concepts in neuroendocrine disruption.

León-Olea M, et al. (2014) *Gen Comp Endocrinol*. (Epub ahead of print).






Exposure to a low dose of bisphenol A impairs pituitary-ovarian axis in prepubertal rats: effects on early folliculogenesis.
Gámez JM, Penalba R, Cardoso N, et al.
Environ Toxicol Pharmacol. 2015 Jan;39(1):9-15.




Clocks underneath: the role of peripheral clocks in the timing of female reproductive physiology.
Sellix MT. Front Endocrinol (Lausanne). 2013 Jul 23;4:91.





Circadian clock function in the peripheral tissues of the female HPG axis. The central circadian clock in the suprachiasmatic nucleus (SCN) drives rhythmic GnRH secretion and subsequent gonadotropin secretion from the pituitary. In addition to these neuroendocrine pacemakers, clocks are also present in the pituitary gonadotroph, uterine endometrium and myometrium, oviduct epithelial cells and ovarian theca, interstitial, and granulosa cells. Clock function has been implicated in GnRH signaling, gonadotropin sensitivity, ovulation, steroid hormone synthesis, embryonic maturation, implantation, and decidualization. Synchronization of central and peripheral oscillators is mediated by several putative humoral and neural cues, driven either directly or indirectly by the SCN. Moreover, feedback signals from the periphery, e.g., steroid hormones of ovarian origin, modulate the timing of the clock in both central and peripheral tissues of the HPG axis.


Sex differences in circadian timing systems: implications for disease.
Bailey M, Silver R. Front Neuroendocrinol. 2014 Jan;35(1):111-39.



Unveiling "Musica Universalis" of the Cell: A Brief History of Biological 12-Hour Rhythms.
 Zhu B, Dacso CC, O'Malley BW.
 J Endocr Soc. 2018 Jun 6;2(7):727-752.



Role of core circadian clock genes in hormone release and target tissue sensitivity in the reproductive axis.
 Sen A, Hoffmann HM.
 Mol Cell Endocrinol. 2020 Feb 5;501:110655.



An integrative view of mammalian seasonal neuroendocrinology.
 Dardente H, Wood S, Ebling F, Sáenz de Miera C.
 J Neuroendocrinol. 2019 May;31(5):e12729.

