

Spring 2026 – Systems Biology of Reproduction
Lecture Outline – Gonadal Developmental Systems Biology
Eric Nilsson – Biol 475/575
10:35-11:50 am, Tuesday & Thursday
February 17, 2026
Week 6

Gonadal Developmental Systems Biology

Early Fetal Gonadal Development

- Morphogenesis
- Transcriptome
- Meiotic Arrest

Testis Gonadal Fetal Development

- Knockout Models and Genes
- Cellular Growth Regulation

Ovary Gonadal Fetal Development

- Developmental Timing and Morphology
- Oocyte Nests and Primordial Follicle Assembly
- Hormone Regulation of Assembly
- Culture Models

Description of Gonadal Development

- Endocrine Disruptors

Required Reading

Pepling and Burton (2018) Fetal/Gonadogenesis, Encyclopedia of Reproduction (Second Edition).
Volume 2, Pages 47-51.

Rotgers and Yao, (2018) Formation of the Testis Primordium, Encyclopedia of Reproduction (Second Edition). Volume 1, Pages 84-87.

REFERENCES

- Rajachandran S, Zhang X, Cao Q, Caldeira-Brant AL, et al. Dissecting the spermatogonial stem cell niche using spatial transcriptomics. *Cell Rep.* 2023 Jul 25;42(7):112737.
- Kumar V, Elkouby YM. Tools to analyze the organization and formation of the germline cyst in zebrafish oogenesis. *Development.* 2023 Jul 1;150(13):dev201349.
- Smaga CR, Bock SL, JM, Parrott BB. Sex Determination and Ovarian Development in Reptiles and Amphibians: From Genetic Pathways to Environmental Influences. *Sex Dev.* 2022 Nov 15:1-21.
- Calonga-Solís V, Fabbri-Scallet H, Ott F, et al. MYRF: A New Regulator of Cardiac and Early Gonadal Development-Insights from Single Cell RNA Sequencing Analysis. *J Clin Med.* 2022 Aug 18;11(16):4858.

- Real FM, Lao-Pérez M, Burgos M, et al. Cell adhesion and immune response, two main functions altered in the transcriptome of seasonally regressed testes of two mammalian species. *J Exp Zool B Mol Dev Evol.* 2023 May;340(3):231-244.
- Liu T, Li R, Liu L, Wu S, Zhang L, et al. The Effect of Temperature on Gonadal Sex Differentiation of Yesso Scallop *Patinopecten yessoensis*. *Front Cell Dev Biol.* 2022 Jan 31;9:803046.
- Tsuji-Hosokawa A, Ogawa Y, Tsuchiya I, Terao M, Takada S. Human SRY Expression at the Sex-determining Period is Insufficient to Drive Testis Development in Mice. *Endocrinology.* 2022 Jan 1;163(1):bqab217.
- Burgos M, Hurtado A, Jiménez R, Barrionuevo FJ. Non-Coding RNAs: lncRNAs, miRNAs, and piRNAs in Sexual Development. *Sex Dev.* 2021 Oct 6;1-16.
- Chumduri C, Turco MY. Organoids of the female reproductive tract. *J Mol Med (Berl).* 2021 Apr;99(4):531-553.
- Jolivet G, Daniel-Carlier N, Harscoët E, et al. Fetal Estrogens are not Involved in Sex Determination But Critical for Early Ovarian Differentiation in Rabbits. *Endocrinology.* 2022 Jan 1;163(1):bqab210.
- Rodprasert W, Toppari J, Virtanen HE. Endocrine Disrupting Chemicals and Reproductive Health in Boys and Men. *Front Endocrinol (Lausanne).* 2021 Oct 7;12:706532.
- Boateng R, Boechat N, Henrich PP, Bolcun-Filas E. Whole Ovary Immunofluorescence, Clearing, and Multiphoton Microscopy for Quantitative 3D Analysis of the Developing Ovarian Reserve in Mouse. *J Vis Exp.* 2021 Sep 3;(175).
- Connan-Perrot S, Léger T, Lelandais P. et al. Six Decades of Research on Human Fetal Gonadal Steroids. *Int J Mol Sci.* 2021 Jun 22;22(13):6681.
- Soto DA, Ross PJ. Similarities between bovine and human germline development revealed by single-cell RNA sequencing. *Reproduction.* 2021 Mar;161(3):239-253.
- Żyżyńska-Galeńska K, Bernat A, Piliszek A, et al. Embryonic Environmental Niche Reprograms Somatic Cells to Express Pluripotency Markers and Participate in Adult Chimaeras. *Cells.* 2021 Feb 25;10(3):490.
- Shen Y-C, Shami AN, Moritz L, et al. TCF21 + mesenchymal cells contribute to testis somatic cell development, homeostasis, and regeneration in mice. *Nat Commun.* 2021 Jun 23;12(1):3876.
- Guo J, Sosa E, Chitiashvili T, et al. Single-cell analysis of the developing human testis reveals somatic niche cell specification and fetal germline stem cell establishment. *Cell Stem Cell.* 2021 Apr 1;28(4):764-778.e4.
- Gao T, Lin M, Wu Y, et al. Transferrin receptor (TFRC) is essential for meiotic progression during mouse spermatogenesis. *Zygote.* 2021 Apr;29(2):169-175.
- Yan R-G, Li B-Y, Yang Q-E. Function and transcriptomic dynamics of Sertoli cells during prospermatogonia development in mouse testis. *Reprod Biol.* 2020 Dec;20(4):525-535.
- Feng C-W, Burnet G, Spiller CM, et al. Identification of regulatory elements required for Stra8 expression in fetal ovarian germ cells of the mouse. *Development.* 2021 Mar 9;148(5):dev194977.
- Liu M, Chen H, Dai H, et al. Effects of bis(2-butoxyethyl) phthalate exposure in utero on the development of fetal Leydig cells in rats. *Toxicol Lett.* 2021 Oct 15;351:65-77.
- Cargnelutti F, Di Nisio A, Pallotti F, et al. Effects of endocrine disruptors on fetal testis development, male puberty, and transition age. *Endocrine.* 2021 May;72(2):358-374.
- Wang J-J, Tian Y, Li M-H, et al. Single-cell transcriptome dissection of the toxic impact of Di (2-ethylhexyl) phthalate on primordial follicle assembly. *Theranostics.* 2021 Mar 5;11(10):4992-5009.

- Tanimoto R, Sekii K, Morohaku K, et al. Blocking estrogen-induced AMH expression is crucial for normal follicle formation. *Development*. 2021 Mar 19;148(6):dev197459.
- Peñalver Bernabé B, Thiele I, Galdones E, Siletz A, et al. Dynamic genome-scale cell-specific metabolic models reveal novel inter-cellular and intra-cellular metabolic communications during ovarian follicle development. *BMC Bioinformatics*. 2019 Jun 10;20(1):307.
- Shima Y, Miyabayashi K, Sato T, Suyama M, Ohkawa Y, Doi M, Okamura H, Suzuki K. Fetal Leydig cells dedifferentiate and serve as adult Leydig stem cells. *Development*. 2018 Dec 5;145(23).
- Tsuji-Hosokawa A, Kashimada K, Kato T, Ogawa Y, et al. Peptidyl arginine deiminase 2 (Padi2) is expressed in Sertoli cells in a specific manner and regulated by SOX9 during testicular development. *Sci Rep*. 2018 Sep 5;8(1):13263.
- Romer KA, de Rooij DG, Kojima ML, Page DC. Isolating mitotic and meiotic germ cells from male mice by developmental synchronization, staging, and sorting. *Dev Biol*. 2018 Nov 1;443(1):19-34.
- Rotgers E, Jørgensen A, Yao HH. At the Crossroads of Fate-Somatic Cell Lineage Specification in the Fetal Gonad. *Endocr Rev*. 2018 Oct 1;39(5):739-759.
- Correia B, Sousa MI, Ramalho-Santos J. The mTOR Pathway in Reproduction: From Gonadal Function to Developmental Coordination. *Reproduction*. 2019 Nov 1. pii: REP-19-0057.R2. doi: 10.1530/REP-19-0057. [Epub ahead of print]
- Pandey V, Tripathi A, Dubey PK. Expression and intracellular localization of Nanos2-homologue protein in primordial germ cells and spermatogonial stem cells. *Zygote*. 2019 Apr;27(2):82-88.
- Martinovic V, Vukusic Pusic T, Restovic I, et al. Expression of Epithelial and Mesenchymal Differentiation Markers in the Early Human Gonadal Development. *Anat Rec (Hoboken)*. 2017 Jul;300(7):1315-1326.
- Ben Maamar M, Lesné L, Hennig K, Desdoits-Lethimonier C, et al. Ibuprofen results in alterations of human fetal testis development. *Sci Rep*. 2017 Mar 10;7:44184.
- Bothun AM, Woods DC. Dynamics of WNT signaling components in the human ovary from development to adulthood. *Histochem Cell Biol*. 2019 Feb;151(2):115-123.
- Endo T, Mikedis MM, Nicholls PK, Page DC, de Rooij DG. Retinoic Acid and Germ Cell Development in the Ovary and Testis. *Biomolecules*. 2019 Nov 24;9(12).
- Pan B, Li J. The art of oocyte meiotic arrest regulation. *Reprod Biol Endocrinol*. 2019 Jan 5;17(1):8.
- Huelgas-Morales G, Greenstein D. Control of oocyte meiotic maturation in *C. elegans*. *Semin Cell Dev Biol*. 2018 Dec;84:90-99.
- Yadu N, Kumar PG. Retinoic acid signaling in regulation of meiosis during embryonic development in mice. *Genesis*. 2019 Jul;57(7-8):e23327.
- Zirkin BR, Papadopoulos V. Leydig cells: formation, function, and regulation. *Biol Reprod*. 2018 Jul 1;99(1):101-111.
- Bowles J, Feng CW, Ineson J, Miles K, Spiller CM, Harley VR, Sinclair AH, Koopman P. Retinoic Acid Antagonizes Testis Development in Mice. *Cell Rep*. 2018 Jul 31;24(5):1330-1341
- Fu A, Oberholtzer SM, Bagheri-Fam S, Rastetter RH, et al. Dynamic expression patterns of *Irx3* and *Irx5* during germline nest breakdown and primordial follicle formation promote follicle survival in mouse ovaries. *PLoS Genet*. 2018 Aug 2;14(8):e1007488.
- Rosario R, Crichton JH, Stewart HL, Childs AJ, Adams IR, Anderson RA. *Dazl* determines primordial follicle formation through the translational regulation of *Tex14*. *FASEB J*. 2019 Dec;33(12):14221-14233.

- Kato Y, Iwamori T, Ninomiya Y, Kohda T, Miyashita J, Sato M, Saga Y. ELAVL2-directed RNA regulatory network drives the formation of quiescent primordial follicles. *EMBO Rep.* 2019 Dec 5;20(12):e48251.
- Watanabe R, Sasaki S, Kimura N. Activation of autophagy in early neonatal mice increases primordial follicle number and improves lifelong fertility. *Biol Reprod.* 2019 Sep 30. pii: ioz179. doi: 10.1093/biolre/ioz179. [Epub ahead of print]
- Bhartiya D, Patel H, Sharma D. Heterogeneity of Stem Cells in the Ovary. *Adv Exp Med Biol.* 2019;1169:213-223.
- Li T, Liu X, Gong X, E Q, Zhang X, Zhang X. microRNA 92b-3p regulates primordial follicle assembly by targeting TSC1 in neonatal mouse ovaries. *Cell Cycle.* 2019 Apr;18(8):824-833.
- Guo C, Liu G, Zhao D, Mi Y, Zhang C, Li J. Interaction of Follicle-Stimulating Hormone and Stem Cell Factor to Promote Primordial Follicle Assembly in the Chicken. *Front Endocrinol (Lausanne).* 2019 Feb 19;10:91.
- Chang HM, Wu HC, Sun ZG, Lian F, Leung PCK. Neurotrophins and glial cell line-derived neurotrophic factor in the ovary: physiological and pathophysiological implications. *Hum Reprod Update.* 2019 Mar 1;25(2):224-242.
- Zhihan T, Xinyi M, Qingying L, Rufeif G, et al. Autophagy participates in cyst breakdown and primordial folliculogenesis by reducing reactive oxygen species levels in perinatal mouse ovaries. *J Cell Physiol.* 2019 May;234(5):6125-6135.
- Sifakis S, Androutsopoulos VP, Tsatsakis AM, Spandidos DA. Human exposure to endocrine disrupting chemicals: effects on the male and female reproductive systems. *Environ Toxicol Pharmacol.* 2017 Apr;51:56-70.
- Singh S, Singh SK. Effect of gestational exposure to perfluorononanoic acid on neonatal mice testes. *J Appl Toxicol.* 2019 Dec;39(12):1663-1671.
- Zhang Y, Mu X, Gao R, Geng Y, et al. Foetal-neonatal exposure of Di (2-ethylhexyl) phthalate disrupts ovarian development in mice by inducing autophagy. *J Hazard Mater.* 2018 Sep 15;358:101-112.
- Liu J, Wang Y, Fang Y, Ni C, Ma L, et al. Gestational exposure to ziram disrupts rat fetal Leydig cell development. *Chemosphere.* 2018 Jul;203:393-401.
- Pekar O, Ow MC, Hui KY, Noyes MB, Hall SE, Hubbard EJA. (2017) Linking the environment, DAF-7/TGF β signaling and LAG-2/DSL ligand expression in the germline stem cell niche. *Development.* 15;144(16):2896-2906.
- Panchal T, Chen X, Alchits E, Oh Y, Poon J, Kouptsova J, Laski FA, Godt D. (2017) Specification and spatial arrangement of cells in the germline stem cell niche of the *Drosophila* ovary depend on the Maf transcription factor Traffic jam. *PLoS Genet.* 19;13(5):e1006790.
- Ziková A, Lorenz C, Hoffmann F, Kleiner W, Lutz I, Stöck M, Kloas W. (2017) Endocrine disruption by environmental gestagens in amphibians - A short review supported by new in vitro data using gonads of *Xenopus laevis*. *Chemosphere.* 181:74-82.
- Pritykin Y, Brito T, Schupbach T, Singh M, Pane A. (2017) Integrative analysis unveils new functions for the *Drosophila* Cutoff protein in noncoding RNA biogenesis and gene regulation. *RNA.* 23(7):1097-1109.
- Ahn J, Park YJ, Chen P, Lee TJ, Jeon YJ, Croce CM, Suh Y, Hwang S, Kwon WS, Pang MG, Kim CH, Lee SS, Lee K. (2017) Comparative expression profiling of testis-enriched genes regulated during the development of spermatogonial cells. *PLoS One.* 12(4):e0175787.

- Jadhao AG, Pinelli C, D'Aniello B, Tsutsui K. (2017) Gonadotropin-inhibitory hormone (GnIH) in the amphibian brain and its relationship with the gonadotropin releasing hormone (GnRH) system: An overview. *Gen Comp Endocrinol.* 240:69-76.
- Lee Y, Dizzell SE, Leung V, Nazli A, Zahoor MA, Fichorova RN, Kaushic C. (2016) Effects of Female Sex Hormones on Susceptibility to HSV-2 in Vaginal Cells Grown in Air-Liquid Interface. *Viruses.* 30;8(9).
- Tian Y, Xu J, Feng S, He S, Zhao S, Zhu L, Jin W, Dai Y, Luo L, Qu JY, Wen Z. (2017) The first wave of T lymphopoiesis in zebrafish arises from aorta endothelium independent of hematopoietic stem cells. *J Exp Med.* 214(11):3347-3360.
- Igarashi M, Takasawa K, Hakoda A, Kanno J, Takada S, Miyado M, Baba T, Morohashi K, Tajima T, Hata K, Nakabayashi K, Matsubara Y, Sekido R, Ogata T, Kashimada K, Fukami M. (2017) Identical NR5A1 Missense Mutations in Two Unrelated 46,XX Individuals with Testicular Tissues. *Hum Mutat.* 38(1):39-42.
- Sawyer EM, Brunner EC, Hwang Y, Ivey LE, Brown O, Bannon M, Akrobetu D, Sheaffer KE, Morgan O, Field CO, Suresh N, Gordon MG, Gunnell ET, Regruto LA, Wood CG, Fuller MT, Hales KG. (2017) Testis-specific ATP synthase peripheral stalk subunits required for tissue-specific mitochondrial morphogenesis in *Drosophila*. *BMC Cell Biol.* 18(1):16.
- Rios-Rojas C, Spiller C, Bowles J, Koopman P. (2015) Germ cells influence cord formation and Leydig cell gene expression during mouse testis development. *Dev Dyn.* 2015 Dec 5. doi: 10.1002/dvdy.24371. [Epub ahead of print]
- Wang Y, Liu W, Yang Q, Yu M, Zhang Z. (2015) Di (2-ethylhexyl) phthalate exposure during pregnancy disturbs temporal sex determination regulation in mice offspring. *Toxicology.* 2;336:10-6.
- Zhang L, Chen M, Wen Q, et al. (2015) Reprogramming of Sertoli cells to fetal-like Leydig cells by Wt1 ablation. *Proc Natl Acad Sci U S A.* 112(13):4003-8.
- Grinspon RP, Rey RA. (2014) When hormone defects cannot explain it: malformative disorders of sex development. *Birth Defects Res C Embryo Today.* 102(4):359-73.
- Park S, Kim S, Jin H, Lee K, Bae J. (2014) Impaired development of female mouse offspring maternally exposed to simazine. *Environ Toxicol Pharmacol.* 38(3):845-51.
- Li L, Bu T, Su H, Chen Z, et al., (2015) In utero exposure to diisononyl phthalate caused testicular dysgenesis of rat fetal testis. *Toxicol Lett.* 22;232(2):466-74.
- Teerds KJ, Keijer J. (2015) Reduced fetal androgen exposure compromises Leydig cell function in adulthood. *Asian J Androl.* 17(2):219-20.
- Jones S, Boisvert A, Duong TB, et al., (2014) Disruption of rat testis development following combined in utero exposure to the phytoestrogen genistein and antiandrogenic plasticizer di-(2-ethylhexyl) phthalate. *Biol Reprod.* 91(3):64.
- Juul A, Almstrup K, Andersson AM, et al. (2014) Possible fetal determinants of male infertility. *Nat Rev Endocrinol.* 10(9):553-62.
- Minkina A, Matson CK, Lindeman RE, et al., (2014) DMRT1 protects male gonadal cells from retinoid-dependent sexual transdifferentiation. *Dev Cell.* 9;29(5):511-20.
- Liao HF, Chen WS, Chen YH, et al., (2014) DNMT3L promotes quiescence in postnatal spermatogonial progenitor cells. *Development.* 141(12):2402-13.
- Smith P, Wilhelm D, Rodgers RJ. (2014) Development of mammalian ovary. *J Endocrinol.* 221(3):R145-61.
- Skakkebaek NE, Rajpert-De Meyts E, Buck Louis GM, et al., (2016) Male Reproductive Disorders and Fertility Trends: Influences of Environment and Genetic Susceptibility. *Physiol Rev* 96(1):55-97.

- Salami SA, Raji Y. (2015) Generational reproductive outcomes in Wistar rats maternally exposed to *Ricinus communis* oil at different stages of gestation. *J Dev Orig Health Dis.* 6(5):443-53.
- Nicol B, Yao HH. (2015) Gonadal Identity in the Absence of Pro-Testis Factor SOX9 and Pro-Ovary Factor Beta-Catenin in Mice. *Biol Reprod.* 93(2):35.
- Aydođan Ahabab M, Barlas N. (2015) Influence of in utero di-n-hexyl phthalate and dicyclohexyl phthalate on fetal testicular development in rats. *Toxicol Lett.* 233(2):125-37.
- Chevalier N, Brucker-Davis F, Lahlou N, et al., (2015) A negative correlation between insulin-like peptide 3 and bisphenol A in human cord blood suggests an effect of endocrine disruptors on testicular descent during fetal development. *Hum Reprod.* 30(2):447-53.
- Scerbo MJ, Freire-Regatillo A, Cisternas CD, et al. (2014) Neurogenin 3 mediates sex chromosome effects on the generation of sex differences in hypothalamic neuronal development. *Front Cell Neurosci.* 8;8:188.
- Klinger FG, Rossi V, De Felici M. (2015) Multifaceted programmed cell death in the mammalian fetal ovary. *Int J Dev Biol.* 59(1-3):51-4.
- Gur EB, Karadeniz M, Turan GA. (2015) Fetal programming of polycystic ovary syndrome. *World J Diabetes.* 10;6(7):936-42.
- Connolly F, Rae MT, Späth K, Boswell L, McNeilly AS, Duncan WC. (2015) In an Ovine Model of Polycystic Ovary Syndrome (PCOS) Prenatal Androgens Suppress Female Fetal Renal Gluconeogenesis. *PLoS One.* 6;10(7):e0132113.
- Mu X, Liao X, Chen X, et al., (2015) DEHP exposure impairs mouse oocyte cyst breakdown and primordial follicle assembly through estrogen receptor-dependent and independent mechanisms. *J Hazard Mater.* 298:232-40.
- Bayne RA, Kinnell HL, Coutts SM, et al., (2015) GDF9 is transiently expressed in oocytes before follicle formation in the human fetal ovary and is regulated by a novel NOBOX transcript. *PLoS One.* 10(3):e0119819.
- Nakamura N, Inselman AL, White GA, et al., (2015) Effects of maternal and lactational exposure to 2-hydroxy-4-methoxybenzone on development and reproductive organs in male and female rat offspring. *Birth Defects Res B Dev Reprod Toxicol.* 104(1):35-51.
- Maliqueo M, Sundström Poromaa I, Vanky E, et al. (2015) Placental STAT3 signaling is activated in women with polycystic ovary syndrome. *Hum Reprod.* 30(3):692-700.
- Dutta S, Mark-Kappeler CJ, Hoyer PB, Pepling ME. (2014) The steroid hormone environment during primordial follicle formation in perinatal mouse ovaries. *Biol Reprod.* 91(3):68.
- Wainwright EN, Jorgensen JS, Kim Y, et al. (2013) SOX9 regulates microRNA miR-202-5p/3p expression during mouse testis differentiation. *Biol Reprod.* 15;89(2):34.
- Miyabayashi K, Katoh-Fukui Y, Ogawa H, et al. (2013) Aristaless related homeobox gene, *Arx*, is implicated in mouse fetal Leydig cell differentiation possibly through expressing in the progenitor cells. *PLoS One.* 28;8(6):e68050.
- Rakoczy J, Fernandez-Valverde SL, Glazov EA, et al. (2013) MicroRNAs-140-5p/140-3p modulate Leydig cell numbers in the developing mouse testis. *Biol Reprod.* 6;88(6):143.
- Song HW, Anderson RA, Bayne RA, et al. (2013) The RHOX homeobox gene cluster is selectively expressed in human oocytes and male germ cells. *Hum Reprod.* 28(6):1635-46.
- Muczynski V, Lecureuil C, Messiaen S, et al. (2012) Cellular and molecular effect of MEHP Involving LXR α in human fetal testis and ovary. *PLoS One.* 7(10):e48266.
- Huang Z, Rivas B, Agoulnik AI. (2012) Insulin-like 3 signaling is important for testicular descent but dispensable for spermatogenesis and germ cell survival in adult mice. *Biol Reprod.* 21;87(6):143.

- Miyado M, Nakamura M, Miyado K, et al. (2012) Maml1 deficiency significantly reduces mRNA expression levels of multiple genes expressed in mouse fetal Leydig cells but permits normal genital and reproductive development. *Endocrinology*. 153(12):6033-40.
- Klinefelter GR, Laskey JW, Winnik WM, et al. (2012) Novel molecular targets associated with testicular dysgenesis induced by gestational exposure to diethylhexyl phthalate in the rat: a role for estradiol. *Reproduction*. 144(6):747-61.
- Chen H, Palmer JS, Thiagarajan RD, et al. (2012) Identification of novel markers of mouse fetal ovary development. *PLoS One*. 2012;7(7):e41683.
- DiTacchio L, Bowles J, Shin S, et al. (2012) Transcription factors ER71/ETV2 and SOX9 participate in a positive feedback loop in fetal and adult mouse testis. *J Biol Chem*. 287(28):23657-66.
- Jameson SA, Natarajan A, Cool J, et al. (2012) Temporal transcriptional profiling of somatic and germ cells reveals biased lineage priming of sexual fate in the fetal mouse gonad. *PLoS Genet*. 8(3):e1002575.
- Sarraj MA, Drummond AE. (2012) Mammalian foetal ovarian development: consequences for health and disease. *Reproduction*. 143(2):151-63.
- Luense LJ, Veiga-Lopez A, Padmanabhan V, Christenson LK. (2011) Developmental programming: gestational testosterone treatment alters fetal ovarian gene expression. *Endocrinology*. 152(12):4974-83.
- Flück CE, Meyer-Böni M, Pandey AV, et al. (2011) Why boys will be boys: two pathways of fetal testicular androgen biosynthesis are needed for male sexual differentiation. *Am J Hum Genet*. 12;89(2):201-18.
- Kashimada K, Svingen T, Feng CW, et al. (2011) Antagonistic regulation of Cyp26b1 by transcription factors SOX9/SF1 and FOXL2 during gonadal development in mice. *FASEB J*. 25(10):3561-9.
- Lehraiki A, Chamailard C, Krust A, et al. (2011) Genistein impairs early testosterone production in fetal mouse testis via estrogen receptor alpha. *Toxicol In Vitro*. 25(8):1542-7.
- Sim H, Argentaro A, Czech DP, et al. (2011) Inhibition of SRY-calmodulin complex formation induces ectopic expression of ovarian cell markers in developing XY gonads. *Endocrinology*. 152(7):2883-93.
- Bagheri-Fam S, Argentaro A, Svingen T, et al. (2011) Defective survival of proliferating Sertoli cells and androgen receptor function in a mouse model of the ATR-X syndrome. *Hum Mol Genet*. 1;20(11):2213-24.
- DeFalco T, Takahashi S, Capel B. (2011) Two distinct origins for Leydig cell progenitors in the fetal testis. *Dev Biol*. 1;352(1):14-26.
- Kashimada K, Pelosi E, Chen H, et al. (2011) FOXL2 and BMP2 act cooperatively to regulate follistatin gene expression during ovarian development. *Endocrinology*. 152(1):272-80.
- Nicholas CR, Haston KM, Pera RA. (2010) Intact fetal ovarian cord formation promotes mouse oocyte survival and development. *BMC Dev Biol*. 8;10:2.
- Nilsson EE, Skinner MK. (2009) Progesterone regulation of primordial follicle assembly in bovine fetal ovaries. *Mol Cell Endocrinol*. 10;313(1-2):9-16.
- Lehraiki A, Racine C, Krust A, et al. (2009) Phthalates impair germ cell number in the mouse fetal testis by an androgen- and estrogen-independent mechanism. *Toxicol Sci*. 111(2):372-82.
- Wilhelm D, Washburn LL, Truong V, et al. (2009) Antagonism of the testis- and ovary-determining pathways during ovotestis development in mice. *Mech Dev*. 126(5-6):324-36.
- Gong W, Pan L, Lin Q, et al. (2013) Transcriptome profiling of the developing postnatal mouse testis using next-generation sequencing. *Sci China Life Sci*. 56(1):1-12.

- Zhang HQ, Zhang XF, Zhang LJ, Chao HH, et al. (2012) Fetal exposure to bisphenol A affects the primordial follicle formation by inhibiting the meiotic progression of oocytes. *Mol Biol Rep.* 39(5):5651-7.
- Itman C, Wong C, Hunyadi B, et al. (2011) Smad3 dosage determines androgen responsiveness and sets the pace of postnatal testis development. *Endocrinology.* 152(5):2076-89.
- Nandi SS, Ghosh P, Roy SS. (2011) Expression of PITX2 homeodomain transcription factor during gonadal development in a sexually dimorphic manner. *Cell Physiol Biochem.* 27(2):159-70.
- Wang C, Roy SK. (2010) Expression of E-cadherin and N-cadherin in perinatal hamster ovary: possible involvement in primordial follicle formation and regulation by follicle-stimulating hormone. *Endocrinology.* 151(5):2319-30.
- Mithraprabhu S, Mendis S, Meachem SJ, et al. (2010) Activin bioactivity affects germ cell differentiation in the postnatal mouse testis in vivo. *Biol Reprod.* 82(5):980-90.
- Lee KH, Park JH, Bunick D, et al. (2009) Morphological comparison of the testis and efferent ductules between wild-type and estrogen receptor alpha knockout mice during postnatal development. *J Anat.* 214(6):916-25.
- Garcia TX, DeFalco T, Capel B, Hofmann MC. (2013) Constitutive activation of NOTCH1 signaling in Sertoli cells causes gonocyte exit from quiescence. *Dev Biol.* 1;377(1):188-201.
- Maatouk DM, Mork L, Hinson A, et al. (2012) Germ cells are not required to establish the female pathway in mouse fetal gonads. *PLoS One.* 7(10):e47238.
- Chawengsaksophak K, Svingen T, Ng ET, et al. (2012) Loss of Wnt5a disrupts primordial germ cell migration and male sexual development in mice. *Biol Reprod.* 10;86(1):1-12.
- Rojas-García PP, Recabarren MP, Sarabia L, et al. (2010) Prenatal testosterone excess alters Sertoli and germ cell number and testicular FSH receptor expression in rams. *Am J Physiol Endocrinol Metab.* 299(6):E998-E1005.
- Spiller CM, Bowles J, Koopman P. (2012) Regulation of germ cell meiosis in the fetal ovary. *Int J Dev Biol.* 56(10-12):779-87.
- Sarraj MA, Drummond AE. (2012) Mammalian foetal ovarian development: consequences for health and disease. *Reproduction.* 143(2):151-63.
- Nilsson EE, Skinner MK. (2009) Progesterone regulation of primordial follicle assembly in bovine fetal ovaries. *Mol Cell Endocrinol.* 10;313(1-2):9-16.
- Wang C, Roy SK. (2009) Expression of bone morphogenetic protein receptor (BMPR) during perinatal ovary development and primordial follicle formation in the hamster: possible regulation by FSH. *Endocrinology.* 150(4):1886-96.
- Grützner F et al. (2008) Reproductive biology in egg-laying mammals. *Sex Dev.* 2008;2(3):115-27.
- Ireland JL et al. (2008) Antral follicle count reliably predicts number of morphologically healthy oocytes and follicles in ovaries of young adult cattle. *Biol Reprod.* 79(6):1219-25.
- Sun QY et al. (2008) Oocyte-specific knockout: a novel in vivo approach for studying gene functions during folliculogenesis, oocyte maturation, fertilization, and embryogenesis. *Biol Reprod.* 79(6):1014-20.
- Rodrigues P et al. (2008) Oogenesis: Prospects and challenges for the future. *J Cell Physiol.* 216(2):355-65.
- Richards JS et al. (2008) Immune-like mechanisms in ovulation. *Trends Endocrinol Metab.* Aug;19(6):191-6.
- Best D et al. (2008) Sdmg1 is a conserved transmembrane protein associated with germ cell sex determination and germline-soma interactions in mice. *Development.* 135(8):1415-25.

OVARY DEVELOPMENT

Fetal/Gonadogenesis

Melissa E Pepling and Joshua JN Burton, Syracuse University, Syracuse, NY, United States

© 2018 Elsevier Inc. All rights reserved.

Glossary

Anti-Müllerian hormone (AMH) A growth factor secreted by Sertoli cells to induce the degeneration of the Müllerian ducts in male embryos.

Coelomic epithelium Mesothelium derived from the splanchnopleura of lateral plate mesoderm that forms the lining of the organs in the coelomic cavity.

Folliculogenesis The growth, development, and maturation of an ovarian follicle.

Genital/gonadal ridge The anlagen of the testis in males and the ovary in females.

Germ cells Cells whose descendants become either sperm or egg.

Granulosa cells Ovary-specific cells that produce estrogen and progesterone, and support female germ cells throughout oogenesis.

Leydig cells The steroidogenic cells of the testis that synthesize testosterone.

Mesonephroi A pair of kidney-like organs derived from the intermediate mesoderm that function temporarily in the embryo, and from which the male and female reproductive tract are derived.

Oogonia Ovarian germ cells derived from PGCs that give rise to primary oocytes.

Primordial germ cell (PGC) Germ cell stage preceding sex differentiation.

Sertoli cells Testis-specific cells that secrete AMH in the developing embryo, and function to support male germ cells throughout spermatogenesis.

Sex-determining region Y (Sry) Gene encoding a transcription factor that initiates male sex determination; also known as testis determining factor (TDF).

Theca cells Endocrine cells associated with ovarian follicles that produce the androgen substrate for estrogen biosynthesis.

Introduction

Mammalian embryonic gonads initially form as a bipotential primordium that is morphologically indistinguishable in male and female embryos. A chromosomally determined sexual fate decision elicits the activation of the testis-specific or the ovary-specific pathway, and initiates the transformation of the common primordial gonads into either the male or female phenotype. The differentiated ovary produces and maintains a pool of oocytes, of which a subset is ovulated for fertilization. Additionally, the ovary functions as an endocrine organ, synthesizing and secreting the steroid hormones estrogen and progesterone that stimulate the development of secondary sex characteristics and prepare the accessory reproductive organs to facilitate pregnancy, birth, and lactation. Consequently, organogenesis of the ovaries is a highly important process for female fertility and reproductive success. Defects during gonad formation and development are pertinent to disorders of sex development (DSDs), and may also contribute to infertility or ovarian cancer. Knowledge of the process of gonadogenesis is therefore crucial to our understanding of reproductive pathologies, and may inform new treatment approaches for associated abnormalities. This article provides an overview of the key events that occur during the formation of the bipotential primordial gonad and the differentiation of the ovary, with particular focus on organogenesis of the ovary in mice, the most extensively studied mammalian model, and in humans.

The Urogenital System

Early in mammalian gestation, the embryo becomes trilaminar as it undergoes gastrulation to form three germ layers: the ectoderm, mesoderm, and endoderm. All embryonic tissues and organs are derived from these germ layers as the cells proliferate, migrate, and differentiate. The mesodermal layer is regionalized into the paraxial, intermediate, and lateral plate mesoderm. The intermediate mesoderm forms the urogenital system which includes the gonads and kidneys, as well as their associated duct systems. The urogenital ridges form bilaterally as longitudinal mesodermal elevations in the dorsal body wall of the embryo. Three sets of kidneys—the pronephroi (transitory and non-functional), mesonephroi (interim kidneys), and the metanephroi (becomes the permanent

kidneys)—develop from the urogenital ridges. Formation of the gonads is concurrent and closely associated with the differentiation of the mesonephroi. Invaginations of the surface epithelium of the mesonephroi give rise to the mesonephric (Wolffian) ducts and the paramesonephric (Müllerian) ducts. In male embryos, the testes produce testosterone which influences the differentiation of the Wolffian ducts to form the internal male genitalia, such as the vas deferens, epididymis, and the seminal vesicle. Additionally, the male gonad produces Müllerian-inhibiting substance (MIS), also known as anti-Müllerian hormone (AMH), which induces the degeneration of the Müllerian ducts. In contrast, the developing female gonad lacks these factors and the Müllerian ducts give rise to the fallopian tubes (oviducts), the uterus, the cervix, and the cranial portion of the vagina while the Wolffian ducts degenerate. Recent evidence suggests that the transcription factor, COUP-TFII, suppresses signaling that is required to maintain the Wolffian ducts and therefore actively regulates differentiation of the female reproductive tract (Zhao et al., 2017).

Formation of the Bipotential Gonad: The Indifferent Stage of Sexual Development

Gonadogenesis begins with the formation of the genital (gonadal) ridges which are the somatic precursor structures of both the testis and ovaries. The genital ridges form as a bulge or outgrowth of the coelomic epithelium on the ventromedial side of the mesonephroi by the fifth week of gestation in humans, and by embryonic day 10 (E10) in mice (Satoh, 1991; Waldeyer, 1870; Kaufman, 1992; Byskov, 1986). This proliferation of the coelomic epithelium and the underlying mesenchyme produces the somatic constituents of the gonads. There are at least two distinct bipotential somatic precursor lineages: (1) supporting cell precursors that differentiate into Sertoli cells in the testis or granulosa cells in the ovary, and (2) steroidogenic progenitors that produce Leydig cells in the testis or theca cells in the ovary (Karl and Capel, 1998; Schmahl et al., 2000). Stromal cells which are not specialized between sexes are also present at the genital ridge, and are responsible for structural patterning and angiogenic vascularization of the gonad. Genital ridge formation is therefore an indispensable prerequisite for development of both testis and ovaries, which remain bipotential at this stage until an intricate network of cellular signals begins to drive sexual differentiation.

Genetic Regulation of Gonad Formation

Several transcription factors and signaling proteins perform crucial regulatory roles in the development of the bipotential gonad. These were recently reviewed (Eggers et al., 2014, Tanaka and Nishinakamura, 2014), and are summarized here (also see Table 1). Steroidogenic factor 1 (*Sf1*), also known as *Nr5a1*, encodes an orphan nuclear hormone receptor that is expressed in both the gonad and steroidogenic tissues such as the adrenal glands (Lu and Yamashita, 2017). *Sf1* null mice exhibit agenesis of the gonads and the adrenal glands. *Sf1* is also an important regulator of *Sox9*, an essential gene for mediating testis differentiation. The Wilms tumor gene (*Wt1*) encodes a zinc finger transcription factor that is involved in the regulation of early gonad development, and differentiation of the testis (Kreidberg et al., 1993). *Wt1* null mice fail to undergo gonadogenesis, and their kidneys do not develop. Additionally, the *Wt1* protein has two isoforms, one lacking and one containing a KTS (lysine, threonine, and serine) amino acid motif. Mice lacking the *Wt1*(+KTS) isoform lose the ability to upregulate the testis-determining gene *Sry*, and exhibit male-to-female sex reversal. The *Wt1*(−KTS) isoform, in conjunction with *Lhx9*, directly activates the *Sf1* gene promoter and regulates the expression of testis-inducing factors such as *Amh*. This isoform is also important for the maintenance of the gonadal primordium. In humans, *WT1* mutations are also implicated in a form of kidney cancer that primarily affects children. *Lhx9* encodes the Lim/homeobox 9 protein which is another regulator of *Sf1* (Birk et al., 2000). *Lhx9* null mice also fail to undergo gonadogenesis. A deficiency in *Lhx9* contributes to the loss of *Amh* and testosterone, which causes XY mice to develop as females.

Table 1 Genes involved in mouse gonad formation

Gene	Protein/Function	Mutant phenotype	References
<i>Cbx2</i>	Chromobox, chromatin modification and remodeling factor	Impaired ovary development, male to female sex reversal.	Katoh-Fukui et al. (1998)
<i>Emx2</i>	Homeobox transcription factor	Lack kidney, uterine, gonad, reproductive tract.	Miyamoto et al. (1997)
<i>Gata4</i>	Zinc finger transcription factor	Inhibition of genital ridge formation.	Hu et al. (2013)
<i>Igfr1</i>	Insulin like growth factor receptor	Decreased proliferation of somatic progenitor cells.	Nef et al. (2003)
<i>Insr</i>	Insulin receptor	Decreased proliferation of somatic progenitor cells.	Nef et al. (2003)
<i>Lhx9</i>	Lim homeobox	Impaired gonadogenesis	Birk et al. (2000)
<i>Sf1</i>	Steroidogenic factor, orphan nuclear hormone receptor	No gonads or adrenal glands	Lu and Yamashita (2017)
<i>Six1/4</i>	Six family homeobox	Double knockouts have smaller gonads.	Kawakami et al. (2000)
<i>Wt1</i>	Zn finger transcription factor	No gonad development.	Kreidberg et al. (1993)

Empty spiracles homeobox 2 (Emx2) encodes a homeobox transcription factor involved in urogenital development (Miyamoto et al., 1997). *Emx2* null mice fail to develop kidneys, ureters, gonads, and genital tracts. Cellular polarity is also lost in the forming gonadal ridges of *Emx2* null mice leading to the abnormal assembly of tight junctions. Two other homeobox containing genes, *Six1* and *Six4*, are mammalian homologs of the *sine oculis homeobox (Six)* family in *Drosophila* (Kawakami et al., 2000). Both function redundantly in mouse embryogenesis. *Six1* and *Six4* double-mutant mouse embryos exhibit decreased gonad size in both sexes and abnormal differentiation of the testis in XY gonads.

The chromobox homolog 2 protein encoded by the *Cbx2* gene is a chromatin modification and remodeling factor that is involved in early gonadal development (Katoh-Fukui et al., 1998). *Cbx2* has been shown to upregulate *Sf1*, *Wt1*, and *Sry*. Deletion of *Cbx2* leads to impaired development of the ovary in XX mouse models, and male-to-female sex reversal in XY mice. *Cbx2* null mice appear to have normal development of the coelomic epithelium, but the gonadal cells later exhibit defective proliferation.

The GATA-binding protein 4 (*Gata4*) is a zinc finger transcription factor whose deletion is lethal to mouse embryos prior to genital ridge formation (Hu et al., 2013). Conditional knockdown of *Gata4* expression in mouse embryos after E8.75 causes the coelomic epithelium to remain as a morphologically undifferentiated monolayer, thereby preventing genital ridge formation. The insulin/insulin-like growth factor (IGF) signaling pathway is another crucial regulator of gonadogenesis (Nef et al., 2003). Mouse embryos without the insulin receptor (*Insr*) and the IGF receptor (*Igfr1*) have decreased proliferation of somatic progenitor cells in the gonads of both genotypic sexes prior to gonadal sex determination, and the testis fail to develop. Loss of insulin/IGF signaling can induce male-to-female sex reversal, and is also associated with reduced *Sry* expression levels. Proper spatiotemporal expression of each of these genes is imperative for establishing the somatic gonad.

Sex Determination

In mammals, dimorphic sex determination and differentiation progresses in three distinct phases. Firstly, the genetic sex of the embryo is determined at fertilization by the chromosomal complement of the zygote. Females are normally endowed with two X chromosomes, while males have an X and a Y sex chromosome. This genetic information facilitates gonadal sex determination during the second phase, by triggering growth of the bipotential gonad towards a testicular or ovarian fate. The short arm of the Y chromosome contains the gene encoding the mammalian testis-determining factor (TDF) known as *Sry* (sex-determining region Y). The spatiotemporal expression pattern of *Sry* is strictly controlled, and is first detected in the mouse gonad around E11, peaking at E11.5, before disappearing after E12.5 (Tanaka and Nishinakamura, 2014). *Sry* upregulates *Sox9*, and is both necessary and sufficient to direct the indifferent gonad towards the male phenotype in mice and humans (Koopman et al., 1991; Sinclair et al., 1990). *Sry* induces testicular morphogenesis by coordinating the differentiation of Sertoli cells and other testis-specific lineages. The presumptive ovarian equivalent of *Sry* has not yet been identified, and there is no evidence of a specific ovary determining factor. However, the ovary differentiation pathway is activated in the absence of functional *Sry* protein, and is primarily driven by the WNT/ β -catenin signaling pathway. Accordingly, although initial morphogenesis of the gonad is macroscopically similar between sexes, the gonadal transcriptome is notably different. The third stage, phenotypic sex determination, begins perinatally and continues throughout sexual maturation as the endocrine products of the gonads stimulate the development of secondary sex characteristics. Mutations or defects that compromise development during any of these three stages could lead to urogenital abnormalities or sex reversal.

The Ovary Differentiation Pathway

Differentiation of the bipotential gonads to the ovaries is not simply a passive process that occurs if there is no *Sry*. There are several ovary specific genes and signaling pathways that elicit female gonadal determination and development and this gonadal fate requires active repression of the testis pathway (Eggers et al., 2014, Chassot et al., 2014) (see Table 2). *Foxl2* encodes the forkhead box L2 transcription factor and is upregulated in the developing ovary (Uhlenhaut et al., 2009). Mutations of this gene in humans lead to blepharophimosis ptosis epicanthus inversus syndrome (BPES), which is characterized by eyelid malformation as well as, in some cases, primary ovarian insufficiency. Conditional knockdown of the *Foxl2* gene causes granulosa cells to become reprogrammed into Sertoli cells.

Table 2 Genes involved in mouse ovary differentiation

Gene	Protein/function	Mutant phenotype	References
<i>Ctnnb1</i>	β -catenin, Wnt signaling pathway	Ectopic expression causes male to female sex reversal.	Liu et al. (2009)
<i>Foxl2</i>	forkhead box transcription factor	Female to male sex reversal of somatic cells	Uhlenhaut et al. (2009)
<i>Fst</i>	Follistatin, activin antagonist	Mutants have female to male sex reversal.	Yao et al. (2004)
<i>Rspo1</i>	R-spondin homolog 1	Mutants have female to male sex reversal	Chassot et al. (2008)
<i>Wnt4</i>	Wnt signaling	Decreased proliferation of somatic progenitor cells.	Tomizuka et al. (2008)

β -Catenin, also known as Catenin beta-1 (Ctnnb1), transcriptionally regulates a variety of genes including *Wnt4* and *Fst* which are important for ovarian development (Liu et al., 2009). Ectopic expression of β -catenin in the developing XY mouse gonad causes male-to-female sex reversal, but the mechanism driving this trans-differentiation is unclear. WNT family members are conserved secreted proteins with roles in various biological processes. WNT3A participates in the stabilization of β -catenin and allows it to transcribe its target genes. WNT4 is known to activate canonical WNT signaling during gonadal development (Tomizuka et al., 2008). Females with heterozygous missense *WNT4* mutations exhibit agenesis of the reproductive tract, while *Wnt4* in the mouse is also important for gonadal morphogenesis via its role in sex determination and female development.

RSPO proteins activate the WNT/ β -catenin signaling pathway by binding to the LGR4, LGR5, and LGR6 G protein-coupled receptors, or by binding to ZNRF3 and RNF43 which are negative-feedback regulators of WNT signaling (Chassot et al., 2008). Disruption of human *RSPO1* can lead to female-to-male sex reversal, suggesting that it plays a crucial role in sex determination and female differentiation. At E12.5 in the mouse *Rspo1* and *Wnt4* become expressed in an ovary specific manner, and their gene products are secreted by the somatic cells of the ovary. *Rspo1* null mice exhibit sex reversal, bearing ovotestes. The *Rspo1*^{-/-} embryo demonstrates abnormalities in the XX gonad such as the presence of testis-like vasculature, and reduced germ cell proliferation. Similarly, a deficiency in *Wnt4* is associated with inducing male-like vascularization of the gonad in XX embryos. There is also evidence suggesting that *Wnt4* functions as a survival factor in female germ cells, as there are nearly three times as many apoptotic germ cells in the *Wnt4* deficient gonad, when compared to wild type gonads.

Follistatin (*Fst*) expression is stimulated by *Wnt4* in the XX gonad from E11.5 onwards, but is not expressed in the XY gonad (Yao et al., 2004). *Fst* encodes an activin-binding protein that antagonizes Activin B. This inhibitory action prevents male-like vascularization of the gonad. Subsequent development of the differentiated ovary will involve the production and development of the ovarian follicles.

Primordial Germ Cells: The Embryonic Precursors of Gametes

Primordial germ cells (PGCs) colonize the bipotential gonad in association with the somatic cells, and differentiate to produce haploid eggs or sperm. During gastrulation in mice, at around E7, inductive cell-cell interactions establish the germ line by specifying the allocation of PGCs. The genetic basis for germ cell specification is thought to involve the expression of the *fragilis* and *stella* genes in PGC founder cell lineages, allowing them to retain pluripotency while neighboring cells acquire a mesodermal fate (Saitou et al., 2002). Intercellular signaling involving Bone Morphogenetic Protein 4 (*Bmp4*) is also critical for initiating the formation of the germ line. *Bmp4* null mouse embryos lack PGCs and an allantois which are both derived from the proximal epiblast (Lawson et al., 1999). The *Bmp* signaling pathway regulates the expression of two key genes that are essential for germ cell fate: PR domain containing protein 1 (*Prdm1*) which acts to repress the somatic cell fate, and PR domain containing protein 14 (*Prdm14*) which transcriptionally regulates germ cell pluripotency and epigenetic reprogramming (Windley and Wilhelm, 2015).

PGCs migrate via the hindgut and the dorsal mesentery to the developing gonadal ridges from approximately E9.5 to E11.5 in the mouse and from the fifth to seventh week of gestation in humans (Clark and Eddy, 1975; Fujimoto et al., 1977). During this time, the number of PGCs increases from approximately 40 to 250 in mice, and from approximately 100 to 5000 in humans (Cummings and Kavlock, 2004; Saitou and Yamaji, 2012). The PGCs rapidly proliferate by mitosis and increase in number to colonize the developing gonads as oogonia or spermatogonia. This proliferation produces a maximum of almost 7 million germ cells by the 5th month of gestation in humans, and around 25,000 germ cells by E13.5 in mice. At E11.5 in the mouse embryo and in the seventh to ninth week of gestation in humans, PGCs display similar expression patterns of germline and pluripotency genes. These germ cell specific genes include *BLIMP1*, *AP2 γ* , *UTF1*, *DAZL*, *Kit*, and *DDX4*, while the pluripotency genes include *OCT4*, *NANOG*, *PRDM14*, and *LIN28* (Dolci et al., 2015). The KIT receptor and its ligand KITL, are well known for promoting cell survival and proliferation during gametogenesis. Loss of KIT function impairs PGC proliferation and migration, results in ectopic PGCs, and affects the growth of early ovarian follicles (Buehr et al., 1993; Huang et al., 1993; McCoshen and McCallion, 1975; Mintz and Russell, 1957). By E13.5 in mice and by the tenth week of gestation in humans, oogonia begin to enter meiosis and are called oocytes. Developing oocytes arrest in the diplotene stage of the first meiotic prophase until ovulation begins during puberty, while spermatogonia enter a state of quiescence and reinitiate development at the onset of puberty.

Development of the Ovaries

The differentiated ovary displays notable complexity in structure and function, and is histologically identifiable around the twelfth week of gestation in humans and E13.5 in mice, by the presence of loose cordlike structures termed ovigerous cords (Loffler and Koopman, 2002; Odor and Blandau, 1969). Ovigerous cords consist of clusters of primordial germ cells that are surrounded by somatic cells. Ovarian PGCs in these clusters are physically interconnected by cytoplasmic bridges, and develop synchronously as germ cell cysts (Pepling and Spradling, 1998). Cysts subsequently undergo programmed breakdown to form primordial follicles, consisting of an individual germ cell or oocyte surrounded by a monolayer of somatic granulosa cells (Pepling and Spradling,

2001). Somatic cells extend thin cytoplasmic prolongations between germ cells in ovigerous cords, thereby facilitating cyst breakdown and follicle assembly (Odor and Blandau, 1969; Pepling and Spradling, 2001). The features of early oocyte development are very similar in mice and humans, with the main difference being that cyst breakdown and primordial follicle formation occur perinatally in the mouse, but before birth in humans (Sarraj and Drummond, 2012).

References

- Birk, O. S., Casiano, D. E., Wassif, C. A., Cogliati, T., Zhao, L., Zhao, Y., Grinberg, A., Huang, S., Kreidberg, J. A., Parker, K. L., Porter, F. D., & Westphal, H. (2000). The LIM homeobox gene *Lhx9* is essential for mouse gonad formation. *Nature*, *403*, 909–913.
- Buehr, M., McLaren, A., Bartley, A., & Darling, S. (1993). Proliferation and migration of primordial germ cells in We/We mouse embryos. *Developmental Dynamics*, *198*, 182–189.
- Byskov, A. G. (1986). Differentiation of mammalian embryonic gonad. *Physiological Reviews*, *66*, 71–117.
- Chassot, A. A., Ranc, F., Gregoire, E. P., Roepers-Gajadien, H. L., Taketo, M. M., Camerino, G., De Rooij, D. G., Schedl, A., & Chaboissier, M. C. (2008). Activation of beta-catenin signaling by *Rspo1* controls differentiation of the mammalian ovary. *Human Molecular Genetics*, *17*, 1264–1277.
- Chassot, A. A., Gillot, I., & Chaboissier, M. C. (2014). R-spondin1, WNT4, and the CTNNB1 signaling pathway: Strict control over ovarian differentiation. *Reproduction*, *148*, R97–110.
- Clark, J. M., & Eddy, E. M. (1975). Fine structural observations on the origin and associations of primordial germ cells of the mouse. *Developmental Biology*, *47*, 136–155.
- Cummings, A. M., & Kavlock, R. J. (2004). Function of sexual glands and mechanism of sex differentiation. *The Journal of Toxicological Sciences*, *29*, 167–178.
- Dolci, S., Campolo, F., & De Felici, M. (2015). Gonadal development and germ cell tumors in mouse and humans. *Seminars in Cell & Developmental Biology*, *45*, 114–123.
- Eggers, S., Ohnesorg, T., & Sinclair, A. (2014). Genetic regulation of mammalian gonad development. *Nature Reviews: Endocrinology*, *10*, 673–683.
- Fujimoto, T., Miyayama, Y., & Fuyuta, M. (1977). The origin, migration and fine morphology of human primordial germ cells. *The Anatomical Record*, *188*, 315–330.
- Hu, Y. C., Okumura, L. M., & Page, D. C. (2013). *Gata4* is required for formation of the genital ridge in mice. *PLoS Genetics*, *9*, e1003629.
- Huang, E. J., Manova, K., Packer, A. I., Sanchez, S., Bachvarova, R. F., & Besmer, P. (1993). The murine steel panda mutation affects kit ligand expression and growth of early ovarian follicles. *Developmental Biology*, *157*, 100–109.
- Karl, J., & Capel, B. (1998). Sertoli cells of the mouse testis originate from the coelomic epithelium. *Developmental Biology*, *203*, 323–333.
- Katoh-Fukui, Y., Tsuchiya, R., Shiroishi, T., Nakahara, Y., Hashimoto, N., Noguchi, K., & Higashinakagawa, T. (1998). Male-to-female sex reversal in M33 mutant mice. *Nature*, *393*, 688–692.
- Kaufman, M. H. (1992). *The atlas of mouse development*. London: Academic Press.
- Kawakami, K., Sato, S., Ozaki, H., & Ikeda, K. (2000). Six family genes—structure and function as transcription factors and their roles in development. *BioEssays*, *22*, 616–626.
- Koopman, P., Gubbay, J., Vivian, N., Goodfellow, P., & Lovell-Badge, R. (1991). Male development of chromosomally female mice transgenic for *Sry*. *Nature*, *351*, 117–121.
- Kreidberg, J. A., Sariola, H., Loring, J. M., Maeda, M., Pelletier, J., Housman, D., & Jaenisch, R. (1993). *WT-1* is required for early kidney development. *Cell*, *74*, 679–691.
- Lawson, K. A., Dunn, N. R., Roelen, B. A., Zeinstra, L. M., Davis, A. M., Wright, C. V., Korving, J. P., & Hogan, B. L. (1999). *Bmp4* is required for the generation of primordial germ cells in the mouse embryo. *Genes & Development*, *13*, 424–436.
- Liu, C. F., Bingham, N., Parker, K., & Yao, H. H. (2009). Sex-specific roles of beta-catenin in mouse gonadal development. *Human Molecular Genetics*, *18*, 405–417.
- Loffler, K. A., & Koopman, P. (2002). Charting the course of ovarian development in vertebrates. *The International Journal of Developmental Biology*, *46*, 503–510.
- Lu, K. L., & Yamashita, Y. M. (2017). Germ cell connectivity enhances cell death in response to DNA damage in the *Drosophila* testis. *eLife*, *6*, e27960.
- McCoshen, J. A., & McCallion, D. J. (1975). A study of the primordial germ cells during their migratory phase in steel mutant mice. *Experientia*, *31*, 589–590.
- Mintz, B., & Russell, E. S. (1957). Gene-induced embryological modifications of primordial germ cells in the mouse. *The Journal of Experimental Zoology*, *134*, 207–237.
- Miyamoto, N., Yoshida, M., Kuratani, S., Matsuo, I., & Aizawa, S. (1997). Defects of urogenital development in mice lacking *Emx2*. *Development*, *124*, 1653–1664.
- Nef, S., Verma-Kurvari, S., Merenmies, J., Vassalli, J. D., Efstratiadis, A., Accilli, D., & Parada, L. F. (2003). Testis determination requires insulin receptor family function in mice. *Nature*, *426*, 291–295.
- Odor, D. L., & Blandau, R. J. (1969). Ultrastructural studies on fetal and early postnatal mouse ovaries. I. Histogenesis and organogenesis. *The American Journal of Anatomy*, *124*, 163–186.
- Pepling, M. E., & Spradling, A. C. (1998). Female mouse germ cells form synchronously dividing cysts. *Development*, *125*, 3323–3328.
- Pepling, M. E., & Spradling, A. C. (2001). Mouse ovarian germ cell cysts undergo programmed breakdown to form primordial follicles. *Developmental Biology*, *234*, 339–351.
- Saitou, M., & Yamaji, M. (2012). Primordial germ cells in mice. *Cold Spring Harbor Perspectives in Biology*, *4*, a008375.
- Saitou, M., Barton, S. C., & Surani, M. A. (2002). A molecular programme for the specification of germ cell fate in mice. *Nature*, *418*, 293–300.
- Sarraj, M. A., & Drummond, A. E. (2012). Mammalian foetal ovarian development: Consequences for health and disease. *Reproduction*, *143*, 151–163.
- Satoh, M. (1991). Histogenesis and organogenesis of the gonad in human embryos. *Journal of Anatomy*, *177*, 85–107.
- Schmahl, J., Eicher, E. M., Washburn, L. L., & Capel, B. (2000). *Sry* induces cell proliferation in the mouse gonad. *Development*, *127*, 65–73.
- Sinclair, A. H., Berta, P., Palmer, M. S., Hawkins, J. R., Griffiths, B. L., Smith, M. J., Foster, J. W., Frischauf, A. M., Lovell-Badge, R., & Goodfellow, P. N. (1990). A gene from the human sex-determining region encodes a protein with homology to a conserved DNA-binding motif. *Nature*, *346*, 240–244.
- Tanaka, S. S., & Nishinakamura, R. (2014). Regulation of male sex determination: Genital ridge formation and *Sry* activation in mice. *Cellular and Molecular Life Sciences*, *71*, 4781–4802.
- Tomizuka, K., Horikoshi, K., Kitada, R., Sugawara, Y., Iba, Y., Kojima, A., Yoshitome, A., Yamawaki, K., Amagai, M., Inoue, A., Oshima, T., & Kakitani, M. (2008). R-spondin1 plays an essential role in ovarian development through positively regulating Wnt-4 signaling. *Human Molecular Genetics*, *17*, 1278–1291.
- Uhlenhaut, N. H., Jakob, S., Anlag, K., Eisenberger, T., Sekido, R., Kress, J., Treier, A. C., Klugmann, C., Klasen, C., Holter, N. I., Riethmacher, D., Schutz, G., Cooney, A. J., Lovell-Badge, R., & Treier, M. (2009). Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation. *Cell*, *139*, 1130–1142.
- Waldeyer, W. (1870). *Eierstock und Ei*. Leipzig: Engelmann.
- Windley, S. P., & Wilhelm, D. (2015). Signaling pathways involved in mammalian sex determination and gonad development. *Sexual Development*, *9*, 297–315.
- Yao, H. H., Matzuk, M. M., Jorgez, C. J., Menke, D. B., Page, D. C., Swain, A., & Capel, B. (2004). Follistatin operates downstream of Wnt4 in mammalian ovary organogenesis. *Developmental Dynamics*, *230*, 210–215.
- Zhao, F., Franco, H. L., Rodriguez, K. F., Brown, P. R., Tsai, M. J., Tsai, S. Y., & Yao, H. H. (2017). Elimination of the male reproductive tract in the female embryo is promoted by COUP-TFII in mice. *Science*, *357*, 717–720.

DEVELOPMENT

Formation of the Testis Primordium

Emmi Rotgers and Humphrey H-C Yao, National Institute of Environmental Health Sciences (NIEHS/NIH), Research Triangle Park, NC, United States

© 2018 Elsevier Inc. All rights reserved.

Formation of the Testis From the Bipotential Gonad

The testes form from a primitive undifferentiated gonad or genital ridge that has the potential to become either testes or ovary. In this article, we will describe the earliest events in testis development, when gonadogenesis, or formation of the gonads, occurs. Because most of the experimental evidence comes from studies using the mouse as a model organism, the morphological and molecular events during gonad formation are discussed mostly with reference to mouse development.

A functional gonad requires the presence of germ cells and somatic cells in order to produce gametes and hormones. The somatic cells of the gonads originate from the coelomic epithelium, a thin layer of epithelial cells that covers the gonad. While the primitive gonad is establishing, germ cell precursors, or primordial germ cells, migrate into the gonad from elsewhere in the embryo (discussed in the second part of this article). Key steps in forming the gonad are (1) proliferation of the coelomic epithelial cells, (2) ingression of the coelomic epithelium into the gonad proper, and (3) subsequent expansion and sexual differentiation of the somatic cells (Fig. 1). This process is controlled by a group of transcription factors and signaling molecules. At the onset of gonadogenesis, the coelomic epithelial cells that are destined to form the gonad become positive for the transcription factor Wilm's tumor 1 (WT1) (Kreidberg et al., 1993). Without *Wt1*, thickening of the coelomic epithelium is reduced dramatically (Kreidberg et al., 1993). The coelomic epithelium proliferates in an anterior to posterior fashion. The anterior-posterior thickening of the coelomic epithelium and the subsequent establishment of the gonad require the action of another transcription factor GATA binding protein 4 (GATA4) (Hu et al., 2013). GATA4 expression in the anterior part of the gonad precedes the initial thickening of the coelomic epithelium, and without *Gata4*, proliferation of the coelomic epithelial cells, and subsequent formation of the gonad fail to occur (Hu et al., 2013). The thickening of the coelomic epithelium is followed by a breakdown of its underlying basement membrane and ingression of the coelomic epithelial cells to form the gonad proper. In addition to WT1 and GATA4, the coelomic epithelial cells also express steroidogenic factor 1 (SF1), an orphan nuclear receptor that is important for gonad formation (Hoivik et al., 2010). Without functional SF1, the gonad does not expand properly and the gonadal cells eventually die (Luo et al., 1994). The early SF1 expression in the coelomic epithelium is promoted by GATA4 and two homeodomain proteins SIX1 and SIX4 (sine oculis homeobox homologs 1 and 4) (Hoivik et al., 2010; Fujimoto et al., 2013).

After its initial thickening, the coelomic epithelium continues to proliferate and provides progenitors to the developing gonad. This proliferation of the coelomic epithelial cells is promoted by Lim homeobox 9 (LHX9) and insulin signaling. Loss of *Lhx9* or insulin receptor and IGF type I receptor (*Igf1r*) result in stunted gonad development due to decreased proliferation of gonadal somatic cells (Birk et al., 2000; Pitetti et al., 2013). What the coelomic epithelial cells become after proliferation is determined by the polarity of the cells. The coelomic epithelium proliferates asymmetrically, where the progenitor cells remain as a part of the coelomic epithelium and maintain an undifferentiated epithelial cell fate. The daughter cells, on the other hand, ingress to the gonad proper, and acquire mesenchymal characteristics. During the ingression process, epithelial cells lose their cell polarity and cell-cell connections, and migrate through the basement membrane. Once they pass through the basement membrane, the ingressing epithelial cells form clusters that become the future somatic cells in the testis (Fig. 1). Multiple factors and pathways are involved in the invagination of the coelomic epithelium. Homeodomain transcription factor EMX2 and possibly SIX1/SIX4 are critical for the proliferation and the subsequent migration of the coelomic epithelial cells through the basement membrane. Without *Emx2*, the coelomic epithelial cells accumulate on the gonadal surface and fail to migrate and differentiate (Kusaka et al., 2010). A tight regulation of Notch signaling pathway is also necessary to maintain the asymmetric divisions and proper balance between progenitor and differentiated cell populations. A defect in the Notch signaling pathway results in abnormal accumulation of the progenitors and a failure of them to differentiate into the somatic cells in the testis (Potter et al., 2016; Lin et al., 2017). The coelomic epithelial cells that enter the gonad during sex determination become Sertoli cells. Proliferation and ingression of coelomic epithelial cells continue after sex determination, but cells ingressing at this time contribute to fetal and adult Leydig cell populations in the interstitium (Karl and Capel, 1998).

After the gonad has formed, it is important to ensure that new somatic cell progenitor cells survive and proliferate to form the testes. The survival of gonadal cells is controlled by SF1 and WT1. In the absence *Sf1* and *Wt1*, the coelomic epithelium begins to thicken during the earliest steps of gonad development, but then these structures gradually disappear through apoptosis of the gonadal somatic cells (Luo et al., 1994; Kreidberg et al., 1993). In addition to survival, sufficient proliferation of gonadal somatic cells is necessary for testis development (Schmahl and Capel, 2003). Without proliferation of the somatic cells in this early phase, such as in knockout mouse models for *Pbx1* and *Cbx2*, the resulting gonads are very small and poorly formed (Schnabel et al., 2003; Katoh-Fukui et al., 1998).




Fig. 1 Formation of the gonad. (A) The coelomic epithelium (CE), which lies on top of the future gonad, has an intact basement membrane prior to gonad formation. (B) Coelomic epithelial cells begin to express GATA4, prior to proliferation and thickening of the coelomic epithelium. (C) The cells that are destined to become gonadal somatic cells begin to express SF1. The basement membrane disintegrates and cells begin to ingress into the gonad. (D) Supporting cell progenitors (pre-Sertoli cells) continue to enter the gonad and proliferate. Primordial germ cells arrive to the gonad and they are surrounded by the pre-Sertoli cells. (E) The proliferating precursor cells from the coelomic epithelium contribute also to interstitial cells. Modified from Piprek, R. P., Kloc, M., and Kubiak, J. Z. 2016. Early development of the gonads: Origin and differentiation of the somatic cells of the genital ridges. *Results and Problems in Cell Differentiation* **58**, 1–22.

Development of the genital ridges lays the foundation for testicular development by ensuring that sufficient numbers of Sertoli cell precursors are established. Once the testis primordium is established, testicular development continues with migration of interstitial cell progenitors, vascular cells and macrophages into the gonad, which is covered in detail in other articles.

Primordial Germ Cell Formation and Migration

While the gonad is forming, primordial germ cells that migrate from the hindgut begin to enter the gonad. Primordial germ cells first gain their identity as a few cells in the epiblast, an embryonic structure next to the embryonic ectoderm (Fig. 2A). Primordial germ




Fig. 2 Specification and migration of primordial germ cells. (A) Primordial germ cells are first specified in the epiblast. This process is controlled by BMP from the extraembryonic ectoderm and WNT3 signaling from the visceral endoderm. (B) As the embryo develops the primordial germ cells migrate to the base of the allantois. (C) When primordial germ cells begin their long migration toward the gonads, they leave the allantois and enter the hindgut. They migrate through the hindgut and mesentery until they reach the gonads. Arrows denote direction of migration. Modified from Tang, W. W., Kobayashi, T., Irie, N., Dietmann, S., and Surani, M. A. 2016. Specification and epigenetic programming of the human germ line. *Nature Reviews. Genetics* **17**, 585–600; Rosen, M. P., and Cedars, M. I. 2011. Female reproductive endocrinology and infertility, In D. G. Gardner and D. Shoback (eds.), *Greenspan's basic & Clinical endocrinology*, 9th edn. (New York, NY: The McGraw-Hill Companies). Chapter 13.

cell specification is directed by cues from the adjacent structures: bone morphogenetic protein (BMP) signaling from the extraembryonic ectoderm and Wnt family member 3 (WNT3) from the visceral endoderm (Tang et al., 2016). These signals are sensed by the presumptive primordial germ cell progenitors, and activate genes that distinguish them from somatic cells.

As the embryo develops, primordial germ cells relocate to the extraembryonic region at the base of the allantois, a membrane that is important for the development of the umbilical cord and placenta. From the allantois, primordial germ cells migrate through the hindgut and mesentery toward the gonads (Fig. 2B). Several factors promote primordial germ cell survival during the long migration process, such as DND microRNA-mediated repression inhibitor 1 (DND1), and Kit ligand (KitL) and stromal cell derived factor 1 (SDF1, also known as CXCL12) (Youngren et al., 2005; Gu et al., 2009; Molyneaux et al., 2003).

There are two theories on the migration process of the primordial germ cells: active migration of the primordial germ cells toward the gonads, which is induced by signals secreted by cells along the migratory path, or passive movement of primordial germ cells due to physical changes of the embryonic structure (Harikae et al., 2013). According to the theory of active migration, primordial germ cells sense chemoattractant signals secreted by somatic cells in the gonad, adjacent mesonephros, and mesentery. In response to these signals with the specific receptors, primordial germ cells actively migrate through the hindgut toward the gonads. Examples of chemoattractant signalling pathways are Kit ligand and c-Kit receptor (also known as Steel and KIT, respectively), and SDF1 and its receptor CXCR4 (chemokine (CXC motif) receptor 4). Kit pathway is involved in primordial germ cell proliferation, survival, and migration throughout the migration period (Gu et al., 2009). Without Kit ligand or its receptor, mice have reduced proliferation of migrating primordial germ cells during fetal life and are sterile as adults. Kit ligands elicit their action within a short range, meaning that the target cell has to be close to the cell secreting the Kit ligand. Interestingly, cells that produce Kit ligand form a path which migrating primordial germ cells follow. Kit ligand-expressing cells surround primordial germ cells that are already in the allantois right after primordial germ cells have become specified (Gu et al., 2009). As migration begins, Kit ligand is expressed both in the gonads and the midline area through which primordial germ cells migrate. Later on, when primordial germ cells reach the gonads, Kit ligand expression is switched off in the migratory path outside the gonads, but maintained in the gonads. Kit ligand not only promotes primordial germ cell survival and proliferation, but also influences their motility. Without Kit ligand, primordial germ cells migrate to the correct direction but at a lower rate than normal, resulting in fewer primordial germ cells reaching the gonads (Gu et al., 2009).

There are several similarities between the roles of Kit/Kit ligand and the SDF1/CXCR signaling pathways in controlling primordial germ cell migration. If SDF1 or its receptor CXCR4 are deleted in mice, very few germ cells reach the gonads and the adult mice are sterile (Ara et al., 2003; Molyneaux et al., 2003). SDF1 also promotes primordial germ cell survival and is expressed in the somatic cells along the migratory path of primordial germ cells in a similar fashion as Kit ligand. However, unlike Kit ligand, SDF1 is involved in establishing the direction of primordial germ cell migration. If ectopic SDF1 is present in embryonic organs other than the gonads, primordial germ cells begin to migrate toward that region erroneously (Molyneaux et al., 2003). Also, in

contrast to Kit ligand which is necessary for primordial germ cell migration from start to finish, SDF1 is dispensable for the early stages of migration, but required for the later stages of migration through the mesentery and hindgut to the gonads (Molyneux et al., 2003; Ara et al., 2003). In conclusion, KIT and SDF1/CXCR4 signaling pathways play complementary roles in primordial germ cell proliferation, survival, and migration.

The second theory for primordial germ cell migration is the passive theory: when primordial germ cells are first incorporated into the hindgut endoderm, proliferation, and expansion of the hindgut passively propels primordial germ cells forward and toward the gonad (Harikae et al., 2013). If the hindgut does not expand, primordial germ cells are stuck at the hindgut entrance. It has been proposed that both of the active and passive theories are involved: the initial long-range migration of the primordial germ cells is promoted by passive translocation, but the survival and final steps of migration before reaching the gonads are promoted by chemoattractants secreted from the gonads. Interestingly, migration of primordial germ cells to the correct site still occurs even though formation of the gonads is compromised (Hu et al., 2013; Kreidberg et al., 1993). An explanation for this could be that production of chemoattractants in not only the gonads but also the adjacent mesentery and mesonephros is sufficient to guide the migrating primordial germ cells to their final destination.

After primordial germ cells reach the gonads, they cease migration and begin to form cell-to-cell associations with the somatic cells. It is hypothesized that reaching the site of highest concentration of the chemoattractants prompts primordial germ cells to stop migrating (Richardson and Lehmann, 2010). In addition, gonadal somatic cells likely suppress primordial germ cell migration, but the molecular mechanisms are not known. In the fetal testis, primordial germ cells become surrounded by Sertoli cells, which are responsible for the formation of testis cords. While they are enclosed in the testis cords, germ cells undergo rapid proliferation, form aggregates called germ cell cysts, and then cease mitosis (Lei and Spradling, 2013). Germ cell development that follows formation of fetal testes is discussed in detail in the following article on spermatogonial development and spermatogenesis.

References

- Ara, T., Nakamura, Y., Egawa, T., Sugiyama, T., Abe, K., Kishimoto, T., Matsui, Y., & Nagasawa, T. (2003). Impaired colonization of the gonads by primordial germ cells in mice lacking a chemokine, stromal cell-derived factor-1 (SDF-1). *Proceedings of the National Academy of Sciences of the United States of America*, *100*, 5319–5323.
- Birk, O. S., Casiano, D. E., Wassif, C. A., Cogliati, T., Zhao, L., Zhao, Y., Grinberg, A., Huang, S., Kreidberg, J. A., Parker, K. L., Porter, F. D., & Westphal, H. (2000). The LIM homeobox gene *Lhx9* is essential for mouse gonad formation. *Nature*, *403*, 909–913.
- Fujimoto, Y., Tanaka, S. S., Yamaguchi, Y. L., Kobayashi, H., Kuroki, S., Tachibana, M., Shinomura, M., Kanai, Y., Morohashi, K., Kawakami, K., & Nishinakamura, R. (2013). Homeoproteins *Six1* and *Six4* regulate male sex determination and mouse gonadal development. *Developmental Cell*, *26*, 416–430.
- Gu, Y., Runyan, C., Shoemaker, A., Surani, A., & Wylie, C. (2009). Steel factor controls primordial germ cell survival and motility from the time of their specification in the allantois, and provides a continuous niche throughout their migration. *Development*, *136*, 1295–1303.
- Harikae, K., Miura, K., & Kanai, Y. (2013). Early gonadogenesis in mammals: Significance of long and narrow gonadal structure. *Developmental Dynamics*, *242*, 330–338.
- Hoivik, E. A., Lewis, A. E., Aumo, L., & Bakke, M. (2010). Molecular aspects of steroidogenic factor 1 (SF-1). *Molecular and Cellular Endocrinology*, *315*, 27–39.
- Hu, Y. C., Okumura, L. M., & Page, D. C. (2013). *Gata4* is required for formation of the genital ridge in mice. *PLoS Genetics*, *9*, e1003629.
- Karl, J., & Capel, B. (1998). Sertoli cells of the mouse testis originate from the coelomic epithelium. *Developmental Biology*, *203*, 323–333.
- Katoh-Fukui, Y., Tsuchiya, R., Shiroishi, T., Nakahara, Y., Hashimoto, N., Noguchi, K., & Higashinakagawa, T. (1998). Male-to-female sex reversal in *M33* mutant mice. *Nature*, *393*, 688–692.
- Kreidberg, J. A., Sariola, H., Loring, J. M., Maeda, M., Pelletier, J., Housman, D., & Jaenisch, R. (1993). *WT-1* is required for early kidney development. *Cell*, *74*, 679–691.
- Kusaka, M., Katoh-Fukui, Y., Ogawa, H., Miyabayashi, K., Baba, T., Shima, Y., Sugiyama, N., Sugimoto, Y., Okuno, Y., Kodama, R., Iizuka-Kogo, A., Senda, T., Sasaoka, T., Kitamura, K., Aizawa, S., & Morohashi, K. (2010). Abnormal epithelial cell polarity and ectopic epidermal growth factor receptor (EGFR) expression induced in *Emx2* KO embryonic gonads. *Endocrinology*, *151*, 5893–5904.
- Lei, L., & Spradling, A. C. (2013). Mouse primordial germ cells produce cysts that partially fragment prior to meiosis. *Development*, *140*, 2075–2081.
- Liu, Y. T., Barske, L., DeFalco, T., & Capel, B. (2017). *Numb* regulates somatic cell lineage commitment during early gonadogenesis in mice. *Development*, *144*, 1607–1618.
- Luo, X., Ikeda, Y., & Parker, K. L. (1994). A cell-specific nuclear receptor is essential for adrenal and gonadal development and sexual differentiation. *Cell*, *77*, 481–490.
- Molyneux, K. A., Zinszner, H., Kunwar, P. S., Schaible, K., Stebler, J., Sunshine, M. J., O'Brien, W., Raz, E., Littman, D., Wylie, C., & Lehmann, R. (2003). The chemokine SDF1/CXCL12 and its receptor CXCR4 regulate mouse germ cell migration and survival. *Development*, *130*, 4279–4286.
- Pitetti, J. L., Calvel, P., Romero, Y., Conne, B., Truong, V., Papaioannou, M. D., Schaad, O., Docquier, M., Herrera, P. L., Wilhelm, D., & Nef, S. (2013). Insulin and IGF1 receptors are essential for XX and XY gonadal differentiation and adrenal development in mice. *PLoS Genetics*, *9*, e1003160.
- Potter, S. J., Kumar, D. L., & DeFalco, T. (2016). Origin and differentiation of androgen-producing cells in the gonads. *Results and Problems in Cell Differentiation*, *58*, 101–134.
- Richardson, B. E., & Lehmann, R. (2010). Mechanisms guiding primordial germ cell migration: Strategies from different organisms. *Nature Reviews. Molecular Cell Biology*, *11*, 37–49.
- Schmahl, J., & Capel, B. (2003). Cell proliferation is necessary for the determination of male fate in the gonad. *Developmental Biology*, *258*, 264–276.
- Schnabel, C. A., Selleri, L., & Cleary, M. L. (2003). *Pbx1* is essential for adrenal development and urogenital differentiation. *Genesis*, *37*, 123–130.
- Tang, W. W., Kobayashi, T., Irie, N., Dietmann, S., & Surani, M. A. (2016). Specification and epigenetic programming of the human germ line. *Nature Reviews. Genetics*, *17*, 585–600.
- Youngren, K. K., Coveney, D., Peng, X., Bhattacharya, C., Schmidt, L. S., Nickerson, M. L., Lamb, B. T., Deng, J. M., Behringer, R. R., Capel, B., Rubin, E. M., Nadeau, J. H., & Matin, A. (2005). The *Ter* mutation in the *dead end* gene causes germ cell loss and testicular germ cell tumours. *Nature*, *435*, 360–364.

Further Reading

- Piprek, R. P., Kloc, M., & Kubiak, J. Z. (2016). Early development of the gonads: Origin and differentiation of the somatic cells of the genital ridges. *Results and Problems in Cell Differentiation*, *58*, 1–22.
- Rosen, M. P., & Cedars, M. I. (2011). Female reproductive endocrinology and infertility. In D. G. Gardner, & D. Shoback (Eds.), *Greenspan's basic & clinical endocrinology* (9th edn). New York, NY: The McGraw-Hill Companies. Chapter 13.

"Systems Biology of Reproduction"

Spring 2024 (Even Years) – Course Syllabus
 Biol 475/575 Undergraduate/Graduate (3 Credit)
 SLN: (475) – 06763, (575) – 06764
 Time - Tuesday and Thursday 10:35 am-11:50 am
 Course Lectures in person and recorded on Canvas/Panopto and Discussion Sessions live in person and on WSU Zoom for all campuses (Hybrid Course)
 Room – CUE 418
 Course Director – Michael Skinner, Abelson Hall 507, 335-1524, skinner@wsu.edu
 Co-Instructor – Eric Nilsson, Abelson Hall 507, 225-1835, nilsson@wsu.edu
Learning Objective -
 Current literature based course on the Systems Biology of Reproduction. Learning Systems approaches to the biology of reproduction from a molecular to physiological level of understanding.

Schedule/Lecture Outline –

January	9 & 11	Week 1	Systems Biology Introduction
	16 & 18	Week 2	Molecular/ Cellular/ Reproduction Systems
	23 & 25	Week 3	Sex Determination Systems
Jan /Feb	30 & 1	Week 4	Male Reproductive Tract Development & Function
February	6 & 8	Week 5	Female Reproductive Tract Development & Function
	13 & 15	Week 6	Gonadal Developmental Systems Biology
	20 & 22	Week 7	Testis Systems Biology
	27 & 29	Week 8	Ovary Systems Biology
March	5 & 7	Week 9	Epigenetics and Transgenerational Gonadal Disease
	11 – 15	Week 10	Spring Break
	19 & 21	Week 11	Gametogenesis/ Stem Cells/ Cloning
	26 & 28	Week 12	Hypothalamus- Pituitary Development & Function
April	2 & 4	Week 13	Reproductive Endocrinology Systems
	9 & 11	Week 14	Fertilization & Implantation Systems
	16 & 18	Week 15	Fetal Development & Birth Systems
	23 & 25	Week 16	Assisted Reproduction/Contraception
Apr/May	30 & 2	Week 17	Exam or Grant Review

Spring 2024 – Systems Biology of Reproduction
 Lecture Outline – Gonadal Developmental Systems Biology
 Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday
 February 13, 2024
 Week 6

Gonadal Developmental Systems Biology

- Early Fetal Gonadal Development
 - Morphogenesis
 - Transcriptome
 - Meiotic Arrest
- Testis Gonadal Fetal Development
 - Knockout Models and Genes
 - Cellular Growth Regulation
- Ovary Gonadal Fetal Development
 - Developmental Timing and Morphology
 - Oocyte Nests and Primordial Follicle Assembly
 - Hormone Regulation of Assembly
 - Culture Models
- Disruption of Gonadal Development
 - Endocrine Disruptors

Required Reading

- Pepling and Burton (2018) Fetal/Gonadogenesis, Encyclopedia of Reproduction (Second Edition). Volume 2, Pages 47-51.
- Rotgers and Yao, (2018) Formation of the Testis Primordium, Encyclopedia of Reproduction (Second Edition). Volume 1, Pages 84-87.

Spring 2024 – Systems Biology of Reproduction
 Discussion Outline – Gonadal Developmental Systems Biology
 Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday
 February 15, 2024
 Week 6

Gonadal Developmental Systems Biology

Primary Papers:

1. Soto and Ross (2021) Reproduction 161:239-253
2. Real, et al. (2023) J Exp Zool B Mol Dev Evol. 340(3):231-244
3. Nilsson, et al. (2013) BMC Genomics 14:496

Discussion

Student 2: Reference #1 above


- What is the technical approach?
- What specific transcriptome observations were made?
- Why is the similarity in cow and human germline development?




Student 3: Reference# 2 above

- What are the technologies used and objectives?
- What environmental exposures were compared?
- What impacts on testis are observed in seasonal breeders?


Student 4: Reference #3 above

- What is the experimental and systems approach?
- What is a cluster analysis?
- What gene networks were identified for primordial follicle assembly










Paracrine regulation of steroidogenic cell development. (a) In the fetal mouse testis, Sertoli cells produce paracrine factors PDGF α and DHH that control the appearance and expansion of fetal Leydig cells in the interstitium after E12. Conversely, AMH from Sertoli cells negatively regulates the steroidogenic cell population. Fetal Leydig cells secrete activin A, which promotes proliferation of Sertoli cells via SMAD4 activation. Germ cells do not contribute to paracrine regulation of steroidogenic cells in the testis. (b) Unlike Leydig cells, which differentiate in the testis right after sex determination, the ovarian steroidogenic theca cells differentiate near birth and full steroidogenesis begins after birth. Also, oocytes participate in theca cell recruitment in the ovary through secreting GDF9, which induces expression of the hedgehog ligands DHH and IHH in the granulosa cells. DHH and IHH transform stromal progenitors into the theca cells that surround the follicles around birth in the mouse. Furthermore, granulosa cells promote theca cell recruitment by also producing PDGF α .




Gonadal supporting cells acquire sex-specific chromatin landscapes during mammalian sex determination.
 Garcia-Moreno SA, Futtner CR, Salamone IM, Gonen N, Lovell-Badge R, Maatouk DM.
 Dev Biol. 2019 Feb 15;446(2):168-179.






DMRT1 protects male gonadal cells from retinoid-dependent sexual transdifferentiation.

Minkina A, Matson CK, Lindeman RE, Ghyselinck NB, Bardwell VJ, Zarkower D. Dev Cell. 2014 Jun 9;29(5):511-20.




MYRF: A New Regulator of Cardiac and Early Gonadal Development-Insights from Single Cell RNA Sequencing Analysis

Catonga-Solis V, Fabbris-Scallett H, Ott F, et al. J Clin Med. 2022 Aug 18;11(16):4858.





UMAP plots of cell clusters in human (A) testes from embryonic stage to adults; and (B) ovaries from embryonic and foetal stages identified by single cell RNA sequencing. UMAP: uniform manifold approximation and projection.




Expression levels of MYRF in testes (A) and ovaries (B) at embryonic and fetal stages, showing that it is highly expressed in coelomic epithelium cells.

Testis Development


At the Crossroads of Fate-Somatic Cell Lineage Specification in the Fetal Gonad.
 Rotgers E, Jørgensen A, Yao HH.
 Endocr Rev. 2018 Oct 1;39(5):739-759.



SOX9 regulates microRNA miR-202-5p/3p expression during mouse testis differentiation.
 Wainwright EN, et al. Biol Reprod. (2013) 15;89(2):34.




Leydig cells: formation, function, and regulation.
 Zirkin BR, Papadopoulos V.
 Biol Reprod. 2018 Jul 1;99(1):101-111.





Fetal and adult periods of testosterone production. Fetal Leydig cells produce the high levels of testosterone that are required for the differentiation of the male genitalia and for brain masculinization. Testosterone production declines with the postnatal decline in numbers of the fetal Leydig cells, reaching a nadir early in the postpartum period. Thereafter, testosterone gradually increases to high levels with the development of the adult Leydig cells from stem cells of the neonatal testis. LH is not required either for the development of fetal Leydig cells or for their initial testosterone production. Later, however, the fetal Leydig cells express LH receptor and respond to LH stimulation.

Fetal Leydig cells dedifferentiate and serve as adult Leydig stem cells.
 Shima Y, Miyabayashi K, Sato T, Suyama M, Ohkawa Y, Doi M, Okamura H, Suzuki K.
 Development. 2018 Dec 5;145(23).



Transplantation of cell populations isolated from P10 testis. (A) Schematic of the strategy for cell transplantation and fate tracing. Cells isolated from P10 testes were transplanted into the interstitial space of the P28 nude mouse testis and analyzed at 8 weeks after transplantation. (B-C) Testis sections at P84 were immunostained for EGFP (green), NRSF1 (red) and laminin (white). The boxed area in B was enlarged, and EGFP and laminin signals are shown in B' and B'', respectively. The boxed area in C was enlarged, and EGFP and NRSF1 signals are shown in C' and C'', respectively. Arrows in B' and B'' indicate laminin⁺ PTMCs. Arrows in C' and C'' indicate NRSF1⁺ Leydig cells. (D-E) Testis sections at P84 were immunostained for EGFP (green) and HSD3B6 (red). The boxed area in D was enlarged, and EGFP and HSD3B6 signals are shown in D' and D'', respectively. The boxed area in E was enlarged, and EGFP and HSD3B6 signals are shown in E' and E'', respectively. Arrows in D' and D'' indicate HSD3B6⁺ ALCs, and arrows in E' and E'' indicate HSD3B6⁺ FLCs. (F) Percentages of the indicated cell populations in the recipient testis. We counted 6-70 cells in each section (three distinct sections from each sample; n=3 for population I and II, n=2 for population III). Scale bars: 50 μm.


Aristaless related homeobox gene, *Arx*, is implicated in mouse fetal Leydig cell differentiation possibly through expressing in the progenitor cells.
Miyabayashi K, et al. PLoS One. (2013) 28;8(6):e68050.




Reprogramming of Sertoli cells to fetal-like Leydig cells by *Wt1* ablation.
Zhang L, Chen M, Wen Q, Li Y, Wang Y, Wang Y, Qin Y, Cui X, Yang L, Huff V, Gao F.
Proc Natl Acad Sci U S A. 2015 Mar 31;112(13):4003-8.

Significance

Genetic control of the differentiation between Sertoli cells and granulosa cells has been reported previously. However, the relationship between Sertoli cells and Leydig cells in the testis has not yet been definitively determined. In the present study, we demonstrate for the first time, to our knowledge, that these two cell types can be mutually reprogrammed and that Wilms' Tumor Gene 1 (*Wt1*) plays a critical role in this process. This study provides a novel concept for cell fate determination in testis development that will improve our understanding of the regulatory mechanisms of gonad development.




Transcription factors ER71/ETV2 and SOX9 participate in a positive feedback loop in fetal and adult mouse testis.
 DiTacchio L, et al. J Biol Chem. (2012) 6;287(28):23657-66.





Role of transforming growth factor-alpha and the epidermal growth factor receptor in embryonic rat testis development.


Levine E, Cupp AS, Miyashiro L, Skinner MK.
 Biol Reprod. 2000 Mar;62(3):477-90.





E13 testis + mesonephros organ cultures after 1 day (A) and after 3 days (B) in culture. The upper structure (T) is the testis attached to the lower, dark-colored mesonephros (M). Representative of more than 10 different experiments. The diameter of the organ was approximately 1000 µm. E14 testis organ cultures treated with control nonimmune IgG (C) or with anti-TGF-α IgG antibody S574 (D) after 4 days in culture. Representative cords are marked by arrows placed on the lighter-colored structures and are shown in C and D. This is a microdissected testis devoid of mesonephros with some manipulation in comparison to that shown in A and B. Representative of 10 experiments



Labeling indices of the percentage of PCNA-positive cells for each cell type at various stages of development. Labeling indices were calculated by counting stained and unstained cells in random regions of replicate slides from at least 3 different sections in 3 different experiments. Data are presented as the mean ± SEM.







Labeling index for the percentage of each cell type positive for PCNA on e14 testis organ cultures treated with control nonimmune or anti-TGF- α antibody for 4 days. Labeling indices were calculated by counting stained and unstained cells in random regions of at least 3 different sections from 3 different experiments. ** Statistically significant difference from control ($P < 0.001$); * statistically significant difference from control ($P < 0.02$) as determined by a t-test




A) E18 wild-type (WT) and B) e18 knockout (KO) mice testes. JB4 sections were stained with toluidine blue. $\times 200$ (published at 80%). Data are representative of more than 10 sections from different testes

Ovary Development




The art of oocyte meiotic arrest regulation.
Pan B, Li J.
Reprod Biol Endocrinol. 2019 Jan 5;17(1):8.






Schematic model depicting the proposed signaling pathway in LH/ICG-induced resumption of meiosis. LH surge results in the dephosphorylation and deactivation of NPR2 via the phosphoprotein of phosphatase (PPP)-family in the cumulus granulosa cell [96]. NPR2 production and secretion in the mural granulosa cell. The LH/LHR-induced activation of cAMP/PKA increases the following compounds in the mural granulosa cell: phosphorylation of the cGMP-PDE5, EGF and EGF-like factors production and secretion, and ER expression [103]. The rapid phosphorylation of the cGMP-PDE5 is responsible for converting cGMP into 5'GMP, causes the outward diffusion of intracytole cGMP, leads to the release of cAMP-PDE activity within the oocyte, in turn, the activated cAMP-PDE catalyses the cAMP into AMP, ultimately results in meiotic resumption [99]. The production of EGF and EGF-like factors activates the EGFR signaling elevates calcium of cumulus granulosa cells further to inactivate NPR2 [108]. Furthermore, the decrease of ER serves as an augment step [103]. cGMP-PDE5: cGMP phosphodiesterase 5; EGF-like factors: amphiregulin (AREG), epiregulin (EREG), and β -collagen (BTC)

Retinoic Acid and Germ Cell Development in the Ovary and Testis.
Endo T, Makedis MM, Nicholis PK, Page DC, de Rooij DG.
Biomolecules. 2019 Nov 24;9(12).



Follicular Assembly and Primordial to Primary Follicle Transition



Number of Oocytes During Stages of Early Folliculogenesis

	Proliferation	Assembly	Puberty
Rodent	75,000	27,000	10,000
Primate	6,800,000	1,000,000	700,000


Only 500 human follicles ovulate.

Perinatal Steroids and Follicular Assembly


	Time of Initiation	[Steroid]	Time of Completion	[Steroid]
Rodent	Birth	5×10^{-7}	Day 4 Post-Natal	2×10^{-8}
Primate	Mid Gestation	4×10^{-7}	Birth	NA

Steroids might be endocrine factor that coordinates follicular assembly.


Progesterone regulation of primordial follicle assembly in bovine fetal ovaries.
Nilsson EE, Skinner MK. Mol Cell Endocrinol. (2009) 10;313(1-2):9-16.




Correlation of progesterone concentration in fetal bovine ovaries and the percentage (%) of oocytes assembled into primordial follicles as a function of estimated gestational age (days). Follicle assembly is initiated as ovarian progesterone concentration declines.



Ovarian Floating Filter Culture System




Culture & Treatment
(1 or 2 weeks)






Analysis


Follicular Kinetics *in vivo/in vitro*.







Oocyte Apoptosis in Progesterone Treated and Control Neonatal Ovary.




Effect of Steroids on Neonatal Ovary Culture







Progesterone Regulated Follicle Assembly



Progesterone Inhibited Oocyte Apoptosis



Model for Primordial Follicle Assembly



Development 124, 4939-4947 (1997)
 Printed in Great Britain © The Company of Biologists Limited 1997
 0950262

4939

FIG α , a germ cell specific transcription factor involved in the coordinate expression of the zona pellucida genes

Li-fang Liang, Selma M. Soyal and Jurrien Dean*
 Laboratory of Cellular and Developmental Biology, NIDDK, National Institutes of Health, Bethesda, MD 20892, USA
 *Author for correspondence (e-mail: juriend@helix.nih.gov)

SUMMARY

The mouse zona pellucida is composed of three glycoproteins, ZP1, ZP2 and ZP3, encoded by single-copy genes whose expression is temporally and spatially restricted to oocytes. All three proteins are required for the formation of the extracellular zona matrix and female mice with a single disrupted zona gene lack a zona and are infertile. An E-box (CANNTG), located approximately 200 bp upstream of the transcription start sites of *Zp1*, *Zp2* and *Zp3*, forms a protein-DNA complex present in oocytes and, to a much lesser extent, in testes. It has been previously shown that the integrity of this E-box in *Zp2* and *Zp3* promoters is required for expression of luciferase reporter genes

microinjected into growing oocytes. The presence of the ubiquitous transcription factor E12 in the complex was used to identify a novel basic helix-loop-helix protein, FIG α (Factor In the Germline α) whose expression was limited to oocytes within the ovary. The ability of FIG α to transactivate reporter genes coupled to each of the three mouse zona promoters in heterologous 10T1 embryonic fibroblasts suggests a role in coordinating the expression of the three zona pellucida genes during oogenesis.

Key words: FIG α , basic helix-loop-helix transcription factor, zona pellucida, oocyte-specific gene expression, mouse







Table 1 Summary of analysis and results following e12.5 female genital ridge transplantation.

Days of Transplantation	Analysis ^a	Re-Agg Gonad	Intact Gonad	In Vivo ^b Control
3	Meiosis	Zygotene-pachytene of meiosis I	Zygotene-pachytene of meiosis I	Zygotene-pachytene of meiosis I [18]
5	Ovarian Cord Formation	Ovarian cords do not form and ovary disorganized	Ovarian cords form with distinct organization of oocyte and granulosa cell clusters	Ovarian cords form with distinct organization of oocyte and granulosa cell clusters [1]
7	Oocyte Survival	Oocyte numbers significantly reduced compared to intact gonad	Oocyte numbers reduced compared to day 5	Oocyte numbers decline due to programmed breakdown [10]
12-21	Oocyte Development in Follicles	Oocytes do not survive to day 12 and follicles not observed	Oocytes survive, form follicles, and mature	Oocytes survive, form follicles, and mature [2]

^a For all experiments, n = 3 transplants with 4 embryos or 8 gonads per transplant
^b In vivo control refers to e15.5, e17.5, e19.5, and P4 - P14 stages

Model for Primordial Follicle Assembly

Oocyte "Nest" → Programmed Oocyte Death → Primordial Follicles




Objective: Identify the genes involved in follicle assembly.

Experimental approach:


- Take ovaries from 0-day old rats.
- Culture for 24h with one of several growth factors.
- Extract RNA.
- Affymetrix microarrays.

OVARIAN FLOATING FILTER CULTURE SYSTEM



Culture & Treatment

2 days



Morphological Analysis


1 day

Collect RNA
Microarrays

Treatments


AMH	Anti-Mullerian hormone
CTGF	Connective tissue growth factor
E2	Estradiol
FGF2	Fibroblast growth factor 2
INHBa	Activin A (inhibin alpha subunit)
P4	Progesterone
TNFa	Tumor necrosis factor alpha
Cont	Control untreated

RMA-GC pre-processing of all 28 chips




Conclusion:
we have one outlier (P4 sample)
To do : omit this sample and
pre-process again

RMA-GC pre-processing of 27 chips (minus an outlier)



Follicle Assembly Differentially Expressed Genes




Analysis Approach

1. Microarray analysis by treatment.
2. Coexpression analysis (modules).
3. Automated literature network analysis.

Number of genes and KEGG pathways overlapped between signature lists

	# PW	AMH	CTGF	E2	FGF2	INHBa	P4	TNFa	3+ Total
# genes	73	9	79	115	83	85	63	138	
AMH	158	4	45	63	43	41	29	73	
CTGF	50	2	7	8	7	7	5	9	
E2	120	10	0	69	49	49	38	79	
FGF2	303	13	4	9	71	70	50	115	
INHBa	287	11	1	4	14	52	40	83	
P4	167	5	3	8	11	14	43	85	
TNFa	116	6	4	4	7	9	3	63	
Total	1081								

P0ovaries, 7treatments, 1081 gene list, KEGG from Entrez ID, gene overlaps from
Number of overlaps with only KEGG 3+ start list




KEGG ID	Pathway name	total # genes	AMH	CTGF	E2	FGF2	INHBa	P4	TNFa
			158	50	120	303	287	167	116
mo01100	Metabolic pathways	59	9	6	11	23	9	10	
mo05200	Pathways in cancer	19	3	3	7	3	3	2	
mo04740	Olfactory transduction	19	2	3	1	2	11	1	
mo04010	MAPK signaling pathway	15		2	7	2	3	1	
mo04510	Focal adhesion	15	1	4	5	4	3		
mo04062	Chemokine signaling pathway	15	3	2	6	4	1	1	
mo04144	Endocytosis	15		5	3	3	2	2	
mo00980	Metabolism of xenobiotics by cytochrome P450	14		1	1	3	3	6	
mo04145	Phagosome	13	1	4	3	4	2	3	
mo04020	Calcium signaling pathway	12	5	3	4	1	1	1	
mo04514	Cell adhesion molecules	12	1	1	4	2	1	3	
mo04360	Axon guidance	11	1		6	3	3	1	
mo04810	Regulation of actin cytoskeleton	11	2	3	4	2	2		
mo00480	Glutathione metabolism	11		1	1	3	3	3	1
mo00230	Purine metabolism	11	3	3	3	2	1		
mo03010	Ribosome	11				11			
mo04060	Cytokine-cytokine receptor interaction	11			6	3	2		
mo04080	Neuroactive ligand-receptor interaction	10	2	1	6	3	1		
mo04976	Bile secretion	10	2	2	4	1	1	1	
mo04380	Osteoclast differentiation	9			5	3	1	1	
mo04640	Hematopoietic cell lineage	9	3	1	2	2	1	1	1
mo04110	Cell cycle	8	1		5	1	1		
mo04512	ECM-receptor interaction	7	1	2	2	2	1		
mo04612	Antigen processing and presentation	7			1	3	3		
mo04972	Pancreatic secretion	7		1	3	2	1		
mo04350	TGF-beta signaling pathway	7	1	1	3	2			

Analysis Approach

1. Microarray analysis by treatment.
2. Coexpression analysis (modules).
3. Automated literature network analysis.

Co-expression heat map







Module	Database	Function	# module genes	p-value
turquoise	KEGG pathway	Ribosome	10	7.52E-08
turquoise	GO: Cell Component	cytosolic ribosome	7	0.0000024
turquoise	GO: Biological Process	response to virus	6	0.00013
turquoise	GO: Biological Process	neural tube closure	4	0.00041
turquoise	GO: Biological Process	negative regulation of binding	5	0.00045
turquoise	Panther Biological Process	Protein biosynthesis	15	0.00071
turquoise	KEGG pathway	Glutathione metabolism	5	0.0011
turquoise	KEGG pathway	TGF-beta signaling pathway	3	0.09
blue	GO: Biological Process	response to oxidative stress	12	0.000013
blue	GO: Biological Process	regulation of anatomical structure morphogene	13	0.000022
blue	Panther Biological Process	Mesoderm development	18	0.00019
blue	Panther Biological Process	Angiogenesis	6	0.00022
blue	GO: Biological Process	response to carbohydrate stimulus	6	0.00036
blue	GO: Biological Process	regulation of axonogenesis	6	0.00005
blue	GO: Cell Component	membrane raft	9	0.00078
blue	GO: Biological Process	negative chemotaxis	3	0.0001
blue	KEGG pathway	Axon guidance	6	0.0037
blue	KEGG pathway	Glutathione metabolism	3	0.031
blue	KEGG pathway	Focal adhesion	6	0.032
blue	KEGG pathway	Fc gamma R-mediated phagocytosis	4	0.036
blue	KEGG pathway	Calcium signaling pathway	5	0.067
blue	KEGG pathway	Cytokine-cytokine receptor interaction	5	0.074
brown	GO: Cell Component	germ cell nucleus	3	0.0007
brown	GO: Biological Process	male meiosis	3	0.0015
brown	GO: Biological Process	neurite regeneration	3	0.0031
brown	GO: Cell Component	condensed nuclear chromosome	3	0.0041
brown	KEGG pathway	Cell cycle	4	0.026
brown	KEGG pathway	Olfactory transduction	12	0.065

Number of genes overlapped between signatures (treatments) and modules										
		turquoise	blue	brown	yellow	green	red	black	pink	magenta
	# genes	240	209	176	65	45	43	27	26	20
AMH	158	14	7	28	19	25	0	0	7	3
CTGF	50	0	4	8	1	1	5	2	3	4
E2	120	1	6	3	20	8	14	15	9	4
FGF2	303	4	169	44	28	9	1	0	2	4
INHba	287	184	24	26	5	2	15	0	14	1
P4	167	16	5	58	11	0	8	6	7	4
TNFa	116	36	10	19	6	4	2	4	0	6


Number of overlaps with all probeset IDs
Bold numbers heavily represented by module.
Underlined numbers heavily represented by signature (treatment).

Analysis Approach



1. Microarray analysis by treatment.
2. Coexpression analysis (modules).
3. Automated literature network analysis.




Follicular Assembly and Primordial to Primary Follicle Transition







Establishment and depletion of the ovarian reserve: physiology and impact of environmental chemicals.
 Ge W, Li L, Dyer PW, De Felici M, Shen W.
 Cell Mol Life Sci. 2019 May;76(9):1729-1746.




Fetal Estrogens are not Involved in Sex Determination But Critical for Early Ovarian Differentiation in Rabbits.
 Jolivet G, Daniel-Carlier N, Harscoët E, et al.
 Endocrinology. 2022 Jan 1;163(1):bqab210.





In utero exposure to diisononyl phthalate caused testicular dysgenesis of rat fetal testis.


Li L, Bu T, Su H, Chen Z, Liang Y, et al. *Toxicol Lett.* 2015 Jan 22;232(2):466-74.



Occurrence of multinucleated gonocytes (MNGs) in representative photomicrographs of testis section and percentage of seminiferous cords exhibiting MNGs. (A) Control group and (B) DINP 1000 mg/kg group. Sections were stained with hematoxylin-eosin stains. Arrow indicates MNG. Scale bar = 50 μ m. (C) Percentage of seminiferous cords exhibiting MNGs on GD21.5 (n = 6); Mean \pm SEM. Significant differences at *P < 0.05, **P < 0.01 and ***P < 0.001, respectively, were shown, when compared to control.

Di (2-ethylhexyl) phthalate exposure during pregnancy disturbs temporal sex determination regulation in mice offspring.


Wang Y, Liu W, Yang Q, Yu M, Zhang Z. *Toxicology.* 2015 Oct 2;336:10-6.



Expression levels and patterns of Sry in male genital ridges in offspring from control and embryonal DEHP treatment group during 11.5–12.5 dpc. A: Expression pattern of Sry examined by whole-mount in situ hybridization (WISH) in male genital ridges. Antisense probe is active versus sense probe (data not shown). Arrows indicate the sex cord in gonad. Bars: 100 μ m. B: Relative expression levels of Sry normalized to the β -actin in male genital ridges analyzed by RT-qPCR. Values significantly different from control at p < 0.05 are indicated by asterisks. Values are litter mean \pm SE (n = 4–6 for control, 350 and 700 mg/kg/d groups).


Foetal-neonatal exposure of Di (2-ethylhexyl) phthalate disrupts ovarian development in mice by inducing autophagy.

Zhang Y, Mu X, Gao R, Geng Y, Liu X, Chen X, Wang Y, Ding Y, Wang Y, He J. *J Hazard Mater.* 2018 Sep 15;358:101-112.



Cellular and molecular effect of MEHP Involving LXR α in human fetal testis and ovary.

Muczynski V, et al. *PLoS One.* (2012) 7(10):e48266.








Table 1 Proteins found to be significantly down- or up-regulated following both 10 and 100 mg/kg DEHP exposures.

Spot	Symbol	Accession numbers	Protein name	Protein MW (kDa)	Percent coverage	Dose				Molecular function
						0	10	100	500	
2	SITP1	Q12CL9	Stress induced phosphoprotein 1	63	66	100	58*	52	Protein-protein binding	
8	SYG	Q58G24	Glycyl t-RNA synthetase	72	15	100	56	66	Protein biosynthesis	
22	LMAN1	Q62902	Leucin mannosidase binding 1	55	30	100	61	71	Protein transport	
23	PEXA3	P11598	Protein disulfide-isomerase A3	57	60	100	67	71	Cellular redox homeostasis	
47	ENO1	P04764	Alpha enolase	47	29	100	60	60	Glycolysis/plasminogen activation	
55	ENOI	P04764	Alpha enolase	47	37	100	78	73	Glycolysis/plasminogen activation	
84	HSPM1	P63018	Heat shock protein 71 kDa	71	38	100	143	162	Chaperone/transcriptional repressor	
108	ACTG	P63259	Actin, cytoplasmic 2	42	33	100	40	37	Responder to calcium/ATP binding	
111	HSP90B1	Q668D0	Heat shock protein 90 kDa, subunit 1	90	7	100	53	59	Chaperone in the endoplasmic reticulum	
112	HSP90B1	Q668D0	Heat shock protein 90 kDa, subunit 1	90	7	100	51	50	Chaperone in the endoplasmic reticulum	
127	TCP1	P28480	T complex protein 1 subunit alpha	60	40	100	62	60	Chaperone involved in protein folding	
130	TRAP1	Q530Z0	Heat shock protein 75 kDa, Mitochondrial	74	26	100	68	58	Chaperone involved in protein folding	
143	SRM	Q99A85	Spermatidie synthase	34	18	100	132	140	Spermatidie biosynthesis	
174	ATP6V1-B2	P62815	V type proton ATPase subunit B	57	12	100	49	47	Hydrogen ion transport/ATPase activity	
191	RPLP0	P19945	60S acidic ribosomal protein P0	34	34	100	67	63	Translational elongation	
196	PDIA6	Q63081	Protein disulfide-isomerase A6	46	33	100	78	70	Chaperone/platelet aggregation	
202	CCT5	Q68FQ0	T complex protein 1 subunit epsilon	59	23	100	71	76	Chaperone involved in protein folding	
242	GSTM4	P08009	Glutathione S-transferase	26	6	100	72	73	Conjugation of reduced glutathione	
319	TUBM	P49411	Elongation factor Tu, Mitochondrial	45	67	100	78	74	Translation of proteins	
333	TRAP1	Q530Z0	Heat protein 75 kDa, Mitochondrial	74	27	100	62	67	Chaperone/ATPase activity	
338	YWHAE	P62260	14-3-3 epsilon	32	32	100	142	177	Signal transduction via protein binding	
349	CCT2	Q530M9	T complex protein 1 subunit beta	57	57	100	80	75	Chaperone involved in protein folding	
365	DPI12	P47942	Dihydrocyprimidinase-related protein 2	62	28	100	70	76	Differentiation/cell migration	
366	RPL13A	Q58R80	WD repeat containing protein 1	66	26	100	134	152	Disassembly of actin	
380	VIM	P31000	Vimentin	54	24	100	59	74	Intermediate filament of cytoskeleton	
381	RPSA	P38983	40 S ribosomal protein SA	33	33	100	73	67	Cell surface receptor for laminin/morphogenesis	

Statistical results represent the analysis of 161 per treatment group (n=16); one testis from four GD19 males pooled and extracted/filtered.
 *The background-corrected, integrated optical density of each spot was averaged within group and expressed as percent of control. The symbols of proteins that were found to be significantly (P<0.05) correlated with and/or predictive of treatment or measured endpoints (bold and underlined).

Human exposure to endocrine disrupting chemicals: effects on the male and female reproductive systems.
Sifakis S, Androutopoulos VP, Tsatsakis AM, Spandidos DA.
Environ Toxicol Pharmacol. 2017 Apr;51:56-70.






Table 1
A summary of the studies that have assessed the associations between the exposure of human populations to pesticides and EDC agents and the incidence of disorders affecting the female reproductive system. (C): case-control study; (CC): Cohort study.

Author, Year	Chemical	N	Design	Findings	Variables	Study Type
Sundaram et al., 2011	BPA	171/150 PCOS, 71 Normal	Case-control	Higher BPA levels in PCOS, association of BPA with androgens	Sexual parameters, BMI	CC
Chen et al., 2013	BPA, phthalates	61 women	Case-control	Higher BPA and oestrogen levels in infertile group	Uterine BPA, E2 levels in serum, endogenous and metabolic responses	CC
Souza et al., 2011	BPA	209 women on fertility treatment	Case-control	Higher BPA levels associated with lower AFC	Uterine BPA, Antral Follicle Count (AFC), FSH	CC
Shih-Chieh et al., 2012a,b	BPA	147 women with IVF cycles	Case-control	Negative association between urinary BPA and serum peak E2 and oocyte yield	Urinary BPA, oocyte maturation, fertilization, embryo quality	CC
Shih-Chieh et al., 2012a,b	BPA	137 women	Case-control	Positive association between urinary BPA and implantation failure	Urinary BPA, β-HCG, implantation failure	CC
Hsieh-Lee et al., 2010	BPA	84 women	Case-control	Urinary BPA is inversely correlated with serum peak E2	Urinary BPA, E2 levels, number of oocyte	CC
Shaw et al., 2011	BPA	44 women with IVF	Case-control	Serum BPA is associated with reduced E2 levels	Serum BPA, E2 levels, number of oocytes	CC
Agarwal et al., 2011	BPA	58 infertile females, 37 male partners	Case-control	Inverse association between BPA and fertilization	Serum BPA, fertilization	CC
Yang et al., 2008	BPA	162 women with breast cancer, 152 controls	Case-control	No association with BPA and breast cancer	Blood BPA, incidence of breast cancer	CC
Artemeva et al., 2008	BPA, BPA, Benzyl phthalate, nonylphenol	201 women with breast cancer, 753 controls	Case-control	Association with PCBs, 4-nonylphenol and breast cancer	Levels of EDC and pesticides and incidence of breast cancer	CC
Whitford et al., 2012	BPA, phthalates	288 mother-infant pairs (72 cases, 216 controls)	Case-control	Positive/Negative association of bisphenol/AES, DCP with male birth weight, respectively	Levels of EDC and pesticides, birth weight, head circumference, birth length	CC
Ingber, Opatovska et al., 2005	BPA	77 women (45 cases with miscarriage, 32 controls)	Case-control	Serum BPA is associated with increased miscarriage	Serum BPA, ADA, NK cells, cytokines, progesterone, TNF, Fcγ	CC
Such-Lau et al., 2013	BPA, phthalates	625 women	Cohort	mCNP, mDEHP, mDHP, mBP, mBP and mCNP were associated with endometriosis	BPA, 1,4-phthalate metabolites, age, BMI, creatinine	CC
Shaw et al., 2013	DAF, DE, DM	359 mothers and children	Cohort	Early life exposure (1-7 years) to DAF, DE, DM is associated with respiratory symptoms in childhood	Urinary DAF, DE, DM and metabolites, respiratory symptoms in childhood	CC
Makris et al., 2012	HCB, DDT, DDE	730 women with IVF	Case-control	Serum HCB is associated with failed implantation	Serum HCB, DDT, DDE and metabolites, fertilization	CC

Single-cell transcriptome dissection of the toxic impact of Di (2-ethylhexyl) phthalate on primordial follicle assembly.
Wang J-J, Tian Y, Li M-H, et al.
Theranostics. 2021 Mar 5;11(10):4992-5009.



In an Ovine Model of Polycystic Ovary Syndrome (PCOS) Prenatal Androgens Suppress Female Fetal Renal Gluconeogenesis.
Connolly F, Rae MT, Späth K, Boswell L, McNeilly AS, Duncan WC.
PLoS One. 2015 Jul 6;10(7):e0132113.




Developmental programming: gestational testosterone treatment alters fetal ovarian gene expression.
Luense LJ, et al. (2011) 152(12):4974-83.

TABLE 1. miRNA differentially expressed in control vs. T- or T plus flutamide-treated fetal ovine ovaries

miRNA	C vs. T FC	C vs. T+FC	Functional analysis literature based ^a	Predicted targets (TargetsCan)
miR-497	3.83	3.71	Type 2 diabetes rat [miR-15b/497 family (1)]	PAPPA, IR, GHR, IGF2R, IRS2, IGF1R, turin
miR-29a	2.98		Diabetes/insulin signaling (1-5), FSH regulated (6)	IGF1, INSG1, leptin
miR-192	2.19		Diabetic neuropathy (4, 7, 8)	IGF1
miR-24-2*	2.12		Diabetes (9), insulin signaling (10), bovine ovary (11)	INSIG1, PPARa, Furin, IGF2BP2
miR-15b	2.06	2.93	Type 2 diabetes rat FSH (1), regulated in ovary (6)	PAPPA, IR, GHR, IGF2R, IRS2, IGF1R, furin
miR-101	1.90		AR regulated (12)	PGRMC2, PPARa
miR-212	1.87		LH regulated in ovary (13)	
miR-451	1.85		Sex dependent in liver (14)	
miR-186	1.82		Type 1 diabetes (15)	INSM1, LEPR, IGF1, IGF1R
miR-672	1.82		Mouse ovary (16)	IGF1R
miR-7	1.67		Insulin signaling (17, 18)	IRS2, IRS1, IGF1R, PAPPA
miR-30b-5p	1.58		Estrogen regulated (19)	LEPR, IGF2R, INSG2, IRS1, IRS2, LDLR, IGF1R, IGF1
miR-22*	1.52		Fetal ovine gonad (20), androgen regulated (21), represses ESR1 (22)	ESR1, GHRHR, PTCG1, IGF2BP1, furin
miR-378	-4.13		Lipid/fatty acid metabolism (23), regulates estrogen production (24)	
miR-760	-2.80		Estrogen regulated (25)	IR
miR-10a	-2.76	-3.36	Androgen regulated (26), ovary (27)	PPARa
miR-182	-2.71		Insulin signaling (10), ovary (27)	INSIG1, IGF2BP1
miR-129*	-2.22		Diabetes (28)	IGF1, GHR, ESR1, INSG2
miR-132	-1.82		LH regulated in ovary (13)	
miR-223	-1.69		Diabetes (9)	IGF1R
miR-363	4.50		Sex differentiation (29)	PTGER4, IRS2, INSG1
miR-20b	3.24		Diabetes (9), ESR1 regulated (30)	PPARa, LDLR, IGF2BP1, ADIPOR2, PPARd
miR-330	1.71		Fetal programming (31)	IGF2BP1
miR-29c	1.64		Diabetic neuropathy (32), diabetes (2)	IGF1, INSG1, leptin
miR-29b	1.62		Diabetes (2, 9), sex dependent in liver (14)	IGF1, INSG1, leptin
miR-191*	1.56		Diabetes (9)	
miR-101a*	1.53		AR regulated (12)	PGRMC2, PPARa
miR-105	1.52		Human ovary (27)	
miR-133a	-1.85		Insulin signaling (33, 34)	IR, IGF1R



C, Controls; IR, estrogen receptor; FC, Fold change; T, testosterone + flutamide. References for table are included in the Supplemental Online Materials.

Genistein impairs early testosterone production in fetal mouse testis via estrogen receptor alpha.
Lehraiki A, et al. Toxicol In Vitro. (2011) 25(8):1542-7.



Disruption of rat testis development following combined in utero exposure to the phytoestrogen genistein and antiandrogenic plasticizer di-(2-ethylhexyl) phthalate.

Jones S, Boisvert A, Duong TB, Francois S, Thrane P, Culty M. Biol Reprod. 2014 Sep;91(3):64.



"Systems Biology of Reproduction"

Spring 2024 (Even Years) – Course Syllabus

Biol 475/575 Undergraduate/Graduate (3 Credit)

SLN: (475) – 06763, (575) – 06764

Time - Tuesday and Thursday 10:35 am-11:50 am

Course Lectures in person and recorded on Canvas/Panopto and Discussion Sessions live in person and on WSU Zoom for all campuses (Hybrid Course)

Room – CUE 418

Course Director – Michael Skinner, Abelson Hall 507, 335-1524, skinner@wsu.edu

Co-Instructor – Eric Nilsson, Abelson Hall 507, 225-1835, nilsson@wsu.edu

Learning Objective -

Current literature based course on the Systems Biology of Reproduction. Learning Systems approaches to the biology of reproduction from a molecular to physiological level of understanding.

Schedule/Lecture Outline –

January	9 & 11	Week 1	Systems Biology Introduction
	16 & 18	Week 2	Molecular/ Cellular/ Reproduction Systems
	23 & 25	Week 3	Sex Determination Systems
Jan /Feb	30 & 1	Week 4	Male Reproductive Tract Development & Function
February	6 & 8	Week 5	Female Reproductive Tract Development & Function
	13 & 15	Week 6	Gonadal Developmental Systems Biology
	20 & 22	Week 7	Testis Systems Biology
	27 & 29	Week 8	Ovary Systems Biology
March	5 & 7	Week 9	Epigenetics and Transgenerational Gonadal Disease
	11 – 15	Week 10	Spring Break
	19 & 21	Week 11	Gametogenesis/ Stem Cells/ Cloning
	26 & 28	Week 12	Hypothalamus- Pituitary Development & Function
April	2 & 4	Week 13	Reproductive Endocrinology Systems
	9 & 11	Week 14	Fertilization & Implantation Systems
	16 & 18	Week 15	Fetal Development & Birth Systems
	23 & 25	Week 16	Assisted Reproduction/Contraception
Apr/May	30 & 2	Week 17	Exam or Grant Review