

**Spring 2026 – Systems Biology of Reproduction**  
**Lecture Outline – Ovarian Systems Biology**  
**Eric Nilsson – Biol 475/575**  
**10:35-11:50 am, Tuesday & Thursday**  
**March 3, 2026**  
**Week 8**

## **Ovarian Systems Biology**

### Cell Biology of the Ovary

- Cell types/organization
- Developmental stages (Folliculogenesis)
- Atresia/apoptosis
- Oogenesis

### Regulation of Folliculogenesis

- Growth properties of ovarian follicles
- Local production and action of growth factors
- Growth regulations during development
- Primordial follicle transition

### Endocrine Regulation of Tissue Function

- Gonadotropin actions (Pituitary/Gonadal Axis)
- Steroid production and action
- Two cell theory modifications
- Hormone actions during development

### Cell-Cell Interactions

- Categorization of different cell-cell interactions in the ovary
- Growth factor regulation follicle development
- Oogenesis and systems biology

## **Required Reading**

Bahr JM. (2018) Ovary, Overview. in: Encyclopedia of Reproduction 2nd Edition, Ed: MK Skinner. Elsevier. Vol 2: 3-7.

## **REFERENCES**

Hammami B, Mostafavi FS, Akbari A, Mousavi SR, Kazemi M. Evaluation of the expression profile of mRNAs and lncRNAs in cumulus cells associated with polycystic ovary syndrome and pregnancy. Iran J Basic Med Sci. 2023;26(10):1144-1154.

- Ghafouri F, Sadeghi M, Bahrami A, et al. Construction of a circRNA- lincRNA-lncRNA-miRNA-mRNA ceRNA regulatory network identifies genes and pathways linked to goat fertility. *Front Genet.* 2023 Jul 21;14:1195480.
- Hossain Md A, Ashik SAA, Mahin MR, et al. Systems biology and in silico-based analysis of PCOS revealed the risk of metabolic disorders. *Heliyon.* 2022 Dec 22;8(12):e12480.
- Das A, Destouni A. Novel insights into reproductive ageing and menopause from genomics. *Hum Reprod.* 2023 Feb 1;38(2):195-203.
- Machlin JH, Shikanov A. Single-cell RNA-sequencing of retrieved human oocytes and eggs in clinical practice and for human ovarian cell atlasing. *Mol Reprod Dev.* 2022 Dec;89(12):597-607.
- Bongrani A, Plotton I, Mellouk N. High androgen concentrations in follicular fluid of polycystic ovary syndrome women. *Reprod Biol Endocrinol.* 2022 Jun 14;20(1):88.
- Pierson Smela MD, Kramme CC, Fortuna PRJ, et al. Directed differentiation of human iPSCs to functional ovarian granulosa-like cells via transcription factor overexpression. *Elife.* 2023 Feb 21;12:e83291.
- Telfer EE, Grosbois J, Odey YL, Rosario R, Anderson RA. Making a good egg: human oocyte health, aging, and in vitro development. *Physiol Rev.* 2023 Oct 1;103(4):2623-2677
- Zhang X, Cao Q, Rajachandran S, Grow EJ, Evans M, Chen H. Dissecting mammalian reproduction with spatial transcriptomics. *Hum Reprod Update.* 2023 Nov 2;29(6):794-810. doi: 10.1093/humupd/dmad017.
- Guarnieri G, Iervolino M, Cavallone S, Unfer V, Vianello A. The "Asthma-Polycystic Ovary Overlap Syndrome" and the Therapeutic Role of Myo-Inositol. *Int J Mol Sci.* 2023 Apr 9;24(8):6959
- Xiao C, Wang J, Zhang C. Synthesis, Regulatory Factors, and Signaling Pathways of Estrogen in the Ovary. *Reprod Sci.* 2023 Feb;30(2):350-360. doi: 10.1007/s43032-022-00932-z.
- Reiter RJ, Sharma R, Romero A, Manucha W, Tan DX, Zuccari DAPC, Chuffa LGA. Aging-Related Ovarian Failure and Infertility: Melatonin to the Rescue. *Antioxidants (Basel).* 2023 Mar 11;12(3):695.
- Bizzarri M, Monti N, Piombarolo A, Angeloni A, Verna R. Myo-Inositol and D-Chiro-Inositol as Modulators of Ovary Steroidogenesis: A Narrative Review. *Nutrients.* 2023 Apr 13;15(8):1875.
- Fedeli V, Catizone A, Querqui A, Unfer V, Bizzarri M. The Role of Inositols in the Hyperandrogenic Phenotypes of PCOS: A Re-Reading of Larner's Results. *Int J Mol Sci.* 2023 Mar 27;24(7):6296.
- Ben Yaakov T, Wasserman T, Akin E, Savir Y. Single-cell analysis of the aged ovarian immune system reveals a shift towards adaptive immunity and attenuated cell function. *Elife.* 2023 Apr 25;12:e74915.
- Beresniak A, Russo M, Forte G, Laganà AS, Oliva MM, Aragona C, Chiantera V, Unfer V. A Markov-model simulation of IVF programs for PCOS patients indicates that coupling myo-Inositol with rFSH is cost-effective for the Italian Health System. *Sci Rep.* 2023 Oct 18;13(1):17789.
- Kaboli Kafshgiri S, Farkhondeh T, Miri-Moghaddam E. Glyphosate effects on the female reproductive systems: a systematic review. *Rev Environ Health.* 2021 Jul 15;37(4):487-500.
- Oliver E, Alves-Lopes JP, Harteveld F, et al. Self-organising human gonads generated by a Matrigel-based gradient system. *BMC Biol.* 2021 Sep 23;19(1):212.
- Mao Z, Li T, Zhao H, Wang X, Kang Y, Kang Y. Methylome and transcriptome profiling revealed epigenetic silencing of LPCAT1 and PCYT1A associated with lipidome alterations in polycystic ovary syndrome. *J Cell Physiol.* 2021 Sep;236(9):6362-6375.
- Greaney J, Subramanian GN, Ye Y, Homer H. Isolation and in vitro Culture of Mouse Oocytes. *Bio Protoc.* 2021 Aug 5;11(15):e4104.

- Kaboli Kafshgiri SK, Farkhondeh T, Miri-Moghaddam E. Glyphosate effects on the female reproductive systems: a systematic review. *Rev Environ Health*. 2021 Jul 15.
- Bizzarri M, Logoteta P, Monastra G, Laganà AS. An innovative approach to polycystic ovary syndrome. *J Obstet Gynaecol*. 2021 Jun 25;1-11.
- Place NJ, Prado AM, Faykoo-Martinez M, et al. Germ cell nests in adult ovaries and an unusually large ovarian reserve in the naked mole-rat. *Reproduction*. 2021 Jan;161(1):89-98.
- McEvoy MJ, Sinderewicz E, Creedon L, et al. Death Processes in Bovine Theca and Granulosa Cells Modelled and Analysed Using a Systems Biology Approach. *Int J Mol Sci*. 2021 May 5;22(9):4888.
- Luciano AM, Barros RG, Soares ACS, et al. Recreating the Follicular Environment: A Customized Approach for In Vitro Culture of Bovine Oocytes Based on the Origin and Differentiation State. *Methods Mol Biol*. 2021;2273:1-15.
- Subramani E, Frigo DE. Mitochondria in Metabolic Syndrome, Reproduction and Transgenerational Inheritance-Ongoing Debates and Emerging Links. *Endocrinology*. 2021 Jan 1;162(1):bqaa182.
- Patil K, Shinde G, Hinduja I, Mukherjee S. Compromised Cumulus-Oocyte Complex Matrix Organization and Expansion in Women with PCOS. *Reprod Sci*. 2021 Nov 8. Online ahead of print.
- Slaidina M, Gupta S, Banisch TU, Lehmann R. A single-cell atlas reveals unanticipated cell type complexity in *Drosophila* ovaries. *Genome Res*. 2021 Oct;31(10):1938-1951.
- Li M, Zeng Z, Zhang A, et al. WGCNA Analysis Identifies Polycystic Ovary Syndrome-Associated Circular RNAs That Interact with RNA-Binding Proteins and Sponge miRNAs. *Int J Gen Med*. 2021 Nov 23;14:8737-8751.
- Rydze RT, Patton BK, Briley SM, et al. Deletion of Gremlin-2 alters estrous cyclicity and disrupts female fertility in mice†. *Biol Reprod*. 2021 Nov 15;105(5):1205-1220.
- Bao Y, Yao X, Li X, et al. INHBA transfection regulates proliferation, apoptosis and hormone synthesis in sheep granulosa cells. *Theriogenology*. 2021 Nov;175:111-122.
- Ford EA, Frost ER, Beckett EL, et al. Transcriptomic profiling of neonatal mouse granulosa cells reveals new insights into primordial follicle activation†. *Biol Reprod*. 2021 Oct 21;ioab193.
- França MM, Mendonça BB. Genetics of ovarian insufficiency and defects of folliculogenesis. *Best Pract Res Clin Endocrinol Metab*. 2021 Oct 14;101594.
- Li S, Wang J, Zhang H, et al. Transcriptome profile of goat folliculogenesis reveals the interaction of oocyte and granulosa cell in correlation with different fertility population. *Sci Rep*. 2021 Aug 3;11(1):15698.
- Jones A, Bernabé BP, Padmanabhan V, Li J, Shikanov A. Capitalizing on transcriptome profiling to optimize and identify targets for promoting early murine folliculogenesis in vitro. *Sci Rep*. 2021 Jun 15;11(1):12517.
- Dompe C, Kulus M, Stefańska K, et al. Human Granulosa Cells-Stemness Properties, Molecular Cross-Talk and Follicular Angiogenesis. *Cells*. 2021 Jun 5;10(6):1396.
- Yu L, Wang L, Tao W, et al. LHCGR and ALMS1 defects likely cooperate in the development of polycystic ovary syndrome indicated by double-mutant mice. *J Genet Genomics*. 2021 May 20;48(5):384-395.
- Chakravarthi VP, Ratri A, Masumi S, et al. Granulosa cell genes that regulate ovarian follicle development beyond the antral stage: The role of estrogen receptor  $\beta$ . *Mol Cell Endocrinol*. 2021 May 15;528:111212.
- Fan X, Moustakas I, Bialecka M, et al. Single-Cell Transcriptomics Analysis of Human Small Antral Follicles. *Int J Mol Sci*. 2021 Nov 4;22(21):11955.

- Alvi SM, Zayed Y, Malik R, Peng C. The emerging role of microRNAs in fish ovary: A mini review. *Gen Comp Endocrinol*. 2021 Sep 15;311:113850.
- Okunomiya A, Horie A, Tani H, et al. Figla promotes secondary follicle growth in mature mice. *Sci Rep*. 2021 May 10;11(1):9842.
- Habara O, Logan CY, Kanai-Azuma M, Nusse R, Takase HM. WNT signaling in pre-granulosa cells is required for ovarian folliculogenesis and female fertility. *Development*. 2021 May 1;148(9):dev198846.
- Wołodko K, Castillo-Fernandez J, Kelsey G, Galvão A. Revisiting the Impact of Local Leptin Signaling in Folliculogenesis and Oocyte Maturation in Obese Mothers. *Int J Mol Sci*. 2021 Apr 20;22(8):4270.
- Wang S, Gong Y, Wang Z, et al. Cell-specific network analysis of human folliculogenesis reveals network rewiring in antral stage oocytes. *J Cell Mol Med*. 2021 Mar;25(6):2851-2860.
- Pla I, Sanchez A, Pors SE, et al. Proteome of fluid from human ovarian small antral follicles reveals insights in folliculogenesis and oocyte maturation. *Hum Reprod*. 2021 Feb 18;36(3):756-770.
- Chen Y, Wang X, Yang C, et al. A mouse model reveals the events and underlying regulatory signals during the gonadotrophin-dependent phase of follicle development. *Mol Hum Reprod*. 2020 Dec 10;26(12):920-937.
- Safdar M, Liang A, Rajput AS, et al. Orexin-A Regulates Follicular Growth, Proliferation, Cell Cycle and Apoptosis in Mouse Primary Granulosa Cells via the AKT/ERK Signaling Pathway. *Molecules*. 2021 Sep 16;26(18):5635.
- Qu J, Niu H, Wang J, Wang Q, Li Y. Potential mechanism of lead poisoning to the growth and development of ovarian follicle. *Toxicology*. 2021 Jun 15;457:152810.
- Patil K, Hinduja I, Mukherjee S. Alteration in angiogenic potential of granulosa-lutein cells and follicular fluid contributes to luteal defects in polycystic ovary syndrome. *Hum Reprod*. 2021 Mar 18;36(4):1052-1064.
- Li Z, Jiang J, Yi X, et al. miR-18b regulates the function of rabbit ovary granulosa cells. *Reprod Fertil Dev*. 2021 Mar;33(5):363-371.
- Wang J, Chu K, Wang Y, et al. Procr-expressing granulosa cells are highly proliferative and are important for follicle development. *iScience*. 2021 Jan 19;24(2):102065.
- Olsen KW, Castillo-Fernandez J, Chan AC, et al. Identification of a unique epigenetic profile in women with diminished ovarian reserve. *Fertil Steril*. 2021 Mar;115(3):732-741.
- Elvin JA, Yan C, Wang P, Nishimori K, Matzuk MM. Molecular characterization of the follicle defects in the growth differentiation factor 9-deficient ovary. *Mol Endocrinol*. 1999 Jun;13(6):1018-34.
- Alaee S, Asadollahpour R, Colagar AH, Talaei-Khozani T. The decellularized ovary as a potential scaffold for maturation of preantral ovarian follicles of prepubertal mice. *Syst Biol Reprod Med*. 2021 Dec;67(6):413-427.
- Akin N, Le AH, Ha UDT, et al. Positive effects of amphiregulin on human oocyte maturation and its molecular drivers in patients with polycystic ovary syndrome. *Hum Reprod*. 2021 Nov 6;deab237.
- Fan X, Moustakas I, Bialecka M, et al. Single-Cell Transcriptomics Analysis of Human Small Antral Follicles. *Int J Mol Sci*. 2021 Nov 4;22(21):11955.
- Luo X, Chang H-M, Yi Y, et al. Bone morphogenetic protein 2 upregulates SERPINE2 expression through noncanonical SMAD2/3 and p38 MAPK signaling pathways in human granulosa-lutein cells. *FASEB J*. 2021 Sep;35(9):e21845.
- Chen M, Dong F, Chen M, et al. PRMT5 regulates ovarian follicle development by facilitating Wt1 translation. *Elife*. 2021 Aug 27;10:e68930.

- Yoshino T, Suzuki T, Nagamatsu G, et al. Generation of ovarian follicles from mouse pluripotent stem cells. *Science*. 2021 Jul 16;373(6552):eabe0237.
- Ansere VA, Ali-Mondal S, Sathiaselan R, et al. Cellular hallmarks of aging emerge in the ovary prior to primordial follicle depletion. *Mech Ageing Dev*. 2021 Mar;194:111425.
- Zeng J, Sun Y, Li X, et al. 2,5-Hexanedione influences primordial follicular development in cultured neonatal mouse ovaries by interfering with the PI3K signaling pathway via miR-214-3p. *Toxicol Appl Pharmacol*. 2020 Dec 15;409:115335.
- Bezerra FTG, Dau AMP, Van Den Hurk R, Silva JRV. Molecular characteristics of oocytes and somatic cells of follicles at different sizes that influence in vitro oocyte maturation and embryo production. *Domest Anim Endocrinol*. 2021 Jan;74:106485.
- Wu H-M, Chang H-M, Leung PCK. Gonadotropin-releasing hormone analogs: Mechanisms of action and clinical applications in female reproduction. *Front Neuroendocrinol*. 2021 Jan;60:100876.
- Fabová Z, Loncová B, Mlynček M, Sirotkin AV. Kisspeptin as autocrine/paracrine regulator of human ovarian cell functions: Possible interrelationships with FSH and its receptor. *Reprod Biol*. 2021 Nov 26;22(1):100580.
- Chaudhary H, Patel J, Jain NK, Joshi R. The role of polymorphism in various potential genes on polycystic ovary syndrome susceptibility and pathogenesis. *J Ovarian Res*. 2021 Sep 26;14(1):125.
- de Carvalho Delgado J, Dos Santos Hamilton TR, Mendes CM, et al. Bone morphogenetic protein 15 supplementation enhances cumulus expansion, nuclear maturation and progesterone production of in vitro-matured bovine cumulus-oocyte complexes. *Reprod Domest Anim*. 2021 May;56(5):754-763.
- Xu F, Lawson MS, Bean Y, et al. Matrix-free 3D culture supports human follicular development from the unilaminar to the antral stage in vitro yielding morphologically normal metaphase II oocytes. *Hum Reprod*. 2021 Apr 20;36(5):1326-1338.
- Anamthathmakula P, Winuthayanon W. Prostaglandin-Endoperoxide Synthase 2 (PTGS2) in the Oviduct: Roles in Fertilization and Early Embryo Development. *Endocrinology*. 2021 Apr 1;162(4):bqab025.
- Chen H, Cheng S, Xiong W, Tan X. The lncRNA-miRNA-mRNA ceRNA network in mural granulosa cells of patients with polycystic ovary syndrome: an analysis of Gene Expression Omnibus data. *Ann Transl Med*. 2021 Jul;9(14):1156.
- Rotgers E, Nicol B, Rodriguez K, Rattan S, Flaws JA, Yao HH-C. Constitutive expression of Steroidogenic factor-1 (NR5A1) disrupts ovarian functions, fertility, and metabolic homeostasis in female mice. *FASEB J*. 2021 Aug;35(8):e21770.
- Chen L, Ni Z, Cai Z, et al. The Mechanism Exploration of Follicular Fluids on Granulosa Cell Apoptosis in Endometriosis-Associated Infertility. *Biomed Res Int*. 2021 Oct 28;2021:6464686.
- Li S, Wang J, Zhang H, et al. Transcriptome profile of goat folliculogenesis reveals the interaction of oocyte and granulosa cell in correlation with different fertility population. *Sci Rep*. 2021 Aug 3;11(1):15698.
- Dong Y, Lyu L, Zhang D, et al. Integrated lncRNA and mRNA Transcriptome Analyses in the Ovary of *Cynoglossus semilaevis* Reveal Genes and Pathways Potentially Involved in Reproduction. *Front Genet*. 2021 May 19;12:671729.
- Buigues A, Marchante M, de Miguel-Gómez L, et al. Stem cell-secreted factor therapy regenerates the ovarian niche and rescues follicles. *Am J Obstet Gynecol*. 2021 Jul;225(1):65.e1-65.e14.
- Zhang Y, Wang Y, Feng X, et al. Oocyte-derived microvilli control female fertility by optimizing ovarian follicle selection in mice. *Nat Commun*. 2021 May 5;12(1):2523.

- Sou IF, Pryce RM, Tee W-W, McClurg UL. Meiosis initiation: a story of two sexes in all creatures great and small. *Biochem J*. 2021 Oct 29;478(20):3791-3805.
- Ben Maamar M, Nilsson EE, Skinner MK. Epigenetic transgenerational inheritance, gametogenesis and germline development†. *Biol Reprod*. 2021 Sep 14;105(3):570-592.
- Qi X, Yun C, Sun L, Xia J, et al. Gut microbiota-bile acid-interleukin-22 axis orchestrates polycystic ovary syndrome. *Nat Med*. 2019 Aug;25(8):1225-1233.
- Ramly B, Afiqah-Aleng N, Mohamed-Hussein ZA. Protein-Protein Interaction Network Analysis Reveals Several Diseases Highly Associated with Polycystic Ovarian Syndrome. *Int J Mol Sci*. 2019 Jun 18;20(12).
- Wang XY, Qin YY. Long non-coding RNAs in biology and female reproductive disorders. *Front Biosci (Landmark Ed)*. 2019 Mar 1;24:750-764.
- Henriques MC, Loureiro S, Fardilha M, Herdeiro MT. Exposure to mercury and human reproductive health: A systematic review. *Reprod Toxicol*. 2019 Apr;85:93-103.
- Amjadi F, Mehdizadeh M, Ashrafi M, et al. Distinct changes in the proteome profile of endometrial tissues in polycystic ovary syndrome compared with healthy fertile women. *Reprod Biomed Online*. 2018 Aug;37(2):184-200.
- Shukla A, Dahiya S, Onteru SK, Singh D. Differentially expressed miRNA-210 during follicular-luteal transition regulates pre-ovulatory granulosa cell function targeting HRas and EFNA3. *J Cell Biochem*. 2018 Nov;119(10):7934-7943.
- Rodriguez A, Briley SM, Patton BK, et al. Loss of the E2 SUMO-conjugating enzyme Ube2i in oocytes during ovarian folliculogenesis causes infertility in mice. *Development*. 2019 Dec 2;146(23).
- Cai H, Liu B, Wang H, Sun G, Feng L, et al. SP1 governs primordial folliculogenesis by regulating pregranulosa cell development in mice. *J Mol Cell Biol*. 2019 Jul 8. pii: mjj059. doi: 10.1093/jmcb/mjz059. [Epub ahead of print]
- Fan Y, Zhang C, Zhu G. Profiling of RNA N6-methyladenosine methylation during follicle selection in chicken ovary. *Poult Sci*. 2019 Nov 1;98(11):6117-6124.
- Sagvekar P, Kumar P, Mangoli V, Desai S, Mukherjee S. DNA methylome profiling of granulosa cells reveals altered methylation in genes regulating vital ovarian functions in polycystic ovary syndrome. *Clin Epigenetics*. 2019 Apr 11;11(1):61.
- Wang T, Liu Y, Lv M, Xing Q, Zhang Z, He X, Xu Y, Wei Z, Cao Y. miR-323-3p regulates the steroidogenesis and cell apoptosis in polycystic ovary syndrome (PCOS) by targeting IGF-1. *Gene*. 2019 Jan 30;683:87-100.
- Zhao J, Xu J, Wang W, Zhao H, et al. Long non-coding RNA LINC-01572:28 inhibits granulosa cell growth via a decrease in p27 (Kip1) degradation in patients with polycystic ovary syndrome. *EBioMedicine*. 2018 Oct;36:526-538.
- Tran DN, Jung EM, Yoo YM, Ahn C, et al. Depletion of follicles accelerated by combined exposure to phthalates and 4-vinylcyclohexene diepoxide, leading to premature ovarian failure in rats. *Reprod Toxicol*. 2018 Sep;80:60-67.
- Khristi V, Chakravarthi VP, Singh P, et al. ESR2 regulates granulosa cell genes essential for follicle maturation and ovulation. *Mol Cell Endocrinol*. 2018 Oct 15;474:214-226.
- Shukla A, Dahiya S, Onteru SK, Singh D. Differentially expressed miRNA-210 during follicular-luteal transition regulates pre-ovulatory granulosa cell function targeting HRas and EFNA3. *J Cell Biochem*. 2018 Nov;119(10):7934-7943.
- Ragazzini R, Pérez-Palacios R, Baymaz IH, et al. EZHIP constrains Polycomb Repressive Complex 2 activity in germ cells. *Nat Commun*. 2019 Aug 26;10(1):3858.

- Spiller C, Bowles J. Sexually dimorphic germ cell identity in mammals. *Curr Top Dev Biol*. 2019;134:253-288.
- Spears N, Lopes F, Stefansdottir A, Rossi V, De Felici M, Anderson RA, Klinger FG. Ovarian damage from chemotherapy and current approaches to its protection. *Hum Reprod Update*. 2019 Nov 5;25(6):673-693.
- Cuomo D, Porreca I, Ceccarelli M, Threadgill DW, et al. Transcriptional landscape of mouse-aged ovaries reveals a unique set of non-coding RNAs associated with physiological and environmental ovarian dysfunctions. *Cell Death Discov*. 2018 Dec 5;4:112.
- Landry DA, Rossi-Perazza L, Lafontaine S, Sirard MA. Expression of atresia biomarkers in granulosa cells after ovarian stimulation in heifers. *Reproduction*. 2018 Sep;156(3):239-248.
- Zhang J, Liu Y, Yao W, Li Q, Liu H, Pan Z. Initiation of follicular atresia: gene networks during early atresia in pig ovaries. *Reproduction*. 2018 Jul;156(1):23-33
- Ford E, Beckett EL, Roman S, McLaughlin EA, Sutherland J. Advances in human primordial follicle activation and premature ovarian insufficiency. *Reproduction*. 2019 Aug 1. pii: REP-19-0201.R2. doi: 10.1530/REP-19-0201. [Epub ahead of print]
- Shimizu K, Nakamura T, Bayasula, Nakanishi N, et al. Molecular mechanism of FSHR expression induced by BMP15 in human granulosa cells. *J Assist Reprod Genet*. 2019 Jun;36(6):1185-1194.
- Sun XF, Li YP, Pan B, Wang YF, Li J, Shen W. Molecular regulation of miR-378 on the development of mouse follicle and the maturation of oocyte in vivo. *Cell Cycle*. 2018;17(18):2230-2242.
- Cui L, Fang L, Mao X, Chang HM, Leung PCK, Ye Y. GDNF-Induced Downregulation of miR-145-5p Enhances Human Oocyte Maturation and Cumulus Cell Viability. *J Clin Endocrinol Metab*. 2018 Jul 1;103(7):2510-2521.
- Huang J, Ding Y, Li Z. The regulation of the follicular synchronization and sensitivity of rats with PCOS by AMH during prolonged pituitary downregulation. *Gene*. 2019 Dec 30;721:144106.
- Roness H, Spector I, Leichtmann-Bardoogo Y, Savino AM, Dereh-Haim S, Meiorow D. Pharmacological administration of recombinant human AMH rescues ovarian reserve and preserves fertility in a mouse model of chemotherapy, without interfering with anti-tumoural effects. *J Assist Reprod Genet*. 2019 Sep;36(9):1793-1803.
- Yan H, Wen J, Zhang T, Zheng W, He M, et al. Oocyte-derived E-cadherin acts as a multiple functional factor maintaining the primordial follicle pool in mice. *Cell Death Dis*. 2019 Feb 15;10(3):160.
- Hardy K, Mora JM, Dunlop C, Carzaniga R, Franks S, Fenwick MA. Nuclear exclusion of SMAD2/3 in granulosa cells is associated with primordial follicle activation in the mouse ovary. *J Cell Sci*. 2018 Sep 7;131(17).
- Podfigurna A, Meczekalski B, Petraglia F, Luisi S. Clinical, hormonal and metabolic parameters in women with PCOS with different combined oral contraceptives (containing chlormadinone acetate versus drospirenone). *J Endocrinol Invest*. 2019 Oct 25. doi: 10.1007/s40618-019-01133-3. [Epub ahead of print]
- Fan X, Bialecka M, Moustakas I, et al. Single-cell reconstruction of follicular remodeling in the human adult ovary. *Nat Commun*. 2019 Jul 18;10(1):3164.
- Matvere A, Teino I, Varik I, Kuuse S, Tiido T, Kristjuhan A, Maimets T. FSH/LH-Dependent Upregulation of Ahr in Murine Granulosa Cells Is Controlled by PKA Signaling and Involves Epigenetic Regulation. *Int J Mol Sci*. 2019 Jun 23;20(12).
- Stanley JA, Arosh JA, Hoyer PB, Banu SK. Ex Vivo Fetal Whole Ovarian Culture Model: An Essential Tool for Studies in Reproductive Toxicology and Pharmacology. *Methods Mol Biol*. 2019;1965:107-127.

- Tanaka Y, Matsuzaki T, Tanaka N, Iwasa T, Kuwahara A, Irahara M. Activin effects on follicular growth in in vitro preantral follicle culture. *J Med Invest*. 2019;66(1.2):165-171.
- Candelaria NR, Padmanabhan A, Stossi F, Ljungberg MC, et al. VCAM1 Is Induced in Ovarian Theca and Stromal Cells in a Mouse Model of Androgen Excess. *Endocrinology*. 2019 Jun 1;160(6):1377-1393.
- Mossa F, Ireland JJ. Physiology and endocrinology symposium: Anti-Müllerian hormone: a biomarker for the ovarian reserve, ovarian function, and fertility in dairy cows. *J Anim Sci*. 2019 Apr 3;97(4):1446-1455.
- Wang T, Liu Y, Lv M, Xing Q, Zhang Z, He X, Xu Y, Wei Z, Cao Y. miR-323-3p regulates the steroidogenesis and cell apoptosis in polycystic ovary syndrome (PCOS) by targeting IGF-1. *Gene*. 2019 Jan 30;683:87-100.
- Schmidt VM, Isachenko E, Rappl G, Rahimi G, Hanstein B, Morgenstern B, Mallmann P, Isachenko V. Construction of human artificial ovary from cryopreserved ovarian tissue: Appearance of apoptosis and necrosis after enzymatic isolation of follicles. *Cryobiology*. 2018 Oct;84:10-14.
- Hamilton KJ, Hewitt SC, Arao Y, Korach KS. (2017) Estrogen Hormone Biology. *Curr Top Dev Biol*. 2017;125:109-146.
- Yang Z, Sun J, Hu Y, Wang F, Wang X, et al. (2017) Histone H1 defect in escort cells triggers germline tumor in Drosophila ovary. *Dev Biol*. 424(1):40-49.
- Bevilacqua A, Bizzarri M. (2016) Physiological role and clinical utility of inositols in polycystic ovary syndrome. *Best Pract Res Clin Obstet Gynaecol*. 37:129-139.
- Mohammadi S, Kayedpoor P, Karimzadeh-Bardei L, Nabiuni M. (2017) The Effect of Curcumin on TNF- $\alpha$ , IL-6 and CRP Expression in a Model of Polycystic Ovary Syndrome as an Inflammation State. *J Reprod Infertil*. 18(4):352-360.
- Li Y, Zhang L, Hu Y, Chen M, Han F, Qin Y, Chen M, Cui X, Duo S, Tang F, Gao F. (2017)  $\beta$ -Catenin directs the transformation of testis Sertoli cells to ovarian granulosa-like cells by inducing Foxl2 expression. *J Biol Chem*. 2017 Oct 27;292(43):17577-17586.
- Herndon MK, Law NC, Donaubaue EM, Kyriess B, Hunzicker-Dunn M. (2016) Forkhead box O member FOXO1 regulates the majority of follicle-stimulating hormone responsive genes in ovarian granulosa cells. *Mol Cell Endocrinol*. 434:116-26.
- Huang K, Wang Y, Zhang T, He M, Sun G, Wen J, Yan H, Cai H, Yong C, Xia G, Wang C. (2018) JAK signaling regulates germline cyst breakdown and primordial follicle formation in mice. *Biol Open*. 17;7(1).
- Richani D, Gilchrist RB. (2108) The epidermal growth factor network: role in oocyte growth, maturation and developmental competence. *Hum Reprod Update*. 24(1):1-14.
- Kranc W, Budna J, Kahan R, et al. (2017) Molecular basis of growth, proliferation, and differentiation of mammalian follicular granulosa cells. *J Biol Regul Homeost Agents*. 31(1):1-8.
- Li L, Wang B, Zhang W, Chen B, et al. (2017) A homozygous NOBOX truncating variant causes defective transcriptional activation and leads to primary ovarian insufficiency. *Hum Reprod*. 32(1):248-255.
- Chang HM, Qiao J, Leung PC. (2016) Oocyte-somatic cell interactions in the human ovary-novel role of bone morphogenetic proteins and growth differentiation factors. *Hum Reprod Update*. 23(1):1-18.
- Vanornny DA, Mayo KE. (2017) The role of Notch signaling in the mammalian ovary. *Reproduction*. 153(6):R187-R204.
- Li D, Xu D, Zou Y, et al. (2017) Non-coding RNAs and ovarian diseases (Review). *Mol Med Rep*. 15(4):1435-1440.

- Tanaka M. (2016) Germline stem cells are critical for sexual fate decision of germ cells. *Bioessays*. 38(12):1227-1233.
- Laven JS. (2016) Primary Ovarian Insufficiency. *Semin Reprod Med*. 34(4):230-4.
- De Leo V, Musacchio MC, Cappelli V, Massaro MG, Morgante G, Petraglia F. (2016) Genetic, hormonal and metabolic aspects of PCOS: an update. *Reprod Biol Endocrinol*. 16;14(1):38.
- Bedoschi G, Navarro PA, Oktay K. (2016) Chemotherapy-induced damage to ovary: mechanisms and clinical impact. *Future Oncol*. 12(20):2333-44.
- Huang X, Liu C, Hao C, Tang Q, Liu R, Lin S, Zhang L, Yan W. (2016) Identification of altered microRNAs and mRNAs in the cumulus cells of PCOS patients: miRNA-509-3p promotes oestradiol secretion by targeting MAP3K8. *Reproduction*. 151(6):643-55.
- Qin Y, Jiao X, Simpson JL, Chen ZJ. (2015) Genetics of primary ovarian insufficiency: new developments and opportunities. *Hum Reprod Update*. 21(6):787-808.
- Bhartiya D, Parte S, Patel H, et al (2016) Novel Action of FSH on Stem Cells in Adult Mammalian Ovary Induces Postnatal Oogenesis and Primordial Follicle Assembly. *Stem Cells Int*. 2016:5096596.
- Price CA. (2016) Mechanisms of fibroblast growth factor signaling in the ovarian follicle. *J Endocrinol*. 228(2):R31-43.
- Ng A, Barker N. (2015) Ovary and fimbrial stem cells: biology, niche and cancer origins. *Nat Rev Mol Cell Biol*. 16(10):625-38.
- Grieve KM, McLaughlin M, Dunlop CE, Telfer EE, Anderson RA. (2015) The controversial existence and functional potential of oogonial stem cells. *Maturitas*. 82(3):278-81.
- Li Y, Fang Y, Liu Y, Yang X. (2015) MicroRNAs in ovarian function and disorders. *J Ovarian Res*. 1;8:51.
- Ozakupinar OB, Maurer AM, Ozsavci D. (2015) Ovarian stem cells: From basic to clinical applications. *World J Stem Cells*. 26;7(4):757-68.
- McGinnis LK, Luense LJ, Christenson LK. (2015) MicroRNA in Ovarian Biology and Disease. *Cold Spring Harb Perspect Med*. 18;5(9):a022962.
- Hummitzsch K, Anderson RA, Wilhelm D, et al. (2015) Stem cells, progenitor cells, and lineage decisions in the ovary. *Endocr Rev*. 36(1):65-91.
- Thorne JT1, Segal TR2, Chang S3, et al. (2015) Dynamic reciprocity between cells and their microenvironment in reproduction. *Biol Reprod*. 92(1):25.
- Slaidina M, Lehmann R. (2014) Translational control in germline stem cell development. *J Cell Biol*. 207(1):13-21.
- Hsueh AJ, Kawamura K, Cheng Y, Fauser BC. (2015) Intraovarian control of early folliculogenesis. *Endocr Rev*. 36(1):1-24.
- Farrell A, McLoughlin N, Milne JJ, Marison IW, Bones J. (2014) Application of multi-omics techniques for bioprocess design and optimization in chinese hamster ovary cells. *J Proteome Res*. 13(7):3144-59.
- Shea LD, Woodruff TK, Shikanov A. (2014) Bioengineering the ovarian follicle microenvironment. *Annu Rev Biomed Eng*. 11;16:29-52.
- Qiao J, Wang ZB, Feng HL, et al. (2014) The root of reduced fertility in aged women and possible therapeutic options: current status and future prospects. *Mol Aspects Med*. 38:54-85.
- Fitzgerald JB, George J, Christenson LK. (2016) Non-coding RNA in Ovarian Development and Disease. *Adv Exp Med Biol*. 886:79-93.
- Qin Y, Jiao X, Simpson JL, Chen ZJ. (2015) Genetics of primary ovarian insufficiency: new developments and opportunities. *Hum Reprod Update*. 21(6):787-808.

- Ren Y, Suzuki H, Jagarlamudi K, et al. (2015) Lhx8 regulates primordial follicle activation and postnatal folliculogenesis. *BMC Biol.* 16;13:39.
- Hua J, Xu B, Yang Y, Ban R, et al. (2015) Follicle Online: an integrated database of follicle assembly, development and ovulation. *Database (Oxford)*. 29;2015:bav036.
- Bilgin EM, Kovanci E. (2015) Genetics of premature ovarian failure. *Curr Opin Obstet Gynecol.* 27(3):167-74.
- Liu C, Peng J, Matzuk MM, Yao HH. (2015) Lineage specification of ovarian theca cells requires multicellular interactions via oocyte and granulosa cells. *Nat Commun.* 28;6:6934.
- Heeren AM, van Iperen L, Klootwijk DB, et al. (2015) Development of the follicular basement membrane during human gametogenesis and early folliculogenesis. *BMC Dev Biol.* 21;15:4.
- Chen Z, Kang X, Wang L, et al. (2015) Rictor/mTORC2 pathway in oocytes regulates folliculogenesis, and its inactivation causes premature ovarian failure. *J Biol Chem.* 6;290(10):6387-96.
- Wang Y, Teng Z, Li G, Mu X, et al. (2015) Cyclic AMP in oocytes controls meiotic prophase I and primordial folliculogenesis in the perinatal mouse ovary. *Development.* 15;142(2):343-51.
- Dumesic DA, Meldrum DR, Katz-Jaffe MG, et al. (2015) Oocyte environment: follicular fluid and cumulus cells are critical for oocyte health. *Fertil Steril.* 103(2):303-16.
- Park M, Suh DS, Lee K, Bae J. (2014) Positive cross talk between FOXL2 and antimüllerian hormone regulates ovarian reserve. *Fertil Steril.* 102(3):847-855.e1.
- Feng XL, Sun YC, Zhang M, et al. (2014) Insulin regulates primordial-follicle assembly in vitro by affecting germ-cell apoptosis and elevating oestrogen. *Reprod Fertil Dev.* 27(8) 1197-1204.
- Sun J, Wei HM, Xu J, Chang JF, et al. (2015) Histone H1-mediated epigenetic regulation controls germline stem cell self-renewal by modulating H4K16 acetylation. *Nat Commun.* 19;6:8856.
- Lan CW, Chen MJ, Tai KY, et al., (2015) Functional microarray analysis of differentially expressed genes in granulosa cells from women with polycystic ovary syndrome related to MAPK/ERK signaling. *Sci Rep.* 13;5:14994.
- Skory RM, Xu Y, Shea LD, Woodruff TK. (2015) Engineering the ovarian cycle using in vitro follicle culture. *Hum Reprod.* 30(6):1386-95.
- Vázquez MJ, Romero-Ruiz A, Tena-Sempere M. (2015) Roles of leptin in reproduction, pregnancy and polycystic ovary syndrome: consensus knowledge and recent developments. *Metabolism.* 64(1):79-91.
- Wigglesworth K, Lee KB, Emori C, Sugiura K, Eppig JJ. (2015) Transcriptomic diversification of developing cumulus and mural granulosa cells in mouse ovarian follicles. *Biol Reprod.* 92(1):23.
- Perry JR, Day F, Elks CE, Sulem P, et al., (2014) Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature.* 514(7520):92-7.
- Dinicola S, Chiu TT, Unfer V, et al., (2014) The rationale of the myo-inositol and D-chiro-inositol combined treatment for polycystic ovary syndrome. *J Clin Pharmacol.* 54(10):1079-92.
- Aiken CE, Tarry-Adkins JL, et al., (2015) Decreased ovarian reserve, dysregulation of mitochondrial biogenesis, and increased lipid peroxidation in female mouse offspring exposed to an obesogenic maternal diet. *FASEB J.* 2015 Dec 23. pii: fj.15-280800. [Epub ahead of print]
- Peterson JS, Timmons AK, Mondragon AA, McCall K. (2015) The End of the Beginning: Cell Death in the Germline. *Curr Top Dev Biol.* 114:93-119.
- Grive KJ, Freiman RN. (2015) The developmental origins of the mammalian ovarian reserve. *Development.* 1;142(15):2554-63.
- Bayne RA, Kinnell HL, Coutts SM, et al., (2015) GDF9 is transiently expressed in oocytes before follicle formation in the human fetal ovary and is regulated by a novel NOBOX transcript. *PLoS One.* 19;10(3):e0119819.

- Zhang H, Risal S, Gorre N, et al., (2014) Somatic cells initiate primordial follicle activation and govern the development of dormant oocytes in mice. *Curr Biol.* 3;24(21):2501-8.
- Estienne A, Pierre A, di Clemente N, et al. (2015) Anti-Müllerian hormone regulation by the bone morphogenetic proteins in the sheep ovary: deciphering a direct regulatory pathway. *Endocrinology.* 156(1):301-13.
- Pan L, Gong W, Zhou Y, Li X, Yu J, Hu S. (2014) A comprehensive transcriptomic analysis of infant and adult mouse ovary. *Genomics Proteomics Bioinformatics.* 12(5):239-48.
- Nilsson EE, Larsen G, Skinner MK. (2014) Roles of Gremlin 1 and Gremlin 2 in regulating ovarian primordial to primary follicle transition. *Reproduction.* 147(6):865-74.
- Puttabyatappa M, Cardoso RC, Padmanabhan V. (2015) Effect of maternal PCOS and PCOS-like phenotype on the offspring's health. *Mol Cell Endocrinol.* 2015 Nov 27. pii: S0303-7207(15)30153-2. doi: 10.1016/j.mce.2015.11.030. [Epub ahead of print]
- Dumesic DA, Oberfield SE, Stener-Victorin E, et al. (2015) Scientific Statement on the Diagnostic Criteria, Epidemiology, Pathophysiology, and Molecular Genetics of Polycystic Ovary Syndrome. *Endocr Rev.* 36(5):487-525.
- Spritzer PM, Motta AB. (2015) Adolescence and polycystic ovary syndrome: current concepts on diagnosis and treatment. *Int J Clin Pract.* 69(11):1236-46.
- Luque-Ramírez M, Escobar-Morreale HF. (2014) Polycystic ovary syndrome as a paradigm for prehypertension, prediabetes, and preobesity. *Curr Hypertens Rep.* 16(12):500.
- Lin F, Fu YH, Han J, et al. (2014) Changes in the expression of Fox O1 and death ligand genes during follicular atresia in porcine ovary. *Genet Mol Res.* 28;13(3):6638-45.
- Roland AV, Moenter SM. (2014) Reproductive neuroendocrine dysfunction in polycystic ovary syndrome: insight from animal models. *Front Neuroendocrinol.* 35(4):494-511.
- Li Y, Fang Y, Liu Y, Yang X. (2015) MicroRNAs in ovarian function and disorders. *J Ovarian Res.* 1;8:51.
- Tiwari M, Prasad S, Tripathi A, et al., (2015) Apoptosis in mammalian oocytes: a review. *Apoptosis.* 20(8):1019-25.
- Johnson AL. (2015) Ovarian follicle selection and granulosa cell differentiation. *Poult Sci.* 94(4):781-5.
- Hsueh AJ, Kawamura K, Cheng Y, Fauser BC. (2015) Intraovarian control of early folliculogenesis. *Endocr Rev.* 36(1):1-24.
- Mereness AL, Murphy ZC, et al., (2016) Conditional Deletion of Bmal1 in Ovarian Theca Cells Disrupts Ovulation in Female Mice. *Endocrinology.* 157(2):913-27.
- Zielak-Steciwo AE, Browne JA, McGettigan PA, et al., (2014) Expression of microRNAs and their target genes and pathways associated with ovarian follicle development in cattle. *Physiol Genomics.* 46(19):735-45.
- Ortega HH, Marelli BE, Rey F, et al., (2015) Molecular aspects of bovine cystic ovarian disease pathogenesis. *Reproduction.* 149(6):R251-64.
- Sørensen AE, Wissing ML, Salö S, Englund AL, Dalgaard LT. (2014) MicroRNAs Related to Polycystic Ovary Syndrome (PCOS). *Genes (Basel).* 25;5(3):684-708.
- Gheorghisan-Galateanu AA, Hinescu ME, Enciu AM. (2014) Ovarian adult stem cells: hope or pitfall? *J Ovarian Res.* 4;7:71.
- Roland AV, Moenter SM. (2014) Reproductive neuroendocrine dysfunction in polycystic ovary syndrome: insight from animal models. *Front Neuroendocrinol.* 35(4):494-511.
- Rodríguez FM, Colombero M, Amweg AN, et al., (2015) Involvement of PAPP-A and IGFR1 in Cystic Ovarian Disease in Cattle. *Reprod Domest Anim.* 50(4):659-68.

- Munger SC, Natarajan A, Looger LL, et al. (2013) Fine time course expression analysis identifies cascades of activation and repression and maps a putative regulator of mammalian sex determination. *PLoS Genet.* 9(7):e1003630.
- Shafiee MN, Chapman C, Barrett D, et al. (2013) Reviewing the molecular mechanisms which increase endometrial cancer (EC) risk in women with polycystic ovarian syndrome (PCOS): time for paradigm shift? *Gynecol Oncol.* 131(2):489-92.
- Telfer EE, Zelinski MB. (2013) Ovarian follicle culture: advances and challenges for human and nonhuman primates. *Fertil Steril.* 99(6):1523-33.
- Telfer EE, McLaughlin M. (2012) Strategies to support human oocyte development in vitro. *Int J Dev Biol.* 2012;56(10-12):901-7.
- Adam M, Saller S, Ströbl S, et al. (2012) Decorin is a part of the ovarian extracellular matrix in primates and may act as a signaling molecule. *Hum Reprod.* 27(11):3249-58.
- Pate JL, Toyokawa K, Walusimbi S, Brzezicka E. (2010) The interface of the immune and reproductive systems in the ovary: lessons learned from the corpus luteum of domestic animal models. *Am J Reprod Immunol.* 64(4):275-86.
- Peddinti D, Memili E, Burgess SC. (2010) Proteomics-based systems biology modeling of bovine germinal vesicle stage oocyte and cumulus cell interaction. *PLoS One.* 21;5(6):e11240.
- Teino I, Matvere A, Kuuse S, et al. (2014) Transcriptional repression of the Ahr gene by LHCGR signaling in preovulatory granulosa cells is controlled by chromatin accessibility. *Mol Cell Endocrinol.* 25;382(1):292-301.
- Grøndahl ML, Borup R, Vikeså J, et al. (2013) The dormant and the fully competent oocyte: comparing the transcriptome of human oocytes from primordial follicles and in metaphase II. *Mol Hum Reprod.* 19(9):600-17.
- Liu Z, Castrillon DH, Zhou W, Richards JS. (2013) FOXO1/3 depletion in granulosa cells alters follicle growth, death and regulation of pituitary FSH. *Mol Endocrinol.* 27(2):238-52.
- Sobinoff AP, Sutherland JM, McLaughlin EA. (2013) Intracellular signalling during female gametogenesis. *Mol Hum Reprod.* 19(5):265-78.
- Richards JS, Liu Z, Kawai T, et al. (2012) Adiponectin and its receptors modulate granulosa cell and cumulus cell functions, fertility, and early embryo development in the mouse and human. *Fertil Steril.* 98(2):471-9.e1.
- Teino I, Kuuse S, Ingerpuu S, et al. (2012) The aryl hydrocarbon receptor regulates mouse Fshr promoter activity through an e-box binding site. *Biol Reprod.* 22;86(3):77.
- Reader KL, Heath DA, Lun S, et al. (2011) Signalling pathways involved in the cooperative effects of ovine and murine GDF9+BMP15-stimulated thymidine uptake by rat granulosa cells. *Reproduction.* 142(1):123-31.
- Psathaki OE, Hübner K, Sabour D, et al. (2011) Ultrastructural characterization of mouse embryonic stem cell-derived oocytes and granulosa cells. *Stem Cells Dev.* 20(12):2205-15.
- Oktem O, Buyuk E, Oktay K. (2011) Preantral follicle growth is regulated by c-Jun-N-terminal kinase (JNK) pathway. *Reprod Sci.* 18(3):269-76.
- Kowalewski MP, Dyson MT, Boos A, Stocco DM. (2010) Vasoactive intestinal peptide (VIP)-mediated expression and function of steroidogenic acute regulatory protein (StAR) in granulosa cells. *Mol Cell Endocrinol.* 26;328(1-2):93-103.
- Fan HY, O'Connor A, Shitanaka M, et al. (2010) Beta-catenin (CTNNB1) promotes preovulatory follicular development but represses LH-mediated ovulation and luteinization. *Mol Endocrinol.* 24(8):1529-42.

- Fan HY, Liu Z, Paquet M, et al. (2009) Cell type-specific targeted mutations of Kras and Pten document proliferation arrest in granulosa cells versus oncogenic insult to ovarian surface epithelial cells. *Cancer Res.* 15;69(16):6463-72.
- Hossain MM, Cao M, Wang Q, et al. (2013) Altered expression of miRNAs in a dihydrotestosterone-induced rat PCOS model. *J Ovarian Res.* 15;6(1):36.
- Nakamura S, Kurokawa H, Asakawa S, et al. (2009) Two distinct types of theca cells in the medaka gonad: germ cell-dependent maintenance of cyp19a1-expressing theca cells. *Dev Dyn.* 238(10):2652-7.
- Zheng W, Zhang H, Liu K. (2014) The two classes of primordial follicles in the mouse ovary: their development, physiological functions, and implications for future research. *Mol Hum Reprod.* Jan 20. [Epub ahead of print]
- Navalakhe RM, Jagtap DD, Nayak SU, et al. (2013) Effect of FSH receptor-binding inhibitor-8 on FSH-mediated granulosa cell signaling and proliferation. *Chem Biol Drug Des.* 82(2):178-88.
- Hunt PA, Lawson C, Gieske M, et al. (2012) Bisphenol A alters early oogenesis and follicle formation in the fetal ovary of the rhesus monkey. *Proc Natl Acad Sci U S A.* 23;109(43):17525-30.
- Adhikari D, Gorre N, Risal S, et al. (2012) The safe use of a PTEN inhibitor for the activation of dormant mouse primordial follicles and generation of fertilizable eggs. *PLoS One.* 7(6):e39034.
- Laan M, Grigorova M, Huhtaniemi IT. (2012) Pharmacogenetics of follicle-stimulating hormone action. *Curr Opin Endocrinol Diabetes Obes.* 19(3):220-7.
- Zhang HQ, Zhang XF, Zhang LJ, et al. (2012) Fetal exposure to bisphenol A affects the primordial follicle formation by inhibiting the meiotic progression of oocytes. *Mol Biol Rep.* 39(5):5651-7.
- Dixit H, Rao L, Padmalatha V, et al (2010) Genes governing premature ovarian failure. *Reprod Biomed Online*20(6):724-40.
- Richards JS, Pangas SA. (2010) The ovary: basic biology and clinical implications. *J Clin Invest.* 120(4):963-72.
- Nilsson EE, Skinner MK. (2009) Progesterone regulation of primordial follicle assembly in bovine fetal ovaries. *Mol Cell Endocrinol.* 10;313(1-2):9-16.
- Nilsson E, Dole G, Skinner MK. (2009) Neurotrophin NT3 promotes ovarian primordial to primary follicle transition. *Reproduction.* 138(4):697-707.
- Fan HY, Liu Z, Shimada M, Sterneck E, et al. (2009) MAPK3/1 (ERK1/2) in ovarian granulosa cells are essential for female fertility. *Science.* 15;324(5929):938-41.
- Barthelmess EK, Naz RK. (2014) Polycystic ovary syndrome: current status and future perspective. *Front Biosci (Elite Ed).* 1;6:104-19.
- Layman LC. (2013) The genetic basis of female reproductive disorders: etiology and clinical testing. *Mol Cell Endocrinol.* 6;370(1-2):138-48.
- Richards JS, Pangas SA. (2010) New insights into ovarian function. *Handb Exp Pharmacol.* (198):3-27.
- Grützner F et al. (2008) Reproductive biology in egg-laying mammals. *Sex Dev.* 2008;2(3):115-27.
- Ireland JL et al. (2008) Antral follicle count reliably predicts number of morphologically healthy oocytes and follicles in ovaries of young adult cattle. *Biol Reprod.* 79(6):1219-25.
- Sun QY et al. (2008) Oocyte-specific knockout: a novel in vivo approach for studying gene functions during folliculogenesis, oocyte maturation, fertilization, and embryogenesis. *Biol Reprod.* 79(6):1014-20.
- Rodrigues P et al. (2008) Oogenesis: Prospects and challenges for the future. *J Cell Physiol.* 216(2):355-65.

- Richards JS et.al. (2008) Immune-like mechanisms in ovulation. *Trends Endocrinol Metab.* Aug;19(6):191-6.
- Best D et. al. (2008) Sdmg1 is a conserved transmembrane protein associated with germ cell sex determination and germline-soma interactions in mice. *Development.* 135(8):1415-25.
- Hunt PA et. al. (2008) Human female meiosis: what makes a good egg go bad? *Trends Genet.* 24(2):86-93.
- Oktem O et. al. (2008) The ovary: anatomy and function throughout human life. *Ann N Y Acad Sci.* 1127:1-9. Review.
- Stocco C (2008) Aromatase expression in the ovary: hormonal and molecular regulation. *Steroids.* 73(5):473-87.
- Tatone C et. al. (2008) Cellular and molecular aspects of ovarian follicle ageing. *Hum Reprod Update.* 14(2):131-42.
- Choi Y et. al. (2008) Germ cell-specific transcriptional regulator sohlh2 is essential for early mouse folliculogenesis and oocyte-specific gene expression. *Biol Reprod.* 79(6):1176-82.
- Monné M et. al. (2008) Crystal structure of the ZP-N domain of ZP3 reveals the core fold of animal egg coats. *Nature.* 4;456(7222):653-7.
- Lee HJ et. al. (2007) Bone marrow transplantation generates immature oocytes and rescues long-term fertility in a preclinical mouse model of chemotherapy-induced premature ovarian failure. *J Clin Oncol.* 1;25(22):3198-204.
- Kimura S et. al. (2007) Androgen receptor function in folliculogenesis and its clinical implication in premature ovarian failure. *Trends Endocrinol Metab.* 18(5):183-9.
- Song JL et. al. (2006) Oogenesis: single cell development and differentiation. *Dev Biol* 300(1):385-405.
- Shimada M et. al. (2006) Paracrine and autocrine regulation of epidermal growth factor-like factors in cumulus oocyte complexes and granulosa cells: key roles for prostaglandin synthase 2 and progesterone receptor. *Mol Endocrinol* 20(6):1352-65.
- Dixit H et. al. (2006) Missense mutations in the BMP15 gene are associated with ovarian failure. *Hum Genet* 119(4):408-15.
- Eggan K et. al. (2006) Ovulated oocytes in adult mice derive from non-circulating germ cells. *Nature* 441(7097):1109-14.
- Kremenskov M et. al. (2006) DNA methylation profiles of donor nuclei cells and tissues of cloned bovine fetuses. *J Reprod Dev* 52(2):259-66.
- Thomas FH and Vanderhyden BC (2006) Oocyte-granulosa cell interactions during mouse follicular development: regulation of kit ligand expression and its role in oocyte growth. *Reprod Biol Endocrinol.* 4:19.
- De La Fuente R (2006) Chromatin modifications in the germinal vesicle (GV) of mammalian oocytes. *Dev Biol* 292(1):1-12.
- Wickenheisser JK et. al. (2006) Human ovarian theca cells in culture. *Trends Endocrinol Metab* 17(2):65-71.
- Hsieh M et. al. (2005) Mice null for Frizzled4 (Fzd4<sup>-/-</sup>) are infertile and exhibit impaired corpora lutea formation and function. *Biol Reprod* 73(6):1135-46.
- Matsuda-Minehata F et. al. (2005) Changes in expression of anti-apoptotic protein, cFLIP, in granulosa cells during follicular atresia in porcine ovaries. *Mol Reprod Dev.* 72(2):145-51.
- Wickenheisser JK et. al. (2005) Retinoids and retinol differentially regulate steroid biosynthesis in ovarian theca cells isolated normal cycling in women and women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 90(8):4858-65.

- Xu J et. al. (2005) Discovery of LH-regulated genes in the primate corpus luteum. *Mol Hum Reprod* 11(3):151-9.
- Richards JS et. al. (2005) Regulated expression of ADAMTS family members in follicles and cumulus oocyte complexes: Evidence for specific and redundant patterns during ovulation. *Biol Reprod* 72(5):1241-55.
- Frase HM and Duncan WC (2005) Vascular morphogenesis in the primate ovary. *Neuroscience* 138(3):773-81.
- Friedmann S et. al. (2005) Ovarian transcriptomes as a tool for a global approach of genes modulated by gonadotropic hormones in human ovarian granulosa cells. *Endocrine* 26(3):359-65.
- Hahn KL et. al. (2005) Lunatic fringe null female mice are infertile due to defects in meiotic maturation. *Development* 132(4):817-28.
- Hunter MG et. al. (2004) Endocrine and paracrine control of follicular development and ovulation rate in farm species. *Anim Reprod Sci* 82-83:461-477.
- Juengel JL et. al. (2004) Physiology of GDF9 and BMP15 signalling molecules. *Anim Reprod Sci* 82-83:447-460.
- Gilchrist RB et. al. (2004) Oocyte-somatic cell interactions during follicle development in mammals. *Anim Reprod Sci* 82-83:431-446.
- Zeleznik AJ (2004) The physiology of follicle selection. *Reprod Biol Endocrinol* 2:31.
- Sakumoto R, Okuda K (2004) Possible actions of tumor necrosis factor-alpha in ovarian function. *J. Reprod Dev* 50:39-46.
- Wei AY and Prints EA (2003) Therapy for polycystic ovarian syndrome. *Curr Opin Pharmacol* 3:678-682.
- Albertini DF, Barrett SL (2003) Oocyte-somatic cell communication. *Reprod Suppl* 61:49-54.
- Wong AS, Auersperg N (2003) Ovarian surface epithelium: family history and early events in ovarian cancer. *Reprod Biol Endocrinol* 1:70.
- Goodarzi MO, Korenman SG (2003) The importance of insulin resistance in polycystic ovary syndrome. *Fertil Steril* 80:255-258.
- Azziz R (2003) Androgen excess is the key element in polycystic ovary syndrome. *Fertil Steril* 80:252-254.
- Epifano O and Dean J (2002) Genetic control of early folliculogenesis in mice. *Trends in Endocrinology & Metabolism* 13:169-173.
- Kezele P, Nilsson E and Skinner MK (2002) Cell-cell interactions in primordial follicle assembly and development. *Frontiers in Bioscience* 7:d1990-1996.
- Richards JS et. al. (2002) Novel signaling pathways that control ovarian follicular development, ovulation, and luteinization. *Recent Prog Horm Res* 57:195-220.
- Findlay JK. et.al. (2002) Recruitment and development of the follicle; the roles of the transforming growth factor- $\beta$  superfamily. *Molecular and cellular endocrinology* 191:35-43.
- Matzuk MM, Burns KH, Viveiros MM and Eppig JJ (2002) Intercellular communication in the mammalian ovary: oocytes carry the conversation. *Science* 296:2178-2180.
- Vanderhyden B. (2002) Molecular basis of ovarian development and function. *Frontiers in Bioscience* 7:d2006-2022.
- Durlinger AL, Visser JA, Themmen AP (2002) Regulation of ovarian function: the role of anti-Müllerian hormone. *Reproduction* 124:601-609.
- Johnson AL, and Bridgham JT (2002) Caspase-mediated apoptosis in the vertebrate ovary. *Reproduction* 124:19-27.

# OVARY

## Ovary, Overview

Janice M Bahr, University of Illinois, Urbana-Champaign, IL, United States

© 2018 Elsevier Inc. All rights reserved.

### Glossary

**Corpus luteum** An endocrine gland formed from the granulosa and theca layers of an ovulated follicle.

**Follicle** A structure in the ovary consisting of the oocyte and surrounding granulosa and theca cell layers.

**Granulosa cells** Somatic cells directly surrounding the oocyte.

**Meiosis** A type of cell division which the oocyte undergoes reducing the number of chromosomes so that the oocyte has one copy of each chromosome.

**Oocyte** The female gamete.

**Ovary** The female gonad.

**Steroids** Molecules with a basic structure similar to that of cholesterol.

**Theca cells** Layer of steroidogenic cells and connective tissue surrounding the granulosa cells and forming the outer layer of the follicle.


## Introduction

Ovaries are female gonads responsible for the generation of female gametes (oocytes) and synthesis of hormones necessary for the regulation of reproductive functions. Since the first description of the ovary reported by Aristotle more than 2000 years ago, information about the ovary has expanded significantly. Knowledge of the formation of the ovary and its endocrine function is essential to understand the mystery of the regeneration of life.

## Anatomy of the Ovary

Most vertebrates develop a pair of ovaries with the exception of some birds, reptiles and a few mammals that only have one ovary. Ovaries lie on either side of the upper pelvic cavity and against the pelvic wall. They are held in place by a mesentery (mesovarium) connected to a broad ligament. Ovaries are one of the most vascular organs in the body. The ovarian artery (or utero-ovarian artery) which arises from the abdominal aorta reaches the ovary along with the mesovarium. Branches of the ovarian artery enter the ovary through the hilus, the same site at which the venous blood exits. Adrenergic and cholinergic nerves also enter the ovary through the hilus.

Even though the size of the ovary varies, the structure of the ovary is similar among mammalian species (**Fig. 1**). The ovary consists of an inner medulla, containing a rich vascular bed within loose connective tissue and an outer cortex, where the ovarian follicles are located. The outermost layer of the cortex is a single squamous or cuboidal surface epithelium derived from the peritoneum. Under the surface epithelium lies the tunica albuginea, a poorly delineated layer of dense connective tissue that gives the ovary a whitish color. The cortex of the ovary is made up of numerous follicles of various sizes and stages of development embedded in the stroma. The stroma is composed of at least three different cell types: connective tissue cells (fibroblasts) performing support functions, smooth muscle cells regulating the contraction of blood vessels and interstitial cells including undifferentiated theca cells and degenerated cells from atretic follicles and regressed corpora lutea. The follicles (follicle is Latin for "little bag") are structurally very conspicuous because of their variation in size. The microscopic appearance of follicles is different depending on the stage of follicular development whereas the basic cellular organization of follicles is the same. A follicle consists of an oocyte and surrounding follicular wall. Between the oocyte and surrounding follicular wall is a thin transparent membrane, the zona pellucida. The follicular wall contains an inner granulosa layer and an outer theca layer. The granulosa layer surrounds the oocyte and is separated from the theca layer by the basement membrane. The number and function of the granulosa cells changes during follicular growth. In mature follicles, the theca layer can be divided into the theca externa and interna. The theca externa consists of concentrically arranged smooth muscle cells innervated with autonomic nerves. The theca interna has epithelioid cells called interstitial cells, which are steroid producing cells. These cells contain LH and insulin receptors



**Fig. 1** A cross-section of the ovary illustrating follicles at different stages of development (from primordial to Graafian follicles), corpus hemorrhagicum, corpus luteum, and corpus albicans. The microscopic structures of follicles are also shown. Adapted from Jones, R.E. (1991) *The ovaries in human reproductive biology*, pp. 39–53. Academic Press, San Diego. p. 42.

and synthesize primarily androgens, of which the predominant steroid is androstenedione. The theca interna has both blood vessels and nerves. The granulosa layer is devoid of blood vessels and nerves at all times.

Once ovulation of the Graafian follicle (tertiary) has occurred, blood derived from torn blood vessels of the theca layer infiltrates the collapsed follicle and results in the formation of the corpus hemorrhagicum, a developing corpus luteum with a bloody core. Luteinizing granulosa and thecal cells begin to divide and invade the antral cavity, which remains after ovulation of the oocyte, forming the corpus luteum (Latin for “yellow body”). Blood vessels from the theca layer grow and penetrate the developing luteal cell mass. If pregnancy does not occur, the corpus luteum degenerates after a certain length of time depending upon the species. The connective tissues replaces the luteal cells and forms the corpus albicans (Latin for “white body”). The ovarian medulla devoid of follicles, contains large, spirally arranged blood vessels, lymphatic vessels and nerves.

## Functions of the Ovary

### Generation of the Female Gametes

#### Oogenesis

Female gametes, or oocytes, provide the maternal genetic material for the formation of an embryo. The ovary nurtures thousands of oocytes and functions as an incubator for their development. The development of oocytes (oogenesis) starts with primordial germ cells, residing in sex cords which divide mitotically producing oogonia. Oogonia then become primary oocytes and undergo the first meiosis. The primary oocytes are arrested at the diplotene stage of the first meiosis until they experience the preovulatory LH surge. Then the first meiosis is reinitiated and the membrane of the oocyte nucleus (germinal vesicle) disintegrates, which is called germinal vesicle breakdown. Meiosis of the oocyte is unequal producing a large haploid secondary oocyte and a tiny haploid first polar body. This polar body can divide again or remain single; in either case, it degenerates. Then the secondary oocyte begins the second meiotic division but this division is arrested at metaphase until after sperm penetration of the oocyte, which occurs in the oviduct. Completion of the second meiosis results in a haploid ovum and the second polar body.

#### Folliculogenesis

Folliculogenesis is a developmental sequence regulated by a number of genes, transcription factors and hormones. During fetal development of humans and postnatal development of mice, oocytes are present in clusters or nests. Majority of these oocytes enter meiosis during embryonic life. As the oocytes separate into individual oocytes, they form primordial follicles and undergo further development called oogenesis. Maturation of oocytes (oogenesis) is closely associated with the development of follicles because factors produced by the oocytes have a major impact on the development of the granulosa and theca layers. Folliculogenesis always begins in the innermost part of the ovarian cortex in mammals. Primordial follicles consist of primary oocytes surrounded by flat squamous pre-granulosa cells. Primordial follicles are the only available source of oocytes during the entire reproductive period of the female. As primordial follicles develop into primary follicles, there are changes in the oocyte. It

increases in diameter and develops an extracellular matrix, the zona pellucida. Reactivation of the oocyte genome causes the oocyte to secrete growth factors which play a crucial role in the growth of the follicle. As primary follicles grow, the granulosa cells divide mitotically so that secondary follicles have two to six layers of cuboidal-shaped granulosa cells. Secondary follicles also acquire an additional somatic cell layer, the theca. There are at least two sources of the theca progenitor cells: somatic precursors of the fetal ovary and mesenchymal cells in the neighboring mesonephros. The formation of this theca layer is dependent upon the presence of growth differentiation factor-9 (GDF-9) produced by the oocyte. The theca layer forms around the basement membrane in secondary follicles and ultimately forms the theca interna and theca externa. Follicular growth from primordial to secondary follicles is gonadotrophin-independent. During the formation of tertiary follicles or preantral follicles, follicles continue to grow in size. As follicles progress from secondary follicles to antral follicles, granulosa cells secrete a fluid that accumulates between cells. Large amounts of additional fluid diffuse out of the thecal blood vessels and are added to the fluid which is called follicular fluid. Follicular fluid contains steroid and protein hormones, anticoagulants, enzymes, and electrolytes and is similar to blood serum in appearance and contents. The follicle filled with follicular fluid is the tertiary or preovulatory follicle. These follicles have a mural granulosa layer of four to six layers and the theca layer is differentiated into an inner theca interna and an outer theca externa. Oocytes in preovulatory follicles are suspended in follicular fluid by a stalk of granulosa cells, the cumulus oophorus. Immediately surrounding the oocyte is a thin ring of granulosa cells, the corona radiata. At this state the follicle is called the Graafian follicle and appears as a transparent vesicle that bulges from the surface of the ovary.

Even though one of the function of the ovary is to produce oocytes, the majority of oocytes never ovulate. The number of oocytes reaches its maximum soon after the ovaries are formed. After that time oocyte number decreases dramatically. At birth, a female has all the oocytes she will have in her life; no new oocytes are formed after birth. The vast majority of oocytes, enclosed in follicles, around 99.9%, are eliminated before ovulation through a process called atresia which is due to the activation of apoptosis in the oocyte and granulosa cells. Follicles can become atretic at any stage of development.

### Production of Hormones

Another function of the ovary is to secrete hormones which act on the hypothalamus and pituitary to regulate the secretion of hormones by these two tissues, thus establishing the hypothalamic-pituitary-ovarian axis. The ovarian hormones also regulate the function of the reproductive tract and ultimately reproduction.


### Protein and peptide hormones

- (i) *Inhibin and activin*: Inhibin and activin were first isolated from gonadal fluids because of their effects on production of follicle stimulating hormone (FSH) by the pituitary in mammals. Inhibins consist of two disulfide-bridged subunits, the  $\alpha$  and  $\beta$  subunits, whereas activins consist of two  $\beta$  subunits. The primary source of inhibin and activin in the ovary is the maturing follicles and the corpus luteum. The function of inhibins is to modulate FSH secretion at the level of the pituitary, whereas the function of activins is to increase FSH secretion at the level of the pituitary. Inhibins and activins have antagonistic actions. Inhibins and activins also function as intraovarian hormones.
- (ii) *Follistatin*: Follistatin is a FSH-modulating polypeptide not related to TGF- $\beta$ . Follistatin acts as a binding protein and a functional antagonist of activin. Granulosa cells in antral follicles and luteal cells secrete follistatin.
- (iii) *Relaxin*: Relaxin is produced by the corpus luteum. The structure of relaxin is very similar to that of insulin but has <20% amino acid homology. In the human, relaxin is the highest during the first trimester of pregnancy after which the concentrations are relative stable. In the rat and the pig, relaxin reaches the highest concentration prior to parturition. Relaxin in these species functions to soften the cervix and vagina for the passage of the fetus during parturition and to promote the growth of nipples. Relaxin also acts on nonreproductive tissues, such as skin and the gastrointestinal tract.
- (iv) *Growth factors*: The ovary not only secretes endocrine hormones to regulate functions of other reproductive organs but also produces growth factors to coordinate the activities of different ovarian compartments. Many growth factors, such as insulin-like growth factors, transforming growth factors and epidermal growth factor are produced by the oocyte and somatic cells in the ovary. This complex intraovarian regulation system is no less important than the extraovarian regulation by the pituitary hormones. These growth factors form a delicate interactive communication web inside the ovary. Without them, the ovarian cells cannot interact with each other and the growth of the ovary is halted.

### Steroid hormones

The ovary uses cholesterol as the precursor for steroid synthesis. Cholesterol is metabolized into progestins, androgens, and estrogens by different compartments of the follicles (Fig. 2).

- (i) *Progestins pregnenolone*. Is the most important progestins (C21 pregnane family) produced by follicles because of its key position as the precursor of all steroid hormones. The most abundant progestin is progesterone, produced as a biosynthetic intermediate by follicles at all growing stages of development and as a secretory end product of the corpus luteum. In the developing follicles, the theca layer is the primary site of progestin productions. Immediately prior to ovulation, the granulosa cells stimulated by LH also synthesize progesterone. After ovulation the corpus luteum



**Fig. 2** Biosynthesis of steroid hormones from cholesterol. This scheme provides a simplistic view of a highly organized and complicated process that requires multiple enzymes. Adapted from Hafez, E.S.E (1993) Folliculogenesis, egg maturation, and ovulation. In *Reproduction in farm animals*, 6th ed., pp. 114–143. Lea and Febiger, Philadelphia, p. 79.


synthesizes copious amounts of progesterone needed to prepare the uterus for implantation and later for the maintenance of pregnancy.

- (ii) *Androgens*. The follicle is a significant source of ovarian androgens (C19 androstane family). Pregnenolone and progesterone are converted into androgen metabolites, dehydroepiandrosterone and androstenedione, respectively. These two metabolites are then transformed into testosterone. The theca layer of the follicle is the primary source of ovarian androgens.
- (iii) *Estrogens*. Physiologically, the estrogens (C18 estrane family) especially estrone and estradiol-17- $\beta$ , are the most important of the ovarian steroids. Androstenedione and testosterone are the immediate biosynthetic precursors of estrone and estradiol-17- $\beta$ , respectively. Their names reflect their roles in the induction of sexual receptivity (estrus) in female mammals. Estrone was the first sex steroid isolated and identified. The granulosa layer is the major site of estrogen synthesis in the mammalian ovary.

## Regulation of Ovarian Functions

### Regulation of Folliculogenesis

Growth of primordial follicles to the preantral stage is independent of gonadotropins and is controlled by intraovarian growth factors. Growth of follicles after the preantral stage depends on appropriate patterns of secretion, sufficient concentrations and adequate ratios of FSH and LH in the blood. FSH plays a major role in early follicular development. FSH stimulates granulosa cell mitosis and accumulation of follicular fluid. Granulosa cells synthesize estrogens in response to FSH which further enhance the mitotic effect of FSH. Moreover, FSH induces granulosa cell sensitivity to LH by increasing LH receptor expression. Abundant LH receptors in granulosa cells prepare the luteinization of granulosa cells in response to the ovulatory LH surge in mammals. In contrast, theca cells are stimulated only by LH and LH receptors are present from the beginning of the formation of the theca layer.



**Fig. 3** “Two-cell, two-hormone” theory of follicular steroidogenesis. LH binds to specific membrane receptors on theca cells and stimulates cyclic AMP production and the conversion of cholesterol to androgens, primarily androstenedione and testosterone. These androgens diffuse into the circulation and across the basement membrane into granulosa cells. FSH binds to specific membrane receptors on granulosa cells and stimulates cyclic AMP production, which leads to increased aromatase enzyme activity and the conversion of theca androgens to estrogens. Adapted from Yen, S. S. C. and Jaffe, R. B. (1986). *Reproductive Endocrinology* (2<sup>nd</sup> ed.), Philadelphia: Saunders.

### Regulation of Steroidogenesis

The steroidogenic output of the ovary is a function of coordinated actions of theca and granulosa cells. Differences in gonadotropin receptors on the membrane, in the activity of steroidogenic enzymes and in compartmentalization in the follicle result in a unique partnership in steroid synthesis between theca and granulosa cells. The principal site of estrogen synthesis in the mammalian ovary is granulosa cells under the control of FSH. Androgen production appears to be the primary steroidogenic function of theca cells in response to LH. Androgens from theca cells provide substrates for granulosa cells to synthesize estrogens. The action of LH on theca androgen production, together with the action of FSH on granulosa estrogen synthesis, forms the basis of the “two-cell, two-hormone” theory for the control of steroidogenesis in the ovary (Fig. 3).

### Further Reading

- Dong, J., Albertini, D. F., Nishimori, K., Kumar, T. R., Lu, N., & Matzuk, M. M. (1996). Growth differentiation factor-9 is required during early ovarian folliculogenesis. *Nature*, 383, 531–535.
- Hafez, E. S. E. (1993). Folliculogenesis, egg maturation, and ovulation. In *Reproduction in farm animals* (6th ed., pp. 114–143). Philadelphia: Lea and Febiger.
- Jones, R. E. (1991). *The ovaries in human reproductive biology*. San Diego: Academic Press. pp. 39–53.
- Liu, C., Peng, J., Matzuk, M.M. and Yao, H.H. (2015). *Nature Communications*. <https://doi.org/10.1038/ncomms7934>.
- Rajkovic, A., Panagas, S. A., & Matzuk, M. M. (2006). Follicular development: Mouse, sheep and human models. In J. D. Neill (Ed.), *Knobil and Neill's physiology of reproduction* (3rd edn). Amsterdam: Elsevier.
- Strauss, J.F. III and Williams, C.J. (2009). *The ovarian life cycle*, Strauss, J.F. III, Barberi, R.L. eds., Yen and Jaffe's reproductive endocrinology, physiology, pathophysiology, and management, 6th edn, Saunders, Philadelphia, PA.
- Williams, C. J., & Erickson, G. F. (2012). In L. J. De Groot, G. Chrousos, & K. Dungan (Eds.), *Morphology and physiology of the ovary*. South Dartmouth, MA: Endotext.

**"Systems Biology of Reproduction"**

Spring 2024 (Even Years) – Course Syllabus  
 Biol 475/575 Undergraduate/Graduate (3 Credit)  
 SLN: (475) – 06763, (575) – 06764  
 Time - Tuesday and Thursday 10:35 am-11:50 am  
 Course Lectures in person and recorded on Canvas/Panopto and Discussion Sessions live in person and on WSU Zoom for all campuses (Hybrid Course)  
 Room – CUE 418

Course Director – Michael Skinner, Abelson Hall 507, 335-1524, [skinner@wsu.edu](mailto:skinner@wsu.edu)  
 Co-Instructor – Eric Nilsson, Abelson Hall 507, 225-1835, [nilsson@wsu.edu](mailto:nilsson@wsu.edu)

**Learning Objective -**  
 Current literature based course on the Systems Biology of Reproduction. Learning Systems approaches to the biology of reproduction from a molecular to physiological level of understanding.

**Schedule/Lecture Outline –**

January 9 & 11	Week 1	Systems Biology Introduction
16 & 18	Week 2	Molecular/ Cellular/ Reproduction Systems
23 & 25	Week 3	Sex Determination Systems
Jan /Feb 30 & 1	Week 4	Male Reproductive Tract Development & Function
February 6 & 8	Week 5	Female Reproductive Tract Development & Function
13 & 15	Week 6	Gonadal Developmental Systems Biology
20 & 22	Week 7	Testis Systems Biology
27 & 29	Week 8	Ovary Systems Biology
March 5 & 7	Week 9	Epigenetics and Transgenerational Gonadal Disease
11 – 15	<b>Week 10</b>	<b>Spring Break</b>
19 & 21	Week 11	Gametogenesis/ Stem Cells/ Cloning
26 & 28	Week 12	Hypothalamus- Pituitary Development & Function
April 2 & 4	Week 13	Reproductive Endocrinology Systems
9 & 11	Week 14	Fertilization & Implantation Systems
16 & 18	Week 15	Fetal Development & Birth Systems
23 & 25	Week 16	Assisted Reproduction/Contraception
Apr/May 30 & 2	Week 17	Exam or Grant Review

Spring 2024 – Systems Biology of Reproduction  
 Lecture Outline – Ovarian Systems Biology  
 Michael K. Skinner – Biol 475/575  
 CUE 418, 10:35-11:50 am, Tuesday & Thursday  
 February 27, 2024  
 Week 8

**Ovarian Systems Biology**

Cell Biology of the Ovary  
 -Cell types/organization  
 -Developmental stages (Folliculogenesis)  
 -Atresia/apoptosis  
 -Oogenesis

Regulation of Folliculogenesis  
 -Growth properties of ovarian follicles  
 -Local production and action of growth factors  
 -Growth regulations during development  
 -Primordial follicle transition

Endocrine Regulation of Tissue Function  
 -Gonadotropin actions (Pituitary/Gonadal Axis)  
 -Steroid production and action  
 -Two cell theory modifications  
 -Hormone actions during development

Cell-Cell Interactions  
 -Categorization of different cell-cell interactions in the ovary  
 -Growth factor regulation follicle development  
 -Oogenesis and systems biology

**Required Reading**

Bahr JM. (2018) Ovary, Overview. in: Encyclopedia of Reproduction 2nd Edition, Ed: MK Skinner. Elsevier. Vol 2: 3-7.

Spring 2024 – Systems Biology of Reproduction  
 Discussion Outline – Ovary Systems Biology  
 Michael K. Skinner – Biol 475/575  
 Zoom/CUE 418, 10:35-11:50 am, Tuesday & Thursday  
 February 29, 2024  
 Week 8

**Ovary Systems Biology**

**Primary Papers:**

1. Pla, et al. (2021) Hum Reprod. 18:36(3):756-770
2. Sagvekar, et al. (2019) Clinical Epigenetics 11:61
3. Nilsson, et al. (2010) PLoS ONE 7:e11637

**Discussion**

Student 8: Reference 1 above


- What approach and technology was used?
- What gene categories and networks were identified?
- What oocyte maturation and folliculogenesis insights were identified?

Student 9: Reference 2 above

- What are the technology used and objectives?
- What epigenetic regulation and gene network were identified?
- What insights are provided into the development of polycystic ovarian disease?

Student 10: Reference 3 above

- What is the experimental and systems approach?
- What new insights provided on primordial follicle development?
- What gene signaling networks were identified for primordial follicle development?














Figure 11-5. Developmental changes in the adult ovary during a complete menstrual cycle. (Reproduced with permission from Carr SA, Wilson JD. Disorders of the ovary and female reproductive tract. In: Wilson JD, Braunwald E, Isselbacher KJ, et al (eds). Harrison's Principles of Internal Medicine, 12th Ed. New York, N.Y., McGraw-Hill, 1991, pp 1778-1785.)







**Cell Death -**  
 1858 Virchow - discuss process active & passive  
 Process = necrosis & degenerative

1860' s-1870' s - Weigert & Cohnheim - Students Virchow  
 Describe further - physiological need

1885 - Walter Flemming - ovary model used  
 1st micrograph apoptosis  
 previously he described - chromatin & mitosis  
 termed cell death process - chromatolysis


- Apoptosis termed -1914 for programmed cell death
- Rediscovered in 1980' s
- Term not best

Apoptosis	Necrosis
Affects scattered individual cells	Affects tracts of contiguous cells
Chromatin and cytoplasmic condensation, cell shrinkage	Cell swelling and rupture of plasma membrane
May require mRNA and protein synthesis	Not dependent upon new mRNA or protein synthesis
Normal ATP level	Decreased ATP level
No inflammation	Elicits inflammatory responses
Endonuclease activation and internucleosomal DNA cleavage (ladder pattern)	Activation of nonspecific DNases with generalized DNA breakdown (smearing)


Method of identification	Function	
<i>bcl-2</i>	Overexpressed in B-cell lymphoma (found at junction of t(14;18) chromosomal translocation) Cociprecipitate with <i>Bcl-2</i> protein	Suppresses apoptosis Prevents <i>bcl-2</i> action and increases apoptosis
<i>BAX</i>	Homologous screening of cDNA library	Prevents apoptosis
<i>bcl-X</i>	Long form Short form	Increases apoptosis ???
<i>A1</i>	Differential screening (hemopoietic cells)	Prevents apoptosis
<i>MCL1</i>	Subtraction screening (early response gene in myeloid cells)	???
<i>ICE</i>	Process cloning of <i>ced-3</i> (interleukin 1 $\alpha$ -converting enzyme)	Promotes apoptosis
<i>ced-9</i>	Prevents apoptosis	
<i>BHRF1</i>	Epstein-Barr viral sequences	Prevents apoptosis
<i>LMWS-HL</i>	African swine fever virus gene	???

**Figure 2**

**Influences of pro-and anti-apoptotic effectors on death receptor-mediated apoptosis.** Apoptosis is controlled by several pro- (green) and anti- (red) apoptotic proteins. The balance of these proteins are important to ensure tissue homeostasis.




**Initiation of follicular atresia: gene networks during early atresia in pig ovaries.** Zhang J, Liu Y, Yao W, Li Q, Liu H, Pan Z. *Reproduction*. 2018 Jul;156(1):23-33.








Bubble chart of potential signaling pathways generated by DEGs. Pathway analysis was performed to associate the unique DEGs with pathways using the KEGG database. The size and color of each bubble represent number of DEGs in each pathway and *P* value respectively. DEG, differentially expressed genes.

**Activation of autophagy in early neonatal mice increases primordial follicle number and improves lifelong fertility.** Watanabe R, Sasaki S, Kimura N. *Biol Reprod*. 2019 Sep 30. [Epub ahead of print]



**Autophagy participates in cyst breakdown and primordial folliculogenesis by reducing reactive oxygen species levels in perinatal mouse ovaries.** Zhihan T, Xinyi M, Qingying L, Rufeif G, Yan Z, Xuemel C, Yanqing G, Yingxiang W, Junlin H. *J Cell Physiol*. 2019 May;234(5):6125-6135.





## Number of Oocytes During Stages of Early Folliculogenesis

	Proliferation	Assembly	Puberty
Rodent	75,000	27,000	10,000
Primate	6,800,000	1,000,000	700,000


Only 500 human follicles ovulate.

## Oogonial Stem Cell

Journal of Clinical Oncology, Vol 25, No 22 (August 1), 2007; pp. 3198-3204  
© 2007 American Society of Clinical Oncology  
DOI: 10.1200/JCO.2006.10.3028



**Bone Marrow Transplantation Generates Immature Oocytes and Rescues Long-Term Fertility in a Preclinical Mouse Model of Chemotherapy-Induced Premature Ovarian Failure**





Ho-Joon Lee, Kaisa Selesniemi, Yuichi Nikura, Teruko Nikura, Rachael Klein, David M. Dombkowski, Jonathan L. Tilly




## The controversial existence and functional potential of oogonial stem cells.

**Grieve KM, McLaughlin M, Dunlop CE, Telfer EE, Anderson RA. Maturitas. 2015 Nov;82(3):278-81.**





**Nuclear exclusion of SMAD2/3 in granulosa cells is associated with primordial follicle activation in the mouse ovary.**  
 Hardy K, Mora JM, Dunlop C, Carzaniga R, Franks S, Fenwick MA.  
*J Cell Sci.* 2018 Sep 7;131(17). pii: jcs218123.



**Kit-Ligand/Stem Cell Factor Induces Primordial Follicle Development and Initiates Folliculogenesis\***

JEFF A. FAIRCOTT and MICHAEL K. SKINNER


Center for Reproductive Biology, Department of Genetics and Cell Biology, Washington State University, Pullman, Washington 99163-4211

**ABSTRACT**  
 Initiation of folliculogenesis through the initiation of primordial follicle development is a complex process involving the interplay of the follicle and reproductive tracts of most mammalian species. The factors that initiate this critical process are largely unknown. The hypothesis tested in the current study was that kit ligand/stem cell factor (KL) promotes the initiation and progression of primordial follicle development in the ovary. Ovaries from 4-day-old rats were maintained in organ culture for 0 and 14 days and treated with either control, recombinant kit ligand (rKL), or granulosa cell factor (GCF) and KITL. Follicles in ovarian sections were counted and histologically classified as primordial (stage 0), early primary (stage 1), primary (stage 2), transitional (stage 3), or antral (stage 4). Fresh ovaries from 4-day-old rats contained 100% primordial follicles (stage 0) and 22% developing follicles (stages 1–4) per section. After 0 and 14 days in culture, sections from control ovaries contained approximately 4% and 56%, respectively, developing follicles (stage 1–4) per section due to spontaneous development of primordial follicles. Spontaneous primordial follicle development was completely blocked by

ACR-2, a c-kit antibody that blocks KL action. This observation suggests that endogenous KL is necessary for the initiation of follicle development in mice. After 14 days of KL treatment, ovaries from control ovaries contained 17% primordial follicles (stage 0) and 57% developing follicles (stage 1–4) per section demonstrating a 60% increase in the number of primordial follicles. Development of primordial follicles was blocked by ACR-2 suggesting that KL may in part initiate granulosa cell development after the initiation of primordial follicle development. Ovaries contained an average of 200 ± 10 follicles per section. The total number of follicles per section did not significantly vary between treatments suggesting that the effects of KL were not due to an alteration in follicle number. In contrast, KL appears to be one of the first factors identified to be involved in the progression of primordial follicle development. Results suggest that KL is necessary and sufficient to induce primordial follicle development and initiate folliculogenesis. (*Endocrinology* 146:4302–4311, 1999)


**LONG-TERM CULTURE**


**RAT OVARY ORGAN CULTURE**






**Somatic cells initiate primordial follicle activation and govern the development of dormant oocytes in mice.**





Zhang H, Risal S, Gorre N, Busayavalasa K, Li X, Shen Y, Bosbach B, Brännström M, Liu K.  
*Curr Biol.* 2014 Nov 3;24(21):2501-8.





FOXO1/3 depletion in granulosa cells alters follicle growth, death and regulation of pituitary FSH. Liu Z, et al. Mol Endocrinol. (2013) Feb;27(2):238-52.






## Candidate Factors for Primordial to Primary Follicle Transition


- KL (Stem Cell Factor)(Kit Ligand )
- bFGF (basic Fibroblast Growth Factor)
- LIF (Leukemia Inhibitory Factor)
- GDF-9 (Growth and Differentiation Factor 9)
- BMP-4 (Bone Morphogenic Protein 4)
- EGF (Epidermal Growth Factor)
- HGF (Hepatocyte Growth Factor)
- KGF (FGF-7)(Keratinocyte Growth Factor)
- IGF-1 (Insulin Like Growth Factor 1)
- VEGF (Vascular Endothelial Growth Factor)
- TNF $\alpha$  (Tumor Necrosis Factor)

## Current Model of Early Folliculogenesis




## Roles of Gremlin 1 and Gremlin 2 in regulating ovarian primordial to primary follicle transition.


Nilsson EE, Larsen G, Skinner MK.  
Reproduction. 2014 Jun;147(6):865-74.



## Neurotrophin NT3 promotes ovarian primordial to primary follicle transition.


Nilsson E, Dole G, Skinner MK.  
Reproduction. 2009 Oct;138(4):697-707.






### Lhx8 regulates primordial follicle activation and postnatal folliculogenesis.

Ren Y, Suzuki H, Jagarlamudi K, Golnoski K, McGuire M, Lopes R, Pachnis V, Rajkovic A  
 BMC Biol. 2015 Jun 16;13:39.




### Dynamic expression patterns of *Irx3* and *Irx5* during germline nest breakdown and primordial follicle formation promote follicle survival in mouse ovaries.

Fu A, Oberholzer SM, Bagheri-Fam S, et al.  
 PLoS Genet. 2018 Aug 2;14(8):e1007488.




### *Dazl* determines primordial follicle formation through the translational regulation of *Tex14*.


Rosario R, Crichton JH, Stewart HL, Childs AJ, Adams IR, Anderson RA.  
 FASEB J. 2019 Dec;33(12):14221-14233.




**ELAVL2-directed RNA regulatory network drives the formation of quiescent primordial follicles.**  
 Kato Y, Iwamori T, Ninomiya Y, Kohda T, Miyashita J, Sato M, Saga Y.  
 EMBO Rep. 2019 Dec 5;20(12):e48251.




**Sirtuins in female meiosis and in reproductive longevity**  
 Berta N Vazquez, Alejandro Vaquero, Karen Schindler  
 Mol Reprod Dev. 2020 Dec;87(12):1175-1187.




**Identification of a unique epigenetic profile in women with diminished ovarian reserve.**  
 Olsen KW, Castillo-Fernandez J, Chan AC, et al.  
 Fertil Steril. 2021 Mar;115(3):732-741.



**The Mammalian Ovary**






## Systems Biology Approach:

1. Comparative mRNA expression with several treatments.
2. Network analysis.
3. Unbiased literature analysis.

### Regulatory Growth Factors

Used Compound	Official Gene Symbol	Gene Title
bFGF	Fgf2	fibroblast growth factor 2
PDGF	Pdgfb	platelet-derived growth factor beta polypeptide
LIF	Lif	Leukemia inhibitory factor
KGF	Fgf7	fibroblast growth factor 7
BMP4	Bmp4	bone morphogenetic protein 4
AMH	Amh	anti-Mullerian hormone
KL	Kitlg	KIT ligand
GDNF	Gdnf	glial cell derived neurotrophic factor
NT3	Ntf3	neurotrophin 3


All compounds had effect on **Primordial to Primary Follicle** Transition



### Number of Genes and Pathways Overlapped


	#PW	AMH	FGF2	BMP4	GDNF	FGF7	KITLG	LIF	PDGFab	CTGF
# Genes		56	41	22	20	36	54	56	41	12
AMH	268		28	18	14	29	37	40	32	7
FGF2	248	10		13	11	19	27	29	17	7
BMP4	79	4	5		9	11	14	16	13	6
GDNF	148	14	7	3		9	9	11	11	3
FGF7	123	36	5	1	5		28	25	17	5
KITLG	271	8	5	3	1	2		39	24	8
LIF	349	7	18	13	4	3	18		30	9
PDGFab	275	18	22	3	14	5	7	10		6
CTGF	155	5	7	2	1	2	6	2	6	

## Gene Co-expression Network




Module	Size, # genes
turquoise *	194
blue *	182
brown *	158
yellow *	150
green	139
red *	112
black	99
pink	85
magenta	68
purple *	45
greenyellow	32
tan	29
salmon	28
cyan	22
midnightblue	20
lightcyan	20
* - chosen	

(Hierarchical Clustering over the module-module similarity matrix heatmap derived from a network.)



“Short” list of 55 most connected in network genes  
CTGF, connective tissue growth factor

## Organ culture experiments to test if CTGF regulates follicle transition:



## Intraovarian control of early folliculogenesis.


Hsueh AJ, Kawamura K, Cheng Y, Fauser BC.  
Endocr Rev. 2015 Feb;36(1):1-24.

**Table 1.** Intraovarian Paracrine Hormones Act Through RTKs, RSKs, GPCRs, Guanylyl Cyclase Receptor NPRB, and Integrins to Regulate Preantral Follicle Growth<sup>a</sup>

Ligands	Receptors
IGF-1, KGF, VEGF, FGF2, FGF10	RTKs
Activins, BMP6, AMH	RSKs (types I and II)
PACAP, VIP	GPCRs
CNP	Guanylyl cyclase (NPRB)
CCN2/CTGF	Integrins




Abbreviations: CTGF, connective tissue growth factor.  
<sup>a</sup> Diverse paracrine growth factors are secreted by granulosa cells; they act through several distinct intracellular signaling pathways to promote follicle development. IGF-1, KGF, VEGF, FGF2, and FGF10 act through their respective RTKs in granulosa cells to regulate follicle development. In contrast, activins, AMH, and BMP6 synthesized by granulosa cells act through type I and type II RSKs in granulosa cells to regulate follicle development. Also, both PACAP and VIP produced by granulosa cells increase cAMP production by granulosa cells to regulate follicular functions. CNP secreted by granulosa cells binds to the guanylyl cyclase NPRB to increase cGMP production and promote follicle development. In contrast, CCN2/CTGF, produced by granulosa cells in response to Hippo signaling disruption, interacts with membrane-bound integrins in granulosa cells to promote follicle growth.

**WNT signaling in pre-granulosa cells is required for ovarian folliculogenesis and female fertility**  
 Habara O, Logan CY, Kanai-Azuma M, Nusse R, Takase HM.  
*Development*. 2021 May 1;148(9):dev198846.







**Proposed role of WNT signaling in folliculogenesis.** WNT-responding pre-GCs produce *Wnt4*, *Wnt6* and *Wnt11* in primordial follicles and oocytes secrete the WNT agonist *RSPO2*. Activation of canonical WNT signaling in pre-GCs promotes their transition into GCs during primordial follicle activation (PFA). In primary follicles, GCs induce the withdrawal of oocytes from a dormant state, as reflected by the translocation of FOX3 from the nucleus to the cytoplasm.

## The Mammalian Ovary



Hypothetical diagram of RTEC induced earlier menopause. The red line represents normal ovarian dynamics during aging while the blue line represents ovarian dynamics in a premature menopause scenario.



MOLCELL 1991  
 Rapid Paper  
**Mesenchymal-epithelial cell interactions in the ovary: estrogen-induced theca cell steroidogenesis**  
 Andrew J. Roberts and Michael K. Skinner  
 Department of Pharmacology, Florida State University School of Medicine, Tallahassee, FL 32310-4401, U.S.A.  
 (Received 22 June 1990; accepted 22 June 1990)

Key words: Mesenchymal-epithelial cell interactions; Ovary; Theca and granulosa cell; Steroidogenesis; Gonadotropin endocrinology

**Summary**  
 The role of mesenchymal-epithelial cell interactions in the control of ovarian physiology was investigated. Theca cells are the mesenchymal (M) stromal (S) cells that surround the ovarian follicle and produce androgens in response to the gonadotropin releasing hormone (LH). Granulosa cells are the epithelial cells that form the follicle, support the developing ovocyte, and utilize androgens produced by theca cells as a substrate for the production of estrogens. Observations presented indicate that androgens produced by granulosa cells dramatically stimulate androgen production by theca cells. Estrogen was found to have greater stimulatory effects on theca cell androgen production than progesterone, and a combination of estrogen and progesterone results in a greater than additive response of the two hormones. Regulation of androgen production by estrogen provides a local feedback loop in the follicle that will significantly influence ovarian steroidogenesis. The mesenchymal (theca-granulosa cell) interactions provide evidence for the importance of mesenchymal (theca-granulosa cell) interactions in which steroid androgens can produce paracrine factors that modulate mesenchymal cell function and differentiation. The theca cell-granulosa cell interaction identified is proposed to be a critical mesenchymal-epithelial cell interaction for the control of ovarian physiology and the maturation phase of the follicle.




Fig. 4. Schematic of the proposed steroid-mediated interactions between ovarian theca cells and granulosa cells. The steroids involved include androgens (A), estrogens (E) and progesterins (P) with both positive (+) and negative (-) effects on steroidogenesis. The gonadotropins involved include luteinizing hormone (LH) and follicle stimulating hormone (FSH).

## “FSH Actions on Granulosa Cells”




TABLE 1. FSH-stimulated functional parameters in cultured granulosa cells

- I. Enhancement of steroidogenesis
  - a. Estrogen biosynthesis
    - Induction of aromatase
    - Progesterone and 20 $\alpha$ -OH-P biosynthesis
      - Induction of cholesterol side-chain cleavage enzymes and mitochondrial cytochrome P450 activity
      - Induction of 17 $\beta$ -hydroxysteroid dehydrogenase
      - Induction of 20 $\alpha$ -hydroxysteroid dehydrogenase
  - b. Induction of specific plasma membrane receptors
    - LH receptor formation
    - Prolactin (lactogenic) receptor formation
    - $\beta_2$ -adrenergic receptor formation (coupling)
    - Lipoprotein receptor formation
    - FSH receptor formation
    - EGF receptor formation
  3. Secretion of nonsteroidal cell products
    - a. Inhibin
    - b. Plasminogen activator
    - c. Prostaglandins
    - d. Proteoglycans (mucopolysaccharides)
- II. Stimulation of general cell functions
  - a. DNA synthesis
  - b. Protein synthesis
  - c. Glucose uptake and lactate formation
  - d. Cell roundup and aggregation
  - e. Gap junction formation
  - f. Microvilli formation
- III. Plasma membrane-related processes
  - a. Adenyl cyclase activation and cAMP formation
  - b. Formation of cAMP binding protein
  - c. Phosphodiesterase activation
  - d. Increases in plasma membrane microviscosity

## “Actions of LH on Granulosa Cells”

TABLE 3. LH-modulated functional parameters in cultured granulosa cells

- I. Enhancement of steroidogenesis
  - a. Estrogen biosynthesis
  - b. Progesterone and 20 $\alpha$ -OH-P biosynthesis
    - Stimulation of cholesterol esterase activity
    - Induction of cholesterol side-chain cleavage enzyme
    - Induction of 17 $\beta$ -hydroxysteroid dehydrogenase
- II. Modulation of plasma membrane receptors
  - a. Increases in prolactin (lactogenic) receptor formation
  - b. Increases in  $\beta_2$ -adrenergic receptor formation (coupling)
  - c. Down-regulation or stimulation of LH receptors
  - d. Increases in lipoprotein receptor formation
- III. Secretion of non-steroidal cell products
  - a. Plasminogen activator
  - b. Mucopolysaccharides
  - c. Prostaglandins
  - d. Relaxin
- IV. Modulation of general cell functions
  - a. Increases in protein synthesis
  - b. Increases in glycolysis
  - c. Alterations of cell shape
  - d. Reduction in size of gap junctions
  - e. Inhibition of DNA synthesis
  - f. Stimulation of ornithine decarboxylase activity
- V. Plasma membrane-related processes
  - a. Adenyl cyclase activation and cAMP formation
  - b. Stimulation of cAMP-dependent protein kinase activity
  - c. Stimulation of phospholipid turnover










TABLE I  
PROPERTIES AND NOMENCLATURE OF SEVERAL COMMON GROWTH FACTORS

Growth Factor	Approx. Size (kDa)	Examples of Physiological Action	
Insulin-like Growth Factor-I	IGF-I	7.5	Skeletal Growth
Insulin-like Growth Factor-II	IGF-II	7.5	Fetal Development
Epidermal Growth Factor	EGF	6	Tissue Growth
Transforming Growth Factor Alpha	TGF- $\alpha$	5	Tissue Growth
Transforming Growth Factor Beta	TGF- $\beta$	25kDa	Growth Inhibition/ Tissue Repair
Fibroblast Growth Factor	FGF	17	Angiogenesis/Tissue Growth
Vascular Endothelial Growth Factor	VEGF	25-50kDa	Angiogenesis/Tissue Growth
Nerve Growth Factor	NGF	13	Neuronal Development
Interleukin-1	IL-1	17	Immune Response/ Inflammation
Platelet Derived Growth Factor	PDGF	30kDa	Tissue Growth
Stem Cell Factor (c-kit ligand)	SCF	30	Tissue Growth/Fetal Development



### Transforming Growth-Beta Factor Family






FIG. 1. Clones of BMP families and nomenclature of BMPs with their TGF- $\beta$  superfamily. Based on sequence homology, different representative BMPs starting with DPP (close at the top) are listed. As the tree moves from top to bottom, the sequence homology between TGF- $\beta$  receptors and the remaining DPP family members decreases. The remaining DPP family members are listed with a 1 to 10 DPP family members and numbered from 1 to 10. The numbers in parentheses indicate the number of family members through which ALKs and SMADs, and overexpression of BMP11 results in neural crest-derived mesoderm (Shimazaki et al., 2004). Thus, they are, functionally and all sequence family, listed to Activin and TGF- $\beta$  superfamily.






Signalling pathways involved in the cooperative effects of ovine and murine GDF9+BMP15-stimulated thymidine uptake by rat granulosa cells.  
 Reader KL, et al.  
 Reproduction. (2011) Jul;142(1):123-31.



**The epidermal growth factor network: role in oocyte growth, maturation and developmental competence**

Hum Reprod Update. 2018 Jan 1;24(1):1-14.



Gene	Product	Function
Amphiregulin (Areg)	EGF-like factor	Binds EGFR and activates ERK1/2
A disintegrin and metalloproteinase with thrombospondin-like repeats (ADAMTS1)	Protein	Secreted protease cleaves versican
Cathepsin L (Ctsl)	Protein	Protease
Chondroitin sulfate proteoglycan (Cspg2)	Versican	Binds HA
Cell differentiation 14 (Cd14)	Adaptor molecule	Binds LPS
CD34 (Cd34)	Membrane protein	Immune cell stem cell marker
CD36 (Cd36)	Membrane protein	Scavenger receptor
CD52 (Cd52)	GPI anchored protein	Unknown
Complement factor Q1 (C1q)	Clq	Adaptor for TLR2/4 binds HA and PTK3
HA synthase 2 (Hase2)	HA	Polymer of matrix of COCs
Interleukin 6 (Il6)	IL-6, cytokine	Inflammation, innate immune
Pentraxin 3 (Ptx3)	Matrix protein	Binds TNFAIP6
Progesterone receptor (Pgr)	Nuclear protein	Transcription factor
Programmed cell death 1 (Pcdl1)	Membrane protein	Anti-autoimmune regulator
Prostaglandin synthase 2 (Pgs2)	PGE2, prostaglandin	Binds (prostaglandin E2 receptor subtypes) EP2 receptor and induces AREG
Rarb1	Nuclear protein	Transcription factor
Toll-like receptors (Tlr2, Tlr4)	Membrane proteins	Bind LPS and Pam3Cys and HA fragments
TNF- $\alpha$ -induced protein 6 (Tnfrsf6)	Matrix protein	Binds and stabilizes HA matrix

**KGF** - keratinocyte growth factor  
28kDA mesenchymal cell derived growth factor (FGF-7); receptor FGFR-2 splice variant  
Receptor only on epithelial type cells

**HGF** - hepatocyte growth factor  
28kDA mesenchymal cell derived growth factor (scatter factor); promote kidney tubulogenesis  
Receptor is c-met protooncogene  
only on epithelium

Both mediate mesenchymal-epithelial interactions

Theca cells- express/secret KGF & HGF  
Granulosa cells- respond  $\uparrow$  growth




FIG. 5. Effects of KGF and HGF on proliferation of theca and granulosa cells. TERT immortalized granulosa cells (GCs). Data shown is an average  $\pm$  SEM of three independent experiments. The numbers of different experiments is indicated at granulosa or theca, each treatment. Values for the control were generally greater than 100% of control DNA.

Parrott et al (1994)  
Endocrinology  
135:569

TABLE 2  
GROWTH FACTORS IN THE OVARY


Growth Factor	Proposed Site Synthesis	Proposed Site of Action	Proposed Function*
IGF-1	Granulosa	Granulosa Theca	+Growth $\rightarrow$ Differentiation +Growth $\rightarrow$ Differentiation
FGF	Granulosa	Granulosa Endothelium	+Growth Angiogenesis
TGF- $\beta$	Theca Granulosa	Granulosa Theca	-Growth $\rightarrow$ Differentiation -Growth $\rightarrow$ Differentiation
TGF- $\alpha$	Theca	Granulosa Theca	+Growth $\rightarrow$ Differentiation +Growth
VEGF	Granulosa	Endothelium	Angiogenesis $\rightarrow$ Growth
NGF	Ovary	Neuron	Innervation
SCF (كسوليا)	Granulosa	Oocyte	Oocyte Maturation




\*A (+) denotes an increase and (-) indicates a decrease.

KGF	Theca	Granulosa	+growth
HGF	Theca	Granulosa	+growth
GDF9	Oocyte	Granulosa/ Theca	+growth/ +differentiation
BMP15	Oocyte	Granulosa	+growth





### The role of Notch signaling in the mammalian ovary

Reproduction, 2017 Jun;153(6):R187-R204.  
Vanony DA, Mayo KE.







	<b>Environmental</b>	<b>Nutritional</b>	<b>Regulatory</b>
<b>Theca-Granulosa</b>	ECM	Androgen	Estrogen Progesterin TGFα TGFβ HGF KGF
<b>Granulosa-Oocyte</b>	Minimal ECM Cytoarchitectural Support	Gap Junctions (<800mw)	cAMP Xanthine Derivatives OMI Kit-ligand (SCF) GDF-9
<b>Granulosa-Granulosa</b>	Minimal ECM Gap Junctions	Gap Junctions (cAMP)	Estrogen IGF FRP



The dormant and the fully competent oocyte: comparing the transcriptome of human oocytes from primordial follicles and in metaphase II.  
Grøndahl ML, et al.  
Mol Hum Reprod. (2013) 19(9):600-17.




Transcriptome profile of goat folliculogenesis reveals the interaction of oocyte and granulosa cell in correlation with different fertility population.  
Li S, Wang J, Zhang H, et al.  
Sci Rep. 2021 Aug 3;11(1):15698.




Follicle Online: an integrated database of follicle assembly, development and ovulation.

Hua J, Xu B, Yang Y, Ban R, Iqbal F, Cooke HJ, Zhang Y, Shi Q.  
Database (Oxford). 2015 Apr 29;2015:bav036.




Cell-specific network analysis of human folliculogenesis reveals network rewiring in antral stage oocytes.



Wang S, Gong Y, Wang Z, et al.  
J Cell Mol Med. 2021 Mar;25(6):2851-2860.




**A single-cell atlas reveals unanticipated cell type complexity in Drosophila ovaries.**  
 Slaidina M, Gupta S, Banisch TU, Lehmann R.  
 Genome Res. 2021 Oct;31(10):1938-1951.




**Single-Cell Transcriptomics Analysis of Human Small Antral Follicles.**  
 Fan X, Moustakas I, Bialecka M, et al.  
 Int J Mol Sci. 2021 Nov 4;22(21):11955.



**Single-cell RNA-sequencing of retrieved human oocytes and eggs in clinical practice and for human ovarian cell atlasing.**  
 Machlin JH, Shikanov A.  
 Mol Reprod Dev. 2022 Dec;89(12):597-607.




Novel insights into reproductive ageing and menopause from genomics.  
 Das A, Destouni A.  
 Hum Reprod. 2023 Feb 1;38(2):195-203.




Ovarian and somatic biological ageing clocks and their putative relationship.

Making a good egg: human oocyte health, aging, and in vitro development.  
 Telfer EE, Grosbois J, Odey YL, Rosario R, Anderson RA.  
 Physiol Rev. 2023 Oct 1;103(4):2623-2677.




Single-cell analysis of the aged ovarian immune system reveals a shift towards adaptive immunity and attenuated cell function.  
 Ben Yaakov T, Wasserman T, Aknin E, Savir Y.  
 Elife. 2023 Apr 25;12:e74915.


**The effect of female age on the ovarian immune milieu**



**The effect of fertile female aging on the ovarian immune system**



Cellular hallmarks of aging emerge in the ovary prior to primordial follicle depletion.  
 Ansere VA, Ali-Mondal S, Sathiaselvan R, et al.  
 Mech Ageing Dev. 2021 Mar;194:111425.



**Polycystic ovary syndrome as a paradigm for prehypertension, prediabetes, and proobesity.**

Luque-Ramirez M, Escobar-Morreale HF. *Curr Hypertens Rep.* 2014 Dec;16(12):500.


**Table 1** Effect of obesity on cardiometabolic risk factors in women with PCOS compared with lean patients

- ↑ Insulin resistance
- ↑ Hyperinsulinism
- ↑ Fasting glucose
- ↑ 2 h glucose after OGTT
- ↑ Prevalence of abnormal glucose tolerance
- ↓ HDL-cholesterol levels
- ↑ Triglycerides levels
- ↑ Office and ABPM systolic and diastolic blood pressure values
- ↑ Prevalence of office and ABPM hypertension
- ↑ Prevalence of NAFLD
- ↑ Prevalence of metabolic syndrome

ABPM ambulatory blood pressure monitoring, NAFLD nonalcoholic fatty liver disease, OGTT standard 75 g oral glucose tolerance test

**Scientific Statement on the Diagnostic Criteria, Epidemiology, Pathophysiology, and Molecular Genetics of Polycystic Ovary Syndrome.**


Dumesic DA, Oberfield SE, Stener-Victorin E, Marshall JC, Laven JS, Legro RS. *Endocr Rev.* 2015 Oct;36(5):487-525.



Pathophysiology of PCOS—a vicious circle. Several theories have been proposed to explain the pathogenesis of PCOS. One of these is that neuroendocrine defects lead to increased pulse frequency and amplitude of LH and relatively low FSH. This causes intrinsic defects in ovarian androgen production. Also, there may be an alteration in cortisol metabolism and excessive adrenal androgen production. Insulin resistance with compensatory hyperinsulinemia further increases ovarian androgen production both directly and indirectly via the inhibition of hepatic SHBG production. Obesity, insulin resistance, and high circulating androgens are associated with increased sympathetic nerve activity. E, estradiol.

**Alteration in angiogenic potential of granulosa-lutein cells and follicular fluid contributes to luteal defects in polycystic ovary syndrome.**


Patil K, Hinduja I, Mukherjee S. *Hum Reprod.* 2021 Mar 18;36(4):1052-1064.



The concentration of vascular endothelial growth factor (VEGF) and fibroblast growth factor 2 (FGF2) levels in FF and serum of control and women with polycystic ovary syndrome (PCOS). FF (n ¼ 20 in each group) and serum (n ¼ 10 in each group) were used to measure levels of VEGF (A and B) and FGF2 (C and D), in women with PCOS compared to controls. Statistical comparison was performed using the Mann-Whitney U test. Data are represented as mean ± SEM and \*P < 0.05; \*\*P < 0.01 considered significant. FF, follicular fluid.

**Compromised Cumulus-Oocyte Complex Matrix Organization and Expansion in Women with PCOS**


Patil K, Shinde G, Hinduja I, Mukherjee S. *Reprod Sci.* 2021 Nov 8. Online ahead of print.



**Functional microarray analysis of differentially expressed genes in granulosa cells from women with polycystic ovary syndrome related to MAPK/ERK signaling.**


Lan CW, Chen MJ, Tai KY, Yu DC, Yang YC, Jan PS, Yang YS, Chen HF, Ho HN. *Sci Rep.* 2015 Oct 13;5:14994.

Genes Symbol	Genes Name	log fold change	adj P value
<b>Upregulated</b>			
AOX1	aldehyde oxidase 1	1.475	0.001813**
Chr154	chromosome 1 open reading frame 58	1.311	0.002377**
SNORD59A	small nuclear RNA, C/D box 59A	1.257	0.002768**
TRIM22	tripartite motif-containing 22	1.212	0.002481**
RN758	RNA, 758, small nuclear	1.180	0.001813**
ANKRD368	ankyrin repeat domain 368	1.157	0.001583**
ENPP5	ectonucleotide pyrophosphatase/cyphosphohydrolase 5	1.115	0.002160**
SLC25A30	solute carrier family 25, member 30	1.085	0.002599**
SNORD21	small nuclear RNA, C/D box 21	1.028	0.001583**
ALPK1	alpha-kinase 1	1.008	0.001583**
<b>Downregulated</b>			
DEK16	DEAF1 like polypeptide 16	-0.762	0.004345**
FRS35	F-box protein 5	-0.709	0.013218*
OR14C36	olfactory receptor, family 14, subfamily C, member 36	-0.627	0.007862**
PCO1H	prostate collagen triple helix	-0.613	0.001583**
RPL4	ribosomal protein L4	-0.563	0.001660**
BC1L15	BC1.2 like 15	-0.552	0.002969**
OR52B	olfactory receptor, family 52, subfamily E, member 8	-0.453	0.014957*
GTF2H1	general transcription factor IIH, polypeptide 1	-0.400	0.025264**
OR5A2	olfactory receptor, family 5, subfamily A, member 2	-0.429	0.011000*
PTH2B	parathyroid hormone 2 receptor	-0.403	0.011824*




**DNA methylome profiling of granulosa cells reveals altered methylation in genes regulating vital ovarian functions in polycystic ovary syndrome.**

Sagvekar P, Kumar P, Mangoli V, Desai S, Mukherjee S. *Clin Epigenetics.* 2019 Apr 11;11(1):61.



**Transcriptional landscape of mouse-aged ovaries reveals a unique set of non-coding RNAs associated with physiological and environmental ovarian dysfunctions.**


Cuomo D, Porreca I, Ceccarelli M, Threadgill DW, et al. *Cell Death Discov.* 2018 Dec 5;4:112.




**Networks of putative target genes regulated by differentially expressed miRNAs.** Validated target genes (grey nodes) mapped to pathway annotations derived from literature and gene ontology using Ingenuity Pathway Analysis (IPA; <http://www.ingenuity.com>). **A** *Mir143* regulatory network **B** and *Mir145* regulatory network. The solid lines connecting molecules represent a direct relation and dotted lines an indirect relation. IPA constructs networks that optimize for both interconnectivity and number of Focus Genes (the grey nodes) under the constraint of a maximal network size. White nodes are added by the algorithm to build a tightly connected molecular network between Focus Genes.

**WGCNA Analysis Identifies Polycystic Ovary Syndrome-Associated Circular RNAs That Interact with RNA-Binding Proteins and Sponge miRNAs.**

Li M, Zeng Z, Zhang A, et al. *Int J Gen Med.* 2021 Nov 23;14:8737-8751.




**The lncRNA-miRNA-mRNA ceRNA network in mural granulosa cells of patients with polycystic ovary syndrome: an analysis of Gene Expression Omnibus data.**  
 Chen H, Cheng S, Xiong W, Tan X.  
 Ann Transl Med. 2021 Jul;9(14):1156.



Flowchart of data processing and analysis. lncRNA, long non-coding RNA; miRNA, microRNA; mRNA, messenger RNA; GO, Gene Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes.

**Evaluation of the expression profile of mRNAs and lncRNAs in cumulus cells associated with polycystic ovary syndrome and pregnancy.**  
 Hammami B, Mostafavi FS, Akbari A, Mousavi SR, Kazemi M.  
 Iran J Basic Med Sci. 2023;26(10):1144-1154.




Microarray data analysis of 7 PCOS and 3 normal GCs samples


Hub gene network. Node color is related to its degree


lncRNA-mRNA interaction network obtained from cytoscape

**Methylome and transcriptome profiling revealed epigenetic silencing of LPCAT1 and PCYT1A associated with lipidome alterations in polycystic ovary syndrome.**  
 Mao Z, Li T, Zhao H, Wang X, Kang Y, Kang Y.  
 J Cell Physiol. 2021 Sep;236(9):6362-6375.



**Systems biology and in silico-based analysis of PCOS revealed the risk of metabolic disorders.**  
 Hossain MA, Al Ashik SA, Mahin MR, Al Amin M, Rahman MH, Khan MA, Emran AA.  
 Heliyon. 2022 Dec 22;8(12):e12480.





### "Systems Biology of Reproduction"

Spring 2024 (Even Years) – Course Syllabus  
 Biol 475/575 Undergraduate/Graduate (3 Credit)  
 SLN: (475) – 06763, (575) – 06764  
 Time - Tuesday and Thursday 10:35 am-11:50 am  
 Course Lectures in person and recorded on Canvas/Panopto and Discussion Sessions live in person and on WSU Zoom for all campuses (Hybrid Course)  
 Room – CUE 418  
 Course Director – Michael Skinner, Abelson Hall 507, 335-1524, [skinner@wsu.edu](mailto:skinner@wsu.edu)  
 Co-Instructor – Eric Nilsson, Abelson Hall 507, 225-1835, [nilsson@wsu.edu](mailto:nilsson@wsu.edu)

**Learning Objective -**  
 Current literature based course on the Systems Biology of Reproduction. Learning Systems approaches to the biology of reproduction from a molecular to physiological level of understanding.

**Schedule/Lecture Outline -**

January	9 & 11	Week 1	Systems Biology Introduction
	16 & 18	Week 2	Molecular/ Cellular/ Reproduction Systems
	23 & 25	Week 3	Sex Determination Systems
Jan /Feb	30 & 1	Week 4	Male Reproductive Tract Development & Function
February	6 & 8	Week 5	Female Reproductive Tract Development & Function
	13 & 15	Week 6	Gonadal Developmental Systems Biology
	20 & 22	Week 7	Testis Systems Biology
	27 & 29	Week 8	Ovary Systems Biology
March	5 & 7	Week 9	Epigenetics and Transgenerational Gonadal Disease
	11 – 15	Week 10	Spring Break
	19 & 21	Week 11	Gametogenesis/ Stem Cells/ Cloning
	26 & 28	Week 12	Hypothalamus- Pituitary Development & Function
April	2 & 4	Week 13	Reproductive Endocrinology Systems
	9 & 11	Week 14	Fertilization & Implantation Systems
	16 & 18	Week 15	Fetal Development & Birth Systems
	23 & 25	Week 16	Assisted Reproduction/Contraception
Apr/May	30 & 2	Week 17	Exam or Grant Review