

Spring 2026 – Systems Biology of Reproduction
Lecture Outline – Epigenetics and Transgenerational Reproductive Disease
Eric Nilsson – Biol 475/575
10:35-11:50 am, Tuesday & Thursday
March 10, 2026
Week 9

Epigenetics and Transgenerational Reproductive Disease

- Environmental Induced Pathology
- Transgenerational Disease Phenotype
- Compound Specificity
- Epigenetic Mechanism
- Epimutations and Exposure Specificity
- Transgenerational Transcriptome
- Transgenerational Testis Disease
- Transgenerational Ovary Disease
- Broader Impact

REQUIRED READING

Ben Maamar M, Nilsson EE, Skinner MK. Epigenetic transgenerational inheritance, gametogenesis and germline development. *Biol Reprod.* 2021 Sep 14;105(3):570-592.

Wang HD, Allard P. Challenging dogmas: How transgenerational epigenetics reshapes our views on life. *J Exp Zool A Ecol Integr Physiol.* 2022 Jan;337(1):70-74.

REFERENCES

- Luo L, Li J, Sun Y, Lv Y, Liu J, Li Y, Zhang C, Zhang W. Maternal genetic intergenerational and transgenerational effects on hormone synthesis in ovarian granulosa cells of offspring exposed to cadmium during pregnancy. *Ecotoxicol Environ Saf.* 2023 Sep 15;263:115278
- Zhang Y, Li J, Shi W, Lu L, et al. Di(2-ethylhexyl) phthalate induces reproductive toxicity and transgenerational reproductive aging in *Caenorhabditis elegans*. *Environ Pollut.* 2023 Nov 1;336:122259.
- Sun Y, Zhang C, Luo L, Lin H, et al. Paternal genetic intergenerational and transgenerational effects of cadmium exposure on hormone synthesis disorders in progeny ovarian granulosa cells. *Environ Pollut.* 2023 Apr 1;322:121175
- Moody SN, Phan JM, Shirtcliff EA, et al. Transgenerational effect of Mothers' experiences of discrimination on Black youths' hormone coupling in response to laboratory stress. *Res Hum Dev.* 2023;20(1-2):25-47.

- Tajai P, Pruksakorn D, Chattipakorn SC, et al. Effects of glyphosate-based herbicides and glyphosate exposure on sex hormones and the reproductive system: From epidemiological evidence to mechanistic insights. *Environ Toxicol Pharmacol*. 2023 Sep;102:104252.
- Dutta S, Sengupta P, Bagchi S, et al. Reproductive toxicity of combined effects of endocrine disruptors on human reproduction. *Front Cell Dev Biol*. 2023 May 12;11:1162015.
- Souza GS, Freitas IMM, Souza JC, et al. Transgenerational effects of maternal exposure to nicotine on structures of pituitary-gonadal axis of rats. *Toxicol Appl Pharmacol*. 2023 Jun 1;468:116525.
- Lopes MFDS, Felix JS, Scaramele NF, et al. Co-expression analysis of lncRNA and mRNA identifies potential adipogenesis regulatory non-coding RNAs involved in the transgenerational effects of tributyltin. *PLoS One*. 2023 Feb 3;18(2):e0281240.
- Yin J, Hong X, Wang J, Li W, et al. DNA methylation 6 mA and histone methylation involved in multi-/trans-generational reproductive effects in *Caenorhabditis elegans* induced by Atrazine. *Ecotoxicol Environ Saf*. 2023 Jan 1;249:114348.
- Devoy C, Raza Y, Kleiner M, et al. The brominated flame retardant, 1,2,5,6-tetrabromocyclooctane (TBCO), causes multigenerational effects on reproductive capacity of Japanese medaka (*Oryzias latipes*). *Chemosphere*. 2023 Feb;313:137561.
- Lim J, Shioda T, Malott KF, et al. Prenatal exposure to benzo[a]pyrene depletes ovarian reserve and masculinizes embryonic ovarian germ cell transcriptome transgenerationally. *Sci Rep*. 2023 May 29;13(1):8671. doi: 10.1038/s41598-023-35494-w.
- Huang W, Shi X, Zhang Q, et al. Transgenerational effects of BDE-47 to zebrafish based on histomorphometry and toxicogenomic analyses. *Chemosphere*. 2023 Dec;344:140401.
- Zhou Y, Fu J, Wang M, et al. Parental and transgenerational impairments of thyroid endocrine system in zebrafish by 2,4,6-tribromophenol. *J Environ Sci (China)*. 2023 Feb;124:291-299.
- Kumar R, Fatima F, Yadav G, et al. Epigenetic Modifications by Estrogen and Androgen in Alzheimer's Disease. *CNS Neurol Disord Drug Targets*. 2023;22(1):6-17.
- Weller JI, Ezra E, Gershoni M. Broad phenotypic impact of the effects of transgenerational heat stress in dairy cattle: a study of four consecutive generations. *Genet Sel Evol*. 2021 Sep 6;53(1):69.
- Lim AI, McFadden T, Link VM, et al. Prenatal maternal infection promotes tissue-specific immunity and inflammation in offspring. *Science*. 2021 Aug 27;373(6558):eabf3002.
- Fellous A, Wegner KM, John U, et al. Windows of opportunity: Ocean warming shapes temperature-sensitive epigenetic reprogramming and gene expression across gametogenesis and embryogenesis in marine stickleback. *Glob Chang Biol*. 2022 Jan;28(1):54-71.
- Martin M, Hill C, Bewley S, et al. Transgenerational adverse effects of valproate? A patient report from 90 affected families. *Birth Defects Res*. 2021 Dec 5. doi: 10.1002/bdr2.1967. Online ahead of print.
- Grishok A. Small RNAs Worm Up Transgenerational Epigenetics Research. *DNA (Basel)*. 2021 Dec;1(2):37-48.
- Anastasiadi D, Venney CJ, Bernatchez L, Wellenreuther M. Epigenetic inheritance and reproductive mode in plants and animals. *Trends Ecol Evol*. 2021 Dec;36(12):1124-1140.
- Parker J, O'Brien C, Gersh FL. Developmental origins and transgenerational inheritance of polycystic ovary syndrome. *Aust N Z J Obstet Gynaecol*. 2021 Dec;61(6):922-926.
- Özdemir I, Steiner FA. Transmission of chromatin states across generations in *C. elegans*. *Semin Cell Dev Biol*. 2021 Nov 22;S1084-9521(21)00284-6.
- Ghai M, Kader F. A Review on Epigenetic Inheritance of Experiences in Humans. *Biochem Genet*. 2021 Nov 18. doi: 10.1007/s10528-021-10155-7. Online ahead of print.

- Zatecka E, Bohuslavova R, Valaskova F, et al. The Transgenerational Transmission of the Paternal Type 2 Diabetes-Induced Subfertility Phenotype. *Front Endocrinol (Lausanne)*. 2021 Nov 5;12:763863.
- Alvarez M, Bleich A, Donohue K. Genetic differences in the temporal and environmental stability of transgenerational environmental effects. *Evolution*. 2021 Nov;75(11):2773-2790.
- Vilcinskis A. Mechanisms of transgenerational immune priming in insects. *Dev Comp Immunol*. 2021 Nov;124:104205.
- Šrut M. Ecotoxicological epigenetics in invertebrates: Emerging tool for the evaluation of present and past pollution burden. *Chemosphere*. 2021 Nov;282:131026.
- Lock M, Argentieri MA, Shields AE. The contribution of ethnography to epigenomics research: toward a new bio-ethnography for addressing health disparities. *Epigenomics*. 2021 Nov;13(21):1771-1786.
- Skinner MK, Nilsson EE. Role of environmentally induced epigenetic transgenerational inheritance in evolutionary biology: Unified Evolution Theory. *Environ Epigenet*. 2021 Oct 30;7(1):dvab012.
- Zheng X, Li Z, Wang G, et al. Sperm epigenetic alterations contribute to inter- and transgenerational effects of paternal exposure to long-term psychological stress via evading offspring embryonic reprogramming. *Cell Discov*. 2021 Oct 27;7(1):101.
- Wellband K, Roth D, Linnansaari T, Curry RA, Bernatchez L. Environment-driven reprogramming of gamete DNA methylation occurs during maturation and is transmitted intergenerationally in Atlantic Salmon. *G3 (Bethesda)*. 2021 Dec 8;11(12):jkab353.
- Gowri V, Monteiro A. Inheritance of Acquired Traits in Insects and Other Animals and the Epigenetic Mechanisms That Break the Weismann Barrier. *J Dev Biol*. 2021 Oct 7;9(4):41.
- Ben Maamar M, Beck D, Nilsson EE, Kubsad D, Skinner MK. Epigenome-wide association study for glyphosate induced transgenerational sperm DNA methylation and histone retention epigenetic biomarkers for disease. *Epigenetics*. 2021 Oct;16(10):1150-1167.
- Conine CC, Rando OJ. Soma-to-germline RNA communication. *Nat Rev Genet*. 2021 Sep 20. doi: 10.1038/s41576-021-00412-1. Online ahead of print.
- Ben Maamar M, Nilsson EE, Skinner MK. Epigenetic transgenerational inheritance, gametogenesis and germline development. *Biol Reprod*. 2021 Sep 14;105(3):570-592.
- Gaspari L, Paris F, Kalfa N, et al. Experimental Evidence of 2,3,7,8-Tetrachlordibenzo-p-Dioxin (TCDD) Transgenerational Effects on Reproductive Health. *Int J Mol Sci*. 2021 Aug 23;22(16):9091.
- Stener-Victorin E, Deng Q. Transmission of Polycystic Ovary Syndrome via Epigenetic Inheritance. *Trends Mol Med*. 2021 Aug;27(8):723-724.
- Mladenov V, Fotopoulos V, Kaiserli E, et al. Deciphering the Epigenetic Alphabet Involved in Transgenerational Stress Memory in Crops. *Int J Mol Sci*. 2021 Jul 1;22(13):7118.
- Baduel P, Colot V. The epiallelic potential of transposable elements and its evolutionary significance in plants. *Philos Trans R Soc Lond B Biol Sci*. 2021 Jun 7;376(1826):20200123.
- Nilsson E, Sadler-Riggelman I, Beck D, Skinner MK. Differential DNA methylation in somatic and sperm cells of hatchery vs wild (natural-origin) steelhead trout populations. *Environ Epigenet*. 2021 May 19;7(1):dvab002.
- Pierron F, Lorient S, Hérouin D, et al. Transgenerational epigenetic sex determination: Environment experienced by female fish affects offspring sex ratio. *Environ Pollut*. 2021 May 15;277:116864.

- Wautier A, Tournaire M, Devouche E, et al. Genital tract and reproductive characteristics in daughters of women and men prenatally exposed to diethylstilbestrol (DES). *Therapie*. Sep-Oct 2020;75(5):439-448.
- Soley FG. Still no evidence for transgenerational inheritance or absence of epigenetic reprogramming in the honey bee. *Proc Natl Acad Sci U S A*. 2021 Jul 13;118(28):e2108608118.
- Wang HD, Allard P. Challenging dogmas: How transgenerational epigenetics reshapes our views on life. *J Exp Zool A Ecol Integr Physiol*. 2022 Jan;337(1):70-74.
- Jawaid A, Jehle K-L, Mansuy IM. Impact of Parental Exposure on Offspring Health in Humans. *Trends Genet*. 2021 Apr;37(4):373-388.
- Mimouni NEH, Paiva I, Barbotin A-L, et al. Polycystic ovary syndrome is transmitted via a transgenerational epigenetic process. *Cell Metab*. 2021 Mar 2;33(3):513-530.e8.
- Beck D, Ben Maamar M, Skinner MK. Integration of sperm ncRNA-directed DNA methylation and DNA methylation-directed histone retention in epigenetic transgenerational inheritance. *Epigenetics Chromatin*. 2021 Jan 12;14(1):6.
- Bertozzi TM, Elmer JL, Macfarlan TS, Ferguson-Smith AC. KRAB zinc finger protein diversification drives mammalian interindividual methylation variability. *Proc Natl Acad Sci U S A*. 2020 Dec 8;117(49):31290-31300.
- Nilsson EE, Skinner MK. Environmentally Induced Epigenetic Transgenerational Inheritance of Reproductive Disease. *Biol Reprod*. 2015 Dec;93(6):145.
- Duempelmann L, Skribbe M, Bühler M. (2020) Small RNAs in the Transgenerational Inheritance of Epigenetic Information. *Trends Genet*. 2020 Jan 14. doi: 10.1016/j.tig.2019.12.001.
- Patel NJ, Hogan KJ, Rizk E, Stewart K, Madrid A, et al. (2020) Ancestral Folate Promotes Neuronal Regeneration in Serial Generations of Progeny. *Mol Neurobiol*. 2020 Jan 10. doi: 10.1007/s12035-019-01812-5.
- Legoff L, D'Cruz SC, Tevosian S, Primig M, Smagulova F. (2019) Transgenerational Inheritance of Environmentally Induced Epigenetic Alterations during Mammalian Development. *Cells* 3;8(12).
- Meyer DN, Crofts EJ, Akemann C, Gurdziel K, Farr R, Baker BB, Weber D, Baker TR. (2019) Developmental exposure to Pb²⁺ induces transgenerational changes to zebrafish brain transcriptome. *Chemosphere* 2;244:125527.
- King SE, Nilsson E, Beck D, Skinner MK. (2019) Adipocyte epigenetic alterations and potential therapeutic targets in transgenerationally inherited lean and obese phenotypes following ancestral exposures. *Adipocyte*. 8(1):362-378.
- Bodden C, Hannan AJ, Reichelt AC. (2020) Diet-Induced Modification of the Sperm Epigenome Programs Metabolism and Behavior. *Trends Endocrinol Metab*. 31(2):131-149.
- Cavalli G, Heard E. (2019) Advances in epigenetics link genetics to the environment and disease. *Nature*. 571(7766):489-499.
- Reece AS, Hulse GK. (2019) Impacts of cannabinoid epigenetics on human development: reflections on Murphy et al. 'cannabinoid exposure and altered DNA methylation in rat and human sperm' epigenetics 2018; 13: 1208-1221. *Epigenetics*. 14(11):1041-1056.
- Cong W, Miao Y, Xu L, Zhang Y, Yuan C, et al. (2019) Transgenerational memory of gene expression changes induced by heavy metal stress in rice (*Oryza sativa* L.). *BMC Plant Biol*. 27;19(1):282.
- Sarkies P. (2020) Molecular mechanisms of epigenetic inheritance: Possible evolutionary implications. *Semin Cell Dev Biol*. 97:106-115.
- King SE, McBirney M, Beck D, Sadler-Riggelman I, Nilsson E, Skinner MK. (2019) Sperm epimutation biomarkers of obesity and pathologies following DDT induced epigenetic transgenerational inheritance of disease. *Environ Epigenet*. 27;5(2):dvz008.

- Skinner MK, Nilsson E, Sadler-Riggleman I, Beck D, Ben Maamar M, McCarrey JR. (2019) Transgenerational sperm DNA methylation epimutation developmental origins following ancestral vinclozolin exposure. *Epigenetics*. 14(7):721-739.
- Ben Maamar M, Nilsson E, Sadler-Riggleman I, Beck D, McCarrey JR, Skinner MK. (2019) Developmental origins of transgenerational sperm DNA methylation epimutations following ancestral DDT exposure. *Dev Biol*. 15;445(2):280-293.
- Lacal I, Ventura R. (2018) Epigenetic Inheritance: Concepts, Mechanisms and Perspectives. *Front Mol Neurosci*. 28;11:292.
- Nilsson E, Klukovich R, Sadler-Riggleman I, Beck D, Xie Y, Yan W, Skinner MK. Environmental toxicant induced epigenetic transgenerational inheritance of ovarian pathology and granulosa cell epigenome and transcriptome alterations: ancestral origins of polycystic ovarian syndrome and primary ovarian insufficiency. *Epigenetics*. 2018;13(8):875-895.
- Gold HB, Jung YH, Corces VG. Not just heads and tails: The complexity of the sperm epigenome. *J Biol Chem*. 2018 Sep 7;293(36):13815-13820.
- Ben Maamar M, King SE, Nilsson E, Beck D, Skinner MK. Epigenetic transgenerational inheritance of parent-of-origin allelic transmission of outcross pathology and sperm epimutations. *Dev Biol*. 2020 Feb 1;458(1):106-119.
- Sadler-Riggleman I, Klukovich R, Nilsson E, Beck D, Xie Y, Yan W, Skinner MK. Epigenetic transgenerational inheritance of testis pathology and Sertoli cell epimutations: generational origins of male infertility. *Environ Epigenet*. 2019 Aug 29;5(3):dvz013.
- Kubsad D, Nilsson EE, King SE, Sadler-Riggleman I, Beck D, Skinner MK. Assessment of Glyphosate Induced Epigenetic Transgenerational Inheritance of Pathologies and Sperm Epimutations: Generational Toxicology. *Sci Rep*. 2019 Apr 23;9(1):6372.
- Baxter FA, Drake AJ. Non-genetic inheritance via the male germline in mammals. *Philos Trans R Soc Lond B Biol Sci*. 2019 Apr 15;374(1770):20180118.
- Klukovich R, Nilsson E, Sadler-Riggleman I, Beck D, Xie Y, Yan W, Skinner MK. Environmental Toxicant Induced Epigenetic Transgenerational Inheritance of Prostate Pathology and Stromal-Epithelial Cell Epigenome and Transcriptome Alterations: Ancestral Origins of Prostate Disease. *Sci Rep*. 2019 Feb 18;9(1):2209.
- Lecoutre S, Petrus P, Rydén M, Breton C. Transgenerational Epigenetic Mechanisms in Adipose Tissue Development. *Trends Endocrinol Metab*. 2018 Oct;29(10):675-685.
- Duempelmann L, Skribbe M, Bühler M. Small RNAs in the Transgenerational Inheritance of Epigenetic Information. *Trends Genet*. 2020 Jan 14. pii: S0168-9525(19)30259-8. doi: 10.1016/j.tig.2019.12.001. [Epub ahead of print]
- Stermer AR, Wilson SK, Klein D, Hall SJ, Boekelheide K. Trichloroethylene exposure alters dimethylated histone three lysine four in protein kinase A signaling pathway chromatin of rat sperm†. *Biol Reprod*. 2019 Nov 21;101(5):875-877.
- van Steenwyk G, Roszkowski M, Manuella F, Franklin TB, Mansuy IM. Transgenerational inheritance of behavioral and metabolic effects of paternal exposure to traumatic stress in early postnatal life: evidence in the 4th generation. *Environ Epigenet*. 2018 Oct 16;4(2):dvy023.
- Beck D, Sadler-Riggleman I, Skinner MK. (2017) Generational comparisons (F1 versus F3) of vinclozolin induced epigenetic transgenerational inheritance of sperm differential DNA methylation regions (epimutations) using MeDIP-Seq. *Environ Epigenet*. 3(3). pii: dvx016.
- Holder LB, Haque MM, Skinner MK. (2017) Machine learning for epigenetics and future medical applications. *Epigenetics*. 12(7):505-514.

- Craig JR, Jenkins TG, Carrell DT, Hotaling JM. (2017) Obesity, male infertility, and the sperm epigenome. *Fertil Steril.* 107(4):848-859.
- Tikoo K, Vikram A, Shrivastava S, Jena G, Shah H, Chhabra R. (2017) Parental High-Fat Diet Promotes Inflammatory and Senescence-Related Changes in Prostate. *Oxid Med Cell Longev.* 2017;2017:4962950.
- Bruner-Tran KL, Gnecco J, Ding T, Glore DR, Pensabene V, Osteen KG. (2017) Exposure to the environmental endocrine disruptor TCDD and human reproductive dysfunction: Translating lessons from murine models. *Reprod Toxicol.* 68:59-71.
- Ho SM, Cheong A, Adgent MA, Veevers J, Suen AA, Tam NNC, Leung YK, Jefferson WN, Williams CJ. (2017) Environmental factors, epigenetics, and developmental origin of reproductive disorders. *Reprod Toxicol.* 68:85-104.
- Siddeek B, Lakhdari N, Inoubli L, et al. (2016) Developmental epigenetic programming of adult germ cell death disease: Polycomb protein EZH2-miR-101 pathway. *Epigenomics.* 8(11):1459-1479.
- Mykhalchenko K, Lizneva D, Trofimova T, Walker W, Suturina L, Diamond MP, Azziz R. (2017) Genetics of polycystic ovary syndrome. *Expert Rev Mol Diagn.* 17(7):723-733.
- Filippou P, Homburg R. (2017) Is foetal hyperexposure to androgens a cause of PCOS? *Hum Reprod Update.* 1;23(4):421-432.
- Lawrenson K, Kar S, McCue K, et al. (2016) Functional mechanisms underlying pleiotropic risk alleles at the 19p13.1 breast-ovarian cancer susceptibility locus. *Nat Commun.* 7:12675.
- Kar SP, Beesley J, Amin Al Olama A, et al. (2016) Genome-Wide Meta-Analyses of Breast, Ovarian, and Prostate Cancer Association Studies Identify Multiple New Susceptibility Loci Shared by at Least Two Cancer Types. *Cancer Discov.* 6(9):1052-67.
- Cucchiara V, Yang JC, Mirone V, Gao AC, Rosenfeld MG, Evans CP. (2017) Epigenomic Regulation of Androgen Receptor Signaling: Potential Role in Prostate Cancer Therapy. *Cancers (Basel).* 16;9(1). pii: E9. doi: 10.3390/cancers9010009.
- Sampath Kumar A, Seah MK, Ling KY, Wang Y, Tan JH, Nitsch S, Lim SL, Lorthongpanich C, Wollmann H, Low DH, Guccione E, Messerschmidt DM. (2017) Loss of maternal Trim28 causes male-predominant early embryonic lethality. *Genes Dev.* 31(1):12-17.
- van den Berg GJ, Pinger PR. (2016) Transgenerational effects of childhood conditions on third generation health and education outcomes. *Econ Hum Biol.* 23:103-120.
- Vrooman LA, Bartolomei MS. (2017) Can assisted reproductive technologies cause adult-onset disease? Evidence from human and mouse. *Reprod Toxicol.* 68:72-84.
- Reichetzedder C, Dwi Putra SE, Li J, Hocher B. (2016) Developmental Origins of Disease - Crisis Precipitates Change. *Cell Physiol Biochem.* 39(3):919-38.
- Illum LR, Bak ST, Lund S, Nielsen AL. (2017) DNA methylation in epigenetic inheritance of metabolic diseases through the male germ line. *J Mol Endocrinol.* 2017 Dec 4. pii: JME-17-0189. doi: 10.1530/JME-17-0189.
- McBirney M, King SE, Pappalardo M, Houser E, Unkefer M, Nilsson E, Sadler-Riggelman I, Beck D, Winchester P, Skinner MK. (2017) Atrazine induced epigenetic transgenerational inheritance of disease, lean phenotype and sperm epimutation pathology biomarkers. *PLoS One* 12(9):e0184306.
- McCarrey JR, Lehle JD, Raju SS, Wang Y, Nilsson EE, Skinner MK. (2016) Tertiary Epimutations - A Novel Aspect of Epigenetic Transgenerational Inheritance Promoting Genome Instability. *PLoS One.* 11(12):e0168038.
- Blake GE, Watson ED. (2016) Unravelling the complex mechanisms of transgenerational epigenetic inheritance. *Curr Opin Chem Biol.* 33:101-7.

- Sailer C, Schmid B, Grossniklaus U. (2016) Apomixis Allows the Transgenerational Fixation of Phenotypes in Hybrid Plants. *Curr Biol.* 26(3):331-7.
- Stuppia L, Franzago M, Ballerini P, Gatta V, Antonucci I. (2015) Epigenetics and male reproduction: the consequences of paternal lifestyle on fertility, embryo development, and children lifetime health. *Clin Epigenetics.* 11;7:120.
- Keil KP, Vezina CM. (2015) DNA methylation as a dynamic regulator of development and disease processes: spotlight on the prostate. *Epigenomics.* 7(3):413-25.
- Cahenzli F, Wenk BA, Erhardt A. (2015) Female butterflies adapt and allocate their progeny to the host-plant quality of their own larval experience. *Ecology.* 96(7):1966-73.
- Huo X, Chen D, He Y, Zhu W, Zhou W, Zhang J. (2015) Bisphenol-A and Female Infertility: A Possible Role of Gene-Environment Interactions. *Int J Environ Res Public Health.* 7;12(9):11101-16.
- Erwin AA, Galdos MA, Wickersheim ML, et al. (2015) piRNAs Are Associated with Diverse Transgenerational Effects on Gene and Transposon Expression in a Hybrid Dysgenic Syndrome of *D. virilis*. *PLoS Genet.* 2015 Aug 4;11(8):e1005332.
- Jobson MA, Jordan JM, Sandrof MA, et al. (2015) Transgenerational Effects of Early Life Starvation on Growth, Reproduction, and Stress Resistance in *Caenorhabditis elegans*. *Genetics.* 201(1):201-12.
- Quinnies KM, Doyle TJ, Kim KH1, Rissman EF. (2015) Transgenerational Effects of Di-(2-Ethylhexyl) Phthalate (DEHP) on Stress Hormones and Behavior. *Endocrinology.* 156(9):3077-83.
- Fénichel P, Brucker-Davis F, Chevalier N. (2015) The history of Distilbène® (Diethylstilbestrol) told to grandchildren--the transgenerational effect. *Ann Endocrinol (Paris).* 76(3):253-9.
- Bale TL. (2015) Epigenetic and transgenerational reprogramming of brain development. *Nat Rev Neurosci.* 16(6):332-44.
- Ziv-Gal A, Wang W, Zhou C, Flaws JA. (2015) The effects of in utero bisphenol A exposure on reproductive capacity in several generations of mice. *Toxicol Appl Pharmacol.* 1;284(3):354-62.
- Tortiglione C. (2014) The heritable effects of nanotoxicity. *Nanomedicine (Lond).* 9(18):2829-41.
- VandeVoort CA, Grimsrud KN, Midic U, et al. (2015) Transgenerational effects of binge drinking in a primate model: implications for human health. *Fertil Steril.* 103(2):560-9.
- Wei Y, Schatten H, Sun QY. (2015) Environmental epigenetic inheritance through gametes and implications for human reproduction. *Hum Reprod Update.* 21(2):194-208.
- Shama LN, Wegner KM. (2014) Grandparental effects in marine sticklebacks: transgenerational plasticity across multiple generations. *J Evol Biol.* 27(11):2297-307.
- Baker TR, King-Heiden TC, Peterson RE, Heideman W. (2014) Dioxin induction of transgenerational inheritance of disease in zebrafish. *Mol Cell Endocrinol.* 398(1-2):36-41
- Bruner-Tran KL, Ding T, Yeoman KB, Archibong A, Arosh JA, Osteen KG. (2014) Developmental exposure of mice to dioxin promotes transgenerational testicular inflammation and an increased risk of preterm birth in unexposed mating partners. *PLoS One.* 15;9(8):e105084.
- Crews D, Gillette R, Miller-Crews I, Gore AC, Skinner MK. (2014) Nature, nurture and epigenetics. *Mol Cell Endocrinol.* 398(1-2):42-52.
- Kawashima T, Berger F. (2014) Epigenetic reprogramming in plant sexual reproduction. *Nat Rev Genet.* 15(9):613-24.
- Nilsson EE, Skinner MK. (2015) Environmentally induced epigenetic transgenerational inheritance of disease susceptibility. *Transl Res.* 165(1):12-7.

- Grandjean P, Barouki R, Bellinger DC, et al. (2015) Life-Long Implications of Developmental Exposure to Environmental Stressors: New Perspectives. *Endocrinology*. 156(10):3408-15.
- Wei Y, Schatten H, Sun QY. (2015) Environmental epigenetic inheritance through gametes and implications for human reproduction. *Hum Reprod Update*. 21(2):194-208.
- Bao J, Bedford MT. (2016) Epigenetic regulation of the histone-to-protamine transition during Spermiogenesis. *Reproduction*. 2016 Feb 5. pii: REP-15-0562. [Epub ahead of print]
- Tseng YT, Liao HF, Yu CY, Mo CF, Lin SP. (2015) Epigenetic factors in the regulation of prospermatogonia and spermatogonial stem cells. *Reproduction*. 150(3):R77-91.
- Katsi V, Felekos I, Siristatidis C, et al. (2015) Preeclampsia: What Does the Father Have to Do with It? *Curr Hypertens Rep*. 17(8):60.
- Beaujean N. (2014) Epigenetics, embryo quality and developmental potential. *Reprod Fertil Dev*. 27(1):53-62.
- Ge ZJ, Schatten H, Zhang CL, Sun QY. (2015) Oocyte ageing and epigenetics. *Reproduction*. 2015 Mar;149(3):R103-14.
- Vaiserman AM. (2015) Epigenetic programming by early-life stress: Evidence from human populations. *Dev Dyn*. 244(3):254-65.
- Campos EI, Stafford JM, Reinberg D. (2014) Epigenetic inheritance: histone bookmarks across generations. *Trends Cell Biol*. 2014 Nov;24(11):664-74.
- She W, Baroux C. (2014) Chromatin dynamics during plant sexual reproduction. *Front Plant Sci*. 2014 Jul 24;5:354.
- Ding GL, Huang HF. (2013) Paternal transgenerational glucose intolerance with epigenetic alterations in second generation offspring of GDM. *Asian J Androl*. 15(4):451-2.
- Di Giacomo M, Comazzetto S, et al. (2013) Multiple epigenetic mechanisms and the piRNA pathway enforce LINE1 silencing during adult spermatogenesis. *Mol Cell*. 23;50(4):601-8.
- Guerrero-Bosagna C, Savenkova M, Haque MM, Nilsson E, Skinner MK. (2013) Environmentally induced epigenetic transgenerational inheritance of altered Sertoli cell transcriptome and epigenome: molecular etiology of male infertility. *PLoS One*. 8(3):e59922.
- Tracey R, Manikkam M, Guerrero-Bosagna C, Skinner MK. (2013) Hydrocarbons (jet fuel JP-8) induce epigenetic transgenerational inheritance of obesity, reproductive disease and sperm epimutations. *Reprod Toxicol*. 36:104-16.
- Feng X, Zilberman D, Dickinson H. (2013) A conversation across generations: soma-germ cell crosstalk in plants. *Dev Cell*. 11;24(3):215-25.
- Soubry A, Schildkraut JM, et al. (2013) Paternal obesity is associated with IGF2 hypomethylation in newborns: results from a Newborn Epigenetics Study (NEST) cohort. *BMC Med*. 6;11:29.
- Chamorro-García R, Sahu M, et al. (2013) Transgenerational inheritance of increased fat depot size, stem cell reprogramming, and hepatic steatosis elicited by prenatal exposure to the obesogen tributyltin in mice. *Environ Health Perspect*. 121(3):359-66.
- Raychaudhuri N, Dubruielle R, et al. (2012) Transgenerational propagation and quantitative maintenance of paternal centromeres depends on Cid/Cenp-A presence in *Drosophila* sperm. *PLoS Biol*. 10(12):e1001434.
- González-Recio O, Ugarte E, Bach A. (2012) Trans-generational effect of maternal lactation during pregnancy: a Holstein cow model. *PLoS One*. 7(12):e51816.
- Hackett JA, Sengupta R, et al. (2013) Germline DNA demethylation dynamics and imprint erasure through 5-hydroxymethylcytosine. *Science*. 25;339(6118):448-52.

- Guerrero-Bosagna C, Covert TR, et al. (2012) Epigenetic transgenerational inheritance of vinclozolin induced mouse adult onset disease and associated sperm epigenome biomarkers. *Reprod Toxicol.* 34(4):694-707.
- Roseboom TJ, Watson ED. (2012) The next generation of disease risk: are the effects of prenatal nutrition transmitted across generations? Evidence from animal and human studies. *Placenta.* 33 Suppl 2:e40-4.
- Uzumcu M, Zama AM, Oruc E. (2012) Epigenetic mechanisms in the actions of endocrine-disrupting chemicals: gonadal effects and role in female reproduction. *Reprod Domest Anim.* 47 Suppl 4:338-47.
- Wolstenholme JT, Edwards M, et al. (2012) Gestational exposure to bisphenol a produces transgenerational changes in behaviors and gene expression. *Endocrinology.* 153(8):3828-38.
- Govorko D, Bekdash RA, Zhang C, Sarkar DK. (2012) Male germline transmits fetal alcohol adverse effect on hypothalamic proopiomelanocortin gene across generations. *Biol Psychiatry.* 1;72(5):378-88.
- Zeh JA, Bonilla MM, et al. (2012) From father to son: transgenerational effect of tetracycline on sperm viability. *Sci Rep.* 2:375.
- Ding GL, Wang FF, et al. (2012) Transgenerational glucose intolerance with Igf2/H19 epigenetic alterations in mouse islet induced by intrauterine hyperglycemia. *Diabetes.* 61(5):1133-42.
- Manikkam M, Guerrero-Bosagna C, Tracey R, Haque MM, Skinner MK. (2012) Transgenerational actions of environmental compounds on reproductive disease and identification of epigenetic biomarkers of ancestral exposures. *PLoS One.* 7(2):e31901.
- van Montfoort AP, Hanssen LL, et al. (2012) Assisted reproduction treatment and epigenetic inheritance. *Hum Reprod Update.* 18(2):171-97.
- Halliday J. (2012) Outcomes for offspring of men having ICSI for male factor infertility. *Asian J Androl.* 14(1):116-20.
- Burdge GC, Hoile SP, Uller T, et al. (2011) Progressive, transgenerational changes in offspring phenotype and epigenotype following nutritional transition. *PLoS One.* 6(11):e28282.
- Low FM, Gluckman PD, Hanson MA. (2011) Developmental plasticity and epigenetic mechanisms underpinning metabolic and cardiovascular diseases. *Epigenomics.* 3(3):279-94.
- Chan D, Delbès G, Landry M, Robaire B, Trasler JM. (2012) Epigenetic alterations in sperm DNA associated with testicular cancer treatment. *Toxicol Sci.* 125(2):532-43.
- Guerrero-Bosagna C, Skinner MK. (2012) Environmentally induced epigenetic transgenerational inheritance of phenotype and disease. *Mol Cell Endocrinol.* 6;354(1-2):3-8.
- Calle A, Fernandez-Gonzalez R, et al. (2012) Long-term and transgenerational effects of in vitro culture on mouse embryos. *Theriogenology.* 1;77(4):785-93.
- Matthews SG, Phillips DI. (2012) Transgenerational inheritance of stress pathology. *Exp Neurol.* 2012 Jan;233(1):95-101.
- Radtke KM, Ruf M, et al. (2011) Transgenerational impact of intimate partner violence on methylation in the promoter of the glucocorticoid receptor. *Transl Psychiatry.* 19;1:e21.
- Stouder C, Paoloni-Giacobino A. (2011) Specific transgenerational imprinting effects of the endocrine disruptor methoxychlor on male gametes. *Reproduction.* 141(2):207-16.
- Skinner MK, Manikkam M, Guerrero-Bosagna C. (2011) Epigenetic transgenerational actions of endocrine disruptors. *Reprod Toxicol.* 31(3):337-43.
- Pentinat T, Ramon-Krauel M, et al. (2010) Transgenerational inheritance of glucose intolerance in a mouse model of neonatal overnutrition. *Endocrinology.* 151(12):5617-23.

- Yazbek SN, Spiezio SH, Nadeau JH, Buchner DA. (2010) Ancestral paternal genotype controls body weight and food intake for multiple generations. *Hum Mol Genet.* 1;19(21):4134-44.
- Zama AM, Uzumcu M. (2010) Epigenetic effects of endocrine-disrupting chemicals on female reproduction: an ovarian perspective. *Front Neuroendocrinol.* 31(4):420-39.
- Dunn GA, Morgan CP, Bale TL. (2011) Sex-specificity in transgenerational epigenetic programming. *Horm Behav.* 59(3):290-5.
- Skinner MK, Manikkam M, Guerrero-Bosagna C. (2010) Epigenetic transgenerational actions of environmental factors in disease etiology. *Trends Endocrinol Metab.* 21(4):214-22.
- Guerrero-Bosagna CM, Skinner MK. (2009) Epigenetic transgenerational effects of endocrine disruptors on male reproduction. *Semin Reprod Med.* 27(5):403-8.
- Jablonka E, Raz G. (2009) Transgenerational epigenetic inheritance: prevalence, mechanisms, and implications for the study of heredity and evolution. *Q Rev Biol.* 84(2):131-76.
- Wang H, Chai Y, Chu X, et al. (2009) Molecular characterization of a rice mutator-phenotype derived from an incompatible cross-pollination reveals transgenerational mobilization of multiple transposable elements and extensive epigenetic instability. *BMC Plant Biol.* 29;9:63.
- Barber RC, Hardwick RJ, et al. (2009) The effects of in utero irradiation on mutation induction and transgenerational instability in mice. *Mutat Res.* 12;664(1-2):6-12.
- Katz DJ, Edwards TM, Reinke V, Kelly WG. (2009) A *C. elegans* LSD1 demethylase contributes to germline immortality by reprogramming epigenetic memory. *Cell.* 17;137(2):308-20.
- Godmann M, Lambrot R, Kimmins S. (2009) The dynamic epigenetic program in male germ cells: Its role in spermatogenesis, testis cancer, and its response to the environment. *Microsc Res Tech.* 72(8):603-19.
- Youngson NA, Whitelaw E. (2008) Transgenerational epigenetic effects. *Annu Rev Genomics Hum Genet.* 9:233-57.
- Schneider S, Kaufmann W, et al. (2008) Vinclozolin--the lack of a transgenerational effect after oral maternal exposure during organogenesis. *Reprod Toxicol.* 2008 Apr;25(3):352-60.
- Nilsson EE, Anway MD, et al. (2008) Transgenerational epigenetic effects of the endocrine disruptor vinclozolin on pregnancies and female adult onset disease. *Reproduction.* 135(5):713-21.
- Anway MD, Skinner MK. (2008) Transgenerational effects of the endocrine disruptor vinclozolin on the prostate transcriptome and adult onset disease. *Prostate.* 1;68(5):517-29.
- Anway MD, Rekow SS, Skinner MK. (2008) Transgenerational epigenetic programming of the embryonic testis transcriptome. *Genomics.* 91(1):30-40.
- Skinner MK. (2008) What is an epigenetic transgenerational phenotype? F3 or F2. *Reprod Toxicol.* 25(1):2-6.
- Crews D, Gore AC, Hsu TS, et al. (2007) Transgenerational epigenetic imprints on mate preference. *Proc Natl Acad Sci U S A.* 3;104(14):5942-6.
- Jirtle RL, Skinner MK. (2007) Environmental epigenomics and disease susceptibility. *Nat Rev Genet.* 8(4):253-62.
- Gluckman PD, Hanson MA, Beedle AS. (2007) Non-genomic transgenerational inheritance of disease risk. *Bioessays.* 29(2):145-54.
- Price TM, Murphy SK, Younglai EV. (2007) Perspectives: the possible influence of assisted reproductive technologies on transgenerational reproductive effects of environmental endocrine disruptors. *Toxicol Sci.* 96(2):218-26.
- Anway MD, Leathers C, Skinner MK. (2006) Endocrine disruptor vinclozolin induced epigenetic transgenerational adult-onset disease. *Endocrinology.* 147(12):5515-23.

- Anway MD, Memon MA, Uzumcu M, Skinner MK. (2006) Transgenerational effect of the endocrine disruptor vinclozolin on male spermatogenesis. *J Androl.* 27(6):868-79.
- Crews D, McLachlan JA. (2006) Epigenetics, evolution, endocrine disruption, health, and disease. *Endocrinology.* 147(6 Suppl):S4-10.
- Anway MD, Skinner MK. (2006) Epigenetic transgenerational actions of endocrine disruptors. *Endocrinology.* 147(6 Suppl):S43-9.
- Magnúsdóttir E, Surani MA. (2014) How to make a primordial germ cell. *Development.* 141(2):245-52.
- Colaneri A, Wang T, et al. (2013) A minimal set of tissue-specific hypomethylated CpGs constitute epigenetic signatures of developmental programming. *PLoS One.* 12;8(9):e72670.
- Skinner MK, Haque CG, Nilsson E, et al. (2013) Environmentally induced transgenerational epigenetic reprogramming of primordial germ cells and the subsequent germ line. *PLoS One.* 15;8(7):e66318.
- Wang H, Wang X, Xu X, Zwaka TP, Cooney AJ. (2013) Epigenetic Re-programming of the Germ Cell Nuclear Factor Gene is Required for Proper Differentiation of Induced Pluripotent Cells. *Stem Cells.* 2013 Mar 14. [Epub ahead of print]
- Cantone I, Fisher AG. (2013) Epigenetic programming and reprogramming during development. *Nat Struct Mol Biol.* 20(3):282-9.
- Park YJ, Herman H, et al. (2012) Sequences sufficient for programming imprinted germline DNA methylation defined. *PLoS One.* 7(3):e33024.
- Niles KM, Chan D, et al. (2011) Critical period of nonpromoter DNA methylation acquisition during prenatal male germ cell development. *PLoS One.* 6(9):e24156.
- Eguizabal C, Shovlin TC, et al. (2009) Generation of primordial germ cells from pluripotent stem cells. *Differentiation.* 78(2-3):116-23.
- Hemberger M, Dean W, Reik W. (2009) Epigenetic dynamics of stem cells and cell lineage commitment: digging Waddington's canal. *Nat Rev Mol Cell Biol.* 10(8):526-37.
- Murphey P, Yamazaki Y, et al. (2009) Epigenetic regulation of genetic integrity is reprogrammed during cloning. *Proc Natl Acad Sci U S A.* 24;106(12):4731-5.
- Shao C, Li Q, Chen S, et al. (2014) Epigenetic modification and inheritance in sexual reversal of fish. *Genome Res.* Feb 2. [Epub ahead of print].
- Sharma A. (2014) Novel transcriptome data analysis implicates circulating microRNAs in epigenetic inheritance in mammals. *Gene.* Jan 29. [Epub ahead of print].
- Wei Y, Yang CR, et al. (2014) Paternally induced transgenerational inheritance of susceptibility to diabetes in mammals. *Proc Natl Acad Sci U S A.* 4;111(5):1873-8.
- Soubry A, Hoyo C, Jirtle RL, Murphy SK. (2014) A paternal environmental legacy: Evidence for epigenetic inheritance through the male germ line. *Bioessays.* Jan 16. [Epub ahead of print].
- Zheng X, Chen L, Li M, et al. (2013) Transgenerational variations in DNA methylation induced by drought stress in two rice varieties with distinguished difference to drought resistance. *PLoS One.* 11;8(11):e80253.
- Skinner MK, Manikkam M, et al. (2013) Ancestral dichlorodiphenyltrichloroethane (DDT) exposure promotes epigenetic transgenerational inheritance of obesity. *BMC Med.* 23;11:228.
- Blewitt M, Whitelaw E. (2013) The use of mouse models to study epigenetics. *Cold Spring Harb Perspect Biol.* 2013 Nov 1;5(11):a017939. doi: 10.1101/cshperspect.a017939.
- Guérin TM, Palladino F, Robert VJ. (2013) Transgenerational functions of small RNA pathways in controlling gene expression in *C. elegans*. *Epigenetics.* 25;9(1). [Epub ahead of print]

- Berghof TV, Parmentier HK, Lammers A. (2013) Transgenerational epigenetic effects on innate immunity in broilers: an underestimated field to be explored? *Poult Sci.* 92(11):2904-13.
- Aiken CE, Ozanne SE. (2014) Transgenerational developmental programming. *Hum Reprod Update.* 20(1):63-75.
- Padmanabhan N, Jia D, et al. (2013) Mutation in folate metabolism causes epigenetic instability and transgenerational effects on development. *Cell.* 26;155(1):81-93.
- Bond DM, Baulcombe DC. (2014) Small RNAs and heritable epigenetic variation in plants. *Trends Cell Biol.* 24(2):100-7.
- Vassoler FM, Sadri-Vakili G. (2013) Mechanisms of transgenerational inheritance of addictive-like behaviors. *Neuroscience.* S0306-4522(13)00658-1. [Epub ahead of print]
- Ayala-García B, López-Santibáñez Guevara M, et al. (2013) Speciation, phenotypic variation and plasticity: what can endocrine disruptors tell us? *Int J Endocrinol.* ;2013:862739.
- Kiani J, Grandjean V, Liebers R, et al. (2013) RNA-mediated epigenetic heredity requires the cytosine methyltransferase Dnmt2. *PLoS Genet.* 9(5):e1003498.
- Feng X, Guang S. (2013) Small RNAs, RNAi and the inheritance of gene silencing in *Caenorhabditis elegans*. *J Genet Genomics.* 20;40(4):153-60.
- Suter L, Widmer A. (2013) Environmental heat and salt stress induce transgenerational phenotypic changes in *Arabidopsis thaliana*. *PLoS One* 9;8(4):e60364.
- Shiota H, Goudarzi A, Rousseaux S, Khochbin S. (2013) Transgenerational inheritance of chromatin states. *Epigenomics.* 5(2):121-2.
- Guerrero-Bosagna C, Savenkova M, et al. (2013) Environmentally induced epigenetic transgenerational inheritance of altered Sertoli cell transcriptome and epigenome: molecular etiology of male infertility. *PLoS One.* ;8(3):e59922.
- Tricker PJ, López CM, (2013) Transgenerational, dynamic methylation of stomata genes in response to low relative humidity. *Int J Mol Sci.* 26;14(4):6674-89.
- Grossniklaus U, Kelly WG, et al. (2013) Transgenerational epigenetic inheritance: how important is it? *Nat Rev Genet.* 14(3):228-35.
- Lim JP, Brunet A. (2013) Bridging the transgenerational gap with epigenetic memory. *Trends Genet.* 29(3):176-86.
- Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner MK. (2013) Plastics derived endocrine disruptors (BPA, DEHP and DBP) induce epigenetic transgenerational inheritance of obesity, reproductive disease and sperm epimutations. *PLoS One.* 8(1):e55387.
- Wan M, Gu H, Wang J, et al. (2013) Inducible mouse models illuminate parameters influencing epigenetic inheritance. *Development.* 140(4):843-52.
- Yan B, Ma L. (2012) The role for maternal piRNA in transgenerational inheritance. *Epigenomics.* 4(5):484-5.
- Sharma A. (2013) Transgenerational epigenetic inheritance: focus on soma to germline information transfer. *Prog Biophys Mol Biol.* 113(3):439-46.
- Bohacek J, Gapp K, Saab BJ, Mansuy IM. (2013) Transgenerational epigenetic effects on brain functions. *Biol Psychiatry.* 15;73(4):313-20.
- Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner MK. (2012) Dioxin (TCDD) induces epigenetic transgenerational inheritance of adult onset disease and sperm epimutations. *PLoS One.* 7(9):e46249.
- Skinner MK, Mohan M, et al. (2012) Epigenetic transgenerational inheritance of somatic transcriptomes and epigenetic control regions. *Genome Biol.* 3;13(10):R91.

- Ou X, Zhang Y, Xu C, et al. (2012) Transgenerational inheritance of modified DNA methylation patterns and enhanced tolerance induced by heavy metal stress in rice (*Oryza sativa* L.). *PLoS One*. 2012;7(9):e41143.
- Manikkam M, Tracey R, et al. (2012) Pesticide and insect repellent mixture (permethrin and DEET) induces epigenetic transgenerational inheritance of disease and sperm epimutations. *Reprod Toxicol*. 34(4):708-19.
- Becker C, Weigel D. (2012) Epigenetic variation: origin and transgenerational inheritance. *Curr Opin Plant Biol*. 15(5):562-7.
- Ashe A, Sapetschnig A, et al. (2012) piRNAs can trigger a multigenerational epigenetic memory in the germline of *C. elegans*. *Cell*. 6;150(1):88-99.
- Crews D, Gillette R, et al (2012) Epigenetic transgenerational inheritance of altered stress responses. *Proc Natl Acad Sci U S A*. 5;109(23):9143-8.
- Kovalchuk I. (2012) Transgenerational epigenetic inheritance in animals. *Front Genet*. 9;3:76.
- Nilsson E, Larsen G, Manikkam M, et al. (2012) Environmentally induced epigenetic transgenerational inheritance of ovarian disease. *PLoS One*. 7(5):e36129.
- Grentzinger T, Armenise C, et al. (2012) piRNA-mediated transgenerational inheritance of an acquired trait. *Genome Res*. 22(10):1877-88.
- Goerlich VC, Nätt D, et al. (2012) Transgenerational effects of early experience on behavioral, hormonal and gene expression responses to acute stress in the precocial chicken. *Horm Behav*. 61(5):711-8.
- Braunschweig M, Jagannathan V, et al. (2012) Investigations on transgenerational epigenetic response down the male line in F2 pigs. *PLoS One*. 7(2):e30583.
- Daxinger L, Whitelaw E. (2012) Understanding transgenerational epigenetic inheritance via the gametes in mammals. *Nat Rev Genet*. 31;13(3):153-62.
- Berger SL. (2012) Transgenerational inheritance of longevity: epigenetic mysteries abound. *Cell Metab*. 4;15(1):6-7.
- Benayoun BA, Brunet A. (2012) Epigenetic memory of longevity in *Caenorhabditis elegans*. *Worm*. 1;1(1):77-81.
- Nelson VR, Nadeau JH. (2010) Transgenerational genetic effects. *Epigenomics*. ;2(6):797-806.
- Guerrero-Bosagna C, Skinner MK. (2012) Environmentally induced epigenetic transgenerational inheritance of phenotype and disease. *Mol Cell Endocrinol*. 6;354(1-2):3-8.
- Greer EL, Maures TJ, et al. (2011) Transgenerational epigenetic inheritance of longevity in *Caenorhabditis elegans*. *Nature*. 19;479(7373):365-71.
- Skinner MK. (2011) Environmental epigenetic transgenerational inheritance and somatic epigenetic mitotic stability. *Epigenetics*. 6(7):838-42.
- Hausser MT, Aufsatz W, Jonak C, Luschnig C. (2011) Transgenerational epigenetic inheritance in plants. *Biochim Biophys Acta*. 1809(8):459-68.
- Skinner MK. (2011) Role of epigenetics in developmental biology and transgenerational inheritance. *Birth Defects Res C Embryo Today*. 93(1):51-5.
- Paszkowski J, Grossniklaus U. (2011) Selected aspects of transgenerational epigenetic inheritance and resetting in plants. *Curr Opin Plant Biol*. 14(2):195-203.
- Matthews SG, Phillips DI. (2012) Transgenerational inheritance of stress pathology. *Exp Neurol*. 233(1):95-101.
- Walker DM, Gore AC. (2011) Transgenerational neuroendocrine disruption of reproduction. *Nat Rev Endocrinol*. 7(4):197-207.

- Carone BR, Fauquier L, et al. (2010) Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell*. 23;143(7):1084-96.
- Pembrey ME. (2010) Male-line transgenerational responses in humans. *Hum Fertil (Camb)*. 13(4):268-71.
- Arai JA, Feig LA. (2011) Long-lasting and transgenerational effects of an environmental enrichment on memory formation. *Brain Res Bull*. 25;85(1-2):30-5.
- Daxinger L, Whitelaw E. (2010) Transgenerational epigenetic inheritance: more questions than answers. *Genome Res*. 20(12):1623-8.
- Pentinat T, Ramon-Krauel M, et al. (2010) Transgenerational inheritance of glucose intolerance in a mouse model of neonatal overnutrition. *Endocrinology*. 151(12):5617-23.
- Guerrero-Bosagna C, Settles M, Lucker B, Skinner MK. (2010) Epigenetic transgenerational actions of vinclozolin on promoter regions of the sperm epigenome. *PLoS One*. 30;5(9). pii: e13100.
- Skinner MK, Guerrero-Bosagna C. (2009) Environmental signals and transgenerational epigenetics. *Epigenomics*. 1(1):111-117.
- Dunn GA, Morgan CP, Bale TL. (2011) Sex-specificity in transgenerational epigenetic programming. *Horm Behav*. 59(3):290-5.
- Sharma A, Singh P. (2009) Detection of transgenerational spermatogenic inheritance of adult male acquired CNS gene expression characteristics using a *Drosophila* systems model. *PLoS One*. 2;4(6):e5763.
- Slatkin M. (2009) Epigenetic inheritance and the missing heritability problem. *Genetics*.182(3):845-50.

Beyond Genes Special Issue

Epigenetic transgenerational inheritance, gametogenesis and germline development[†]

Millissia Ben Maamar, Eric E. Nilsson and Michael K. Skinner*

Center for Reproductive Biology, School of Biological Sciences, Washington State University, Pullman, WA, USA

***Correspondence:** Center for Reproductive Biology, School of Biological Sciences, Washington State University, Pullman, WA, 99164-4236, USA. Tel: 509-335-1524; E-mail: skinner@wsu.edu

[†]**Grant support:** This study was supported by John Templeton Foundation (50183 and 61174) (<https://templeton.org/>) grants to M.K.S. and NIH (ES012974) (<https://www.nih.gov/>) grant to M.K.S. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Received 25 January 2021; Revised 12 April 2021; Accepted 22 April 2021

Abstract

One of the most important developing cell types in any biological system is the gamete (sperm and egg). The transmission of phenotypes and optimally adapted physiology to subsequent generations is in large part controlled by gametogenesis. In contrast to genetics, the environment actively regulates epigenetics to impact the physiology and phenotype of cellular and biological systems. The integration of epigenetics and genetics is critical for all developmental biology systems at the cellular and organism level. The current review is focused on the role of epigenetics during gametogenesis for both the spermatogenesis system in the male and oogenesis system in the female. The developmental stages from the initial primordial germ cell through gametogenesis to the mature sperm and egg are presented. How environmental factors can influence the epigenetics of gametogenesis to impact the epigenetic transgenerational inheritance of phenotypic and physiological change in subsequent generations is reviewed.

Summary sentence

How environmental factors can influence the epigenetics of gametogenesis to impact the epigenetic transgenerational inheritance of phenotypic and physiological change in subsequent generations is reviewed.

Key words: Gametogenesis, Spermatogenesis, Oogenesis, PGCs, Epigenetics, Transgenerational, Review.

Introduction

The germ line is an enduring link between all generations of a species. After the fertilization of the oocyte by the sperm, a totipotent zygote will give rise to all cell lineages of the organism, including the germ line itself. The primordial germ cells (PGCs) are the precursor pluripotent stem cells for the sperm and egg. They are established during the perigastrulation epiblast stage of the mammalian embryo. The PGCs specification is regulated by a unique and complex gene network induced by signals from extra-embryonic tissues [1]. In

the gonads, these PGCs will differentiate into the male prospermatogonia or female oogonia in response to Sertoli or granulosa cell signaling. The prospermatogonia continue in the gametogenesis process and undergo spermatogenesis to develop into the mature sperm. The oogonia continue into the gametogenesis process and undergo oogenesis to develop into the mature oocyte. Therefore, gametogenesis can be seen as a crucial first step for the perpetuation of the mammalian life cycle [2].

The crucial aspect of gametogenesis is the production of genetically and epigenetically competent gametes, which are necessary

for normal fertilization and the organism's development. Epigenetics is defined as the factors and processes around DNA that require genome activity independent of DNA sequence, and are mitotically stable. The components include DNA methylation, histone modifications, and non-coding RNA chromatin structure that regulate gene activity independent of DNA sequence [3]. Maternal and paternal gametes display genomic imprinted differences due to DNA methylation and other epigenetic modifications established in the germ line during gametogenesis, but this is only a small component of the epigenome and its regulation of biology.

Interestingly, previous studies have shown that epigenetic modifications can occur during gametogenesis under the influence of environmental factors (stress, diet, pollutants, etc.), which can lead to phenotypical defects in the individuals exposed and in the subsequent generations through the germline. This non-genetic form of inheritance is termed epigenetic transgenerational inheritance and is mediated through epigenetic alterations (i.e. epimutations) in the sperm or egg.

The current review presents the molecular basis of germ cell development (i.e. gametogenesis), and also how the environment can induce stable epimutations and modified phenotypes through the transgenerational inheritance phenomenon.

Mammalian gametogenesis and primordial germ cell development

Primordial germ cell specification

The totipotent stem cells derived from the zygote are the product of fertilization of the oocyte by the sperm, which gives rise to all cell lineages of an organism, and the germline itself. Thus, the specification of primordial germ cells (PGCs) is a pivotal first step for the acquisition of the germline pluripotent cell and the continuation of the mammalian life cycle [1]. In metazoans, two different processes form the germline in the male and female, giving rise to sperm and oocytes. *Caenorhabditis elegans* and *Drosophila melanogaster* have been used to describe the mechanism for PGC specification in invertebrates, and Zebrafish and *Xenopus* in non-mammalian vertebrates [4–7]. At the onset of development, preformation of germ plasm segregates the germ and the soma. The germ plasm is comprised of RNA, proteins, and organelles that are grouped in a specific location within the oocytes, then allocated to a few cells in the germline of the developing embryo [8]. In this instance, the germline is always differentiated from somatic cells across generations. Alternatively, in mammals, the germline is induced within a population of pluripotent cells. The ectopic expression of germline genes in the soma can be tumorigenic [9], however, specification failure in the germline is a reproductive dead end. This specification process requires a precise orchestration to ensure a timely restriction from the soma. Somatic cells not allocated to the germline will undergo differentiation and perish, whereas the germline has the ability of establishing a new organism in the next generation [10]. The primordial germ cells are specified during early embryonic development. Bone morphogenic protein (BMP) signaling is indispensable for PGC specification, and targeted disruption of *Bmp2*, *Bmp4*, *Bmp8b*, or BMP signaling transducers *Smad1*, *Smad4*, *Smad5* or *Alk2*, all demonstrate loss or reduced numbers of PGC [11–16].

The first phase of gametogenesis happens in early embryogenesis, during the formation and migration of PGCs into the gonadal ridge [17, 18]. Mouse models have been primarily used to study the mammalian germ cell development. In the early post-implantation

embryo epiblast, PGCs specification is initiated. At mouse embryonic day E6.25 in some pluripotent epiblast cells, bone morphogenic protein (BMP) and WNT signals from extra-embryonic tissues to induce the expression of a key regulator of PGC fate: the PR domain zinc-finger protein 1 (PRDM1, also known as BLIMP1) [2, 19]. Two other factors are upregulated next, PRDM14 and the transcription factor AP2 γ (encoded by *Tfap2c*) [20, 21]. The germ cell fate is then induced by the transcription factor network formed by PRDM1, PRDM14, and AP2 γ [22–24]. This tripartite network suppresses somatic gene expression such as *Hoxa1*, *Hoxb1*, *Lim1*, *Evx1*, *Fgf8*, and *Snail* genes, while initiating the germ cell transcriptional program but also setting off a genome-wide epigenetic reprogramming [13, 20, 22, 25–28]. Interestingly, Blimp1 and Prdm14 have distinct binding patterns relative to promoters, whereas Blimp1 is important for the repression of almost all genes usually downregulated in PGCs, as well as for the restoration of pluripotency and epigenetic reprogramming (Figure 1) [29]. The restoration of pluripotency and epigenetic reprogramming are regulated by Prdm14, independently from Blimp1, that defines a novel genetic pathway with strict specificity to the germ cell lineage [30]. In mice, the knockout (KO) of Blimp1, Prdm14, or Ap2 γ result in PGCs specification defects highlighting the fact that these three factors are dominant coordinators of the transcriptional program for the establishment of the germ cell fate. In addition, the concomitant overexpression of these three factors in cells *in vitro* induces mouse germ cell formation in the absence of cytokines [22] shows again the importance of these three transcription factors. After embryonic day 7 (E7) in the mouse, the PGCs are then specified and express PGC-characteristic markers, such as stage-specific embryonic antigen 1 (SSEA1) or developmental pluripotency associated 3 (DPPA3 or STELLA) [31–34]. The expression of several pluripotent genes is also maintained in the PGCs such as *Nanog*, *Oct4* and SRY (sex-determining region-Y) [35–38]. Sybirna and collaborators recently revealed a crucial role for PRDM14 in human germ cell fate, where a loss of function affects the efficiency of specification and results in an aberrant hPGCLC transcriptome. Moreover, their study showed that PRDM14 targets are not conserved between mouse and human, which highlights the evolutionary divergence in the molecular network for PGC specification [39].

Although, most studies have been conducted in mice, recent findings in non-rodent mammals have highlighted the similarities and differences between species. In humans, the PGC formation occurs in the third week of gestation. *In vitro* studies of human PGCs have shown that these cells originate from mesodermal precursor cells, and BMP and WNT signaling pathways are also essential for specification [1, 40, 41]. In contrast to mice, human PGCs lack Sox2 expression. Therefore, the species differences between human and mice PGC transcriptional network may be explained by the differences in either pluripotency circuitry or embryonic origin [40].

Primordial germ cell migration

Just after their specification, between E7.5 and E10.5, while proliferating the PGCs migrate through the hindgut and genital ridge then into the developing gonads to differentiate into gametes [27, 42, 43]. Two germ cell–soma signaling pathways cKIT-STEEL and SDF-CXCR4 are required for the normal migration of PGCs. In mice, germ cells express c-KIT, whereas STEEL is expressed in the somatic cells lining the route to the gonad. The interaction between c-KIT and STEEL is fundamental for PGC proliferation, survival, and migration from the primitive streak to the genital ridge




Figure 1. Human germline development. Just after fertilization, a zygote is formed. At week 1, the blastocyst develops and contains pluripotent epiblast cells, which will give rise to all lineages in the embryo, including the germ line. At week 2, the blastocyst implants into the uterine wall. The human primordial germ cells (hPGCs) are probably specified around the time of gastrulation around week 3. At week 4, the hPGCs are localized near the yolk sac wall close to the allantois. After that stage, the hPGCs migrate through the hindgut to the developing genital ridges. At this developmental stage, the migrating hPGCs go through a genome-wide epigenetic reprogramming, including global DNA demethylation, to erase imprints and other somatic epigenetic marks. During the fetus development and adult life, the germ line will undergo meiosis and gametogenesis to differentiate into sperm and eggs. At the same time, the genome is remethylated and acquires appropriate epigenetic signatures for the generation of a totipotent zygote upon fertilization (modified from [1]).

[44–48]. Sterility because of a lack of spermatogonial stem cells and thus differentiated germ cells has been observed in homozygous *cKit* and *Steel* mutant mice [46, 49–51]. The chemoattractant stromal cell-derived factor 1 (SDF-1) expressed at the genital ridges in the surrounding mesenchyme also facilitates the directional PGC migration. SDF-1 is detected by its receptor C-X-C chemokine receptor type 4 (CXCR4) expressed on the surface of PGCs [47]. A knockdown of the activity of CXCR4b and of the SDF-1a ligand has been shown to result in severe PGC migration defects such as very few PGCs reaching the genital ridge [52]. Alternatively, the migration of PGCs can be redirected toward sites of ectopically expressed SDF-1a [52–54]. This ectopic expression of SDF-1 can account for the development of some extra-gonadal cell tumors in humans [27].

During their migration, PGCs continue to proliferate and reach 500 cells in each fetal gonad at E10.5 in the mouse [55]. At this stage, the PGC differentiate into oogonia in females or gonocytes (i.e. prospermatogonia) in males. To form germline cysts, between E10.5

and E14.5, the fetal germ cells undergo five additional mitotic divisions with incomplete cytokinesis [56]. These cysts cluster together and will form germ cell nests in both female and male fetal gonads [55–57]. These germ cell nests will then resolve and generate the primary oocytes or prospermatogonia in the differentiated female and male gonads, respectively.

Primordial germ cells and epigenetics

DNA methylation. The first DNA methylation erasure period is incomplete and happens in the early embryo, leaving paternally and maternally imprinted genes intact and in all somatic lineages subsequently derived. The second DNA methylation erasure period is more comprehensive and occurs during the germline specification. The function of the DNA methylation erasure is to generate in the case of the embryo a totipotent stem cell population, and for PGCs a pluripotent stem cell population. However, despite these two




Figure 2. Epigenetic reprogramming (DNA methylation erasure) during primordial germ cell development at gonadal sex determination and following fertilization in the early embryo (modified from [283]).

epigenetic methylation erasure processes, epigenetic information can be passed down to the offspring, similar to imprinted genes, even though the mechanisms behind this process remain to be elucidated (Figure 2).

Upon specification, during the rapid proliferation of PGCs, the first DNA methylation erasure happens and consists of a passive DNA methylation between E6.5 and E10.5 in the mouse, which results from repressing *de novo* DNA methyltransferases DNMT3a/b [58–62]. Still, the maintenance of DNA methyltransferase 1 (DNMT1) prevents the dilution of DNA methylation modifications on parentally imprinted regions and meiotic gene promoters [63]. Moreover, a loss of *Dnmt1* in PGCs results in a premature meiotic entry in females and precocious differentiation in prospermatogonia in males, and causes infertility [63].

Further demethylation erasure takes place between E10.5 and E12.5 in the mouse, while the PGCs migrate to the genital ridge and begin sex determination. At this period, DNA methylation is at its lowest level due to enzymatic activity from TET1 and TET2 removing DNA methylation [64–66]. During this DNA demethylation erasure window, the maternally and paternally imprinted loci and resistant promoter regions are erased [65, 67]. Interestingly, several studies have demonstrated that a loss of TET1 and 2 in the germ cells does not impact infertility, and few loci showed altered epigenetic states [65, 68–70]. Despite the near complete DNA demethylation erasure process, some genome loci remain methylated in human and mouse PGCs and are referred as ‘escapees’ [58, 64, 71, 72]. In both species, these escapees have been shown to be associated with retrotransposable elements [58, 64, 71–73], subtelomeric regions [71] and pericentromeric satellite repeats [72] also display these escapees.

The PGCs extensive epigenetic reprogramming includes a genome-wide loss of approximately 90% of 5-methylcytosine (5mC) [58, 64, 65, 68, 71, 74–77]. Even though the underlying molecular

mechanisms of this process have until recently remained unclear, a set of germline reprogramming responsive (GRR) genes is crucial for the correct progression of PGC development and gametogenesis. These genes show unique promoter sequence characteristics, with high levels of both 5mC and 5-hydroxymethylcytosine (5hmC). These genes are targets of TET1, the ten eleven translocation (TET) enzymes, which oxidize 5mCs and promote locus-specific reversal of DNA methylation [78]. The PRC1 is the canonical polycomb repressive complex PRC1, which promotes compact local chromatin structures and longer-range chromatin interactions [79]. The loss of DNA methylation combined with PRC1 repression is uniquely required for GRR gene activation. In this epigenetically poised state, TET1 is required to potentiate a full and efficient activation. TET1 seems to be especially important in female PGCs [68], since they start meiotic prophase soon after completion of epigenetic reprogramming, thus requiring the timely expression of these genes. A slight hypermethylation at GRR gene promoters in the mouse E14.5 *Tet1*^{-/-} PGCs, Hill and collaborators also demonstrate that TET1 stimulates transcription of GRR genes via a DNA demethylation-independent mechanism [80, 81]. In addition, TET1 may also enhance transcription through regulation of the levels of 5mC and 5hmC at non-promoter cis-active elements, such as enhancers [79]. TET1 might also have a critical role in the subsequent removal of aberrant residual and/or *de novo* DNA methylation [79]. This suggests that global reprogramming events require efficient protection from *de novo* DNA methylation to stabilize the newly acquired epigenetic state after the removal of 5mC. PGC epigenetic reprogramming entails complex erasure of epigenetic information and suggests that to enable gametogenesis, a timely and efficient activation of GRR genes is required [79].

Usually, a loss of DNA methylation in somatic cells induces an ectopic expression of retrotransposons, an anarchic proliferation, and apoptosis [82]. However, PGCs develop normally despite this

hypomethylated state. The hypothesis is that the chromatin reorganization could enhance the genome stability and ensure proper chromosome alignment and segregation during mitosis, as well as global transcriptional quiescence during this developmental period [25, 58, 64, 71, 74, 83]. Studies in different models have shown that this process seem to be conserved in multiple species. Experiments with human PGCs have found that these DNA demethylation events follow the same patterns as the ones found in the rodents [72, 73, 84].

Histone marks. In a murine model, before the DNA methylation erasure, the pre-migratory PGCs start a process of reprogramming that erases epigenetic marks. One of the central epigenetic changes in pre-migratory and early migratory mouse PGCs is the loss of H3K9me2. This event is followed by an accumulation of H3K27me3 signal [74, 83, 85]. Eguizabal and collaborators showed in early human gonadal PGCs similar chromatin changes in the human early developing germ line [86]. An erasure of genomic imprints and dynamic changes in chromatin modifications are observed in the mouse PGCs after their entry in the genital ridge [74]. After week 9 of gestation, and similarly to porcine PGCs, human PGCs lose H3K27me3 [86].

Spermatogenesis & spermiogenesis

Male fertility relies on the production by the testis of large numbers of normal spermatozoa. This process is known as spermatogenesis, which can be divided in three major steps: (i) mitosis with the multiplication of the spermatogonia, (ii) meiosis to reduce the number of chromosomes from diploid to haploid cells, which starts with type B spermatogonia into the prophase of the first meiotic division. The cells are then called primary spermatocytes that then divide to form secondary spermatocytes, which will divide and undergo meiosis to form the round spermatids, (iii) and finally, the spermiogenesis, which refers to the successful maturation of round spermatids into spermatozoa [87]. All of these steps are central in the spermatogenic process, and any defect during the spermatogenesis can result in the reduction or absence of sperm production, or production of abnormal sperm (Figure 3).

Endocrine and paracrine regulation

FSH. Many studies in the rat have defined the stages at which testosterone and follicle stimulating hormone (FSH) act during spermatogenesis. The general consensus, until the mid-1990s, was that FSH was paramount for the initiation of spermatogenesis, but its role in the adult was to maintain a normal quantitative germ cell production [88, 89]. However, the study of transgenic mice lacking FSH or its receptor (FSHR) showed that the males were fertile but with a reduced germ cell number [90–94]. Closer observations of these KO mice revealed a reduction in spermatogonia, spermatocytes, and spermatids numbers, which suggests that FSH increases the number of spermatogonia and facilitates their entry into meiosis [90–94]. However, studies on *hpg*. SCARKO (hypogonadal mice lacking gonadotrophins and intratesticular androgen crossed with mice lacking androgen receptors specifically on the Sertoli cells) or *hpg*. ARKO mice (hypogonadal mice lacking gonadotrophins and intratesticular androgen crossed with mice lacking androgen receptors ubiquitously) have shown that FSH does not induce round spermatid formation [95–98].

While a lack of FSH action has been shown to impact spermatogenesis, it is difficult to determine the role of this hormone on

maintaining this process. This is explained by the fact that FSH or its receptor are missing from the start of reproductive development. During normal adult spermatogenesis, apoptosis is a sporadic event, occurring mainly among spermatogonia. Lack of FSH, before sexual maturity and during the first wave of spermatogenesis, which is accompanied by an outburst of focal apoptosis among germ cells, increases the level of cells dying which may impact the adult spermatogenesis [99]. Other studies in rats have suggested that FSH-treatment acts to increase spermatogonia and spermatocyte numbers but displays a limited or incomplete effect on spermatogenesis [100, 101]. While these findings mostly agree with the data with the FSHRKO mice, there are significant differences. In these two studies, it was proposed that FSH could promote the completion of meiosis in rats, which was not observed in the transgenic mice models [101, 102]. The mechanism of action of FSH remains unclear even though FSH can act indirectly through Sertoli cells to increase spermatogonial differentiation/proliferation, but also modify rates of germ cell apoptosis [103–107]. A further role for FSH in the testis might be maintenance of Sertoli cell water balance as an accumulation of fluid was observed in FSHRKO mice cells [108]. As a result, this can alter cell morphology and interactions between germ cells and Sertoli cells, thus could reduce normal spermatogenic efficiency.

Androgen. Androgen plays an essential role in development and maintenance of spermatogenesis, which has been emphasized by a study demonstrating that the precocious expression of androgen receptors (ARs) in Sertoli cells leads to premature spermatogenic cell development [109]. The role of androgens has been clearly demonstrated in any animal model in which androgen levels are reduced, such as through hypophysectomy, GnRH-treatment (agonist or antagonist), ethanedimethane sulfonate treatment (EDS) (which ablates Leydig cells), or in gonadotrophin-deficient mice. In all of these cases, significant loss of pachytene spermatocytes and round spermatids, especially at stages VII and VIII of spermatogenesis, can be reversed by a treatment with testosterone [98, 110–117]. Moreover, in mice lacking functional androgen receptors (*tfm* or ARKO), there is a significant loss of spermatocytes which are also unable to complete meiosis and form round spermatids [118–121]. Androgen maintains indirectly through the Sertoli cells meiosis, which appears to ensure the survival of pachytene spermatocytes and enable diplotene spermatocytes to enter meiotic division [121]. However, the role of androgens in spermiogenesis and spermiation remains unclear in regard to its role on the germ cell niche. Most testis cell types express androgen receptors, except the germ cells. Several studies have concluded that androgen action in the testis is only mediated through somatic cell populations [107, 122, 123].

Estrogen. The physiological role of estrogens in the adult testis has yet to be completely understood. However, some studies suggest that estrogen action is required in the neonate to enable a normal spermatogenesis in adulthood. Estrogen has multiple indirect effects through endocrine regulation and through other tissues on the testis, which makes the study of its action even more difficult. In the adult *hpg* mouse, estrogens stimulate spermatogenesis by actually stimulating the FSH release from the pituitary [124, 125]. Interestingly, exogenous estrogens inhibit spermatogenesis in normal adult animals by inhibiting LH secretion and intratesticular testosterone levels [126]. In some species, the aromatase activity present in the testis can convert androgens to estrogens, such as the horse testes,




Figure 3. Gametogenesis and spermatogenic germ cell stages (modified from [162]).

which produce large concentrations of estrogens [127, 128]. Several cell types in the testis, including the germ cells, express nuclear estrogen receptors ($ER\alpha$ and $ER\beta$), and the membrane cell receptor GPR30 [129, 130]. In ArKO mice (lacking aromatase), the males are initially fertile, but spermatogenesis degenerates and an arrest is observed in the early stages of spermiogenesis and multinucleated cells in the tubular lumen appear [131]. Moreover, during neonatal period, estrogen-dependent $ER\alpha$ signaling is required for a normal adult spermatogenesis and fertility [132].

Spermiogenesis seems to be clearly affected by estrogens. In fact, after irradiation damage to the testis, estrogens are able to stimulate spermatogonial differentiation [133, 134], which is not correlated to intratesticular testosterone suppression [135]. Numerous studies have shown that estrogens are involved in the early development of spermatogenesis and are able to affect spermatogenesis. However, their role in the normal adult spermatogenesis still needs to be determined.

Activin. The majority of testicular cell types produce activins and activin-related proteins [136]. Even though they can act as hormones, they also behave as growth factors in regard to spermatogenesis. Sertoli cells and germ cells express activin receptors [136]. Culture of stem spermatogonia cells, spermatogonia and spermatocytes has shown that these cells are sensitive to activin [137–139]. Follistatin (FST) and follistatin-like 3 (FSTL3) are two activin-binding proteins that can act as antagonists to activin activity. However, overexpression of FST does not reduce local activin levels but causes infertility without clear effects on FSH levels [140]. In KO mice for FSLT3, an

increase in germ cell numbers was observed which was correlated to the increase in Sertoli cell numbers [141]. These different findings suggest that activins have probably a regulatory role in maintaining spermatogenesis and ensuring normal Sertoli cell development and activity.

Spermatogonial formation and renewal. In the mammalian testis, once the primordial germ cells migrate there during fetal development, they associate themselves with the mesenchymal cells, which will later give rise to the Sertoli cells. At this point, the sex cords are formed. The primordial germ cells then differentiate into prospermatogonia and remain centrally positioned in the cords surrounded by immature Sertoli cells. After a period of proliferation, postnatal in the rodent and prenatal in the human, the prospermatogonia migrate to the basement membrane of the sex cords to divide and form type A spermatogonia (Figure 3). Depending on the species, differences have been reported. In the human, A pale, A dark, and B spermatogonia types have been identified [142]. In the rodent testis, multiple type A spermatogonia, intermediate and type B spermatogonia have been reported and their appearance in the testis is temporally controlled. In the rat, around postnatal day 4 or 5, spermatogonial proliferation begins with type B spermatogonia identified at P6. After a series of mitotic divisions, whose mechanisms are only starting to be understood, during puberty the type B spermatogonia develop the capacity to develop into the preleptotene stage of the meiotic process.

In the testis, the migration of primordial germ cells depends on their surface expression of c-kit protein, which is the receptor for

stem cell factor (SCF), produced by the immature Sertoli cells. Mutations of the c-kit receptor will result in failure of spermatogenesis due to the absence of germ cells from the testis [143]. An upregulation of SCF mRNA on E5 has been and is concurrent with the beginning of spermatogonial division [144], which indicates an interaction between c-kit and SCF to mediate and modulate spermatogonial proliferation. Other essential growth factors in the fetus testis include transferring growth factor alpha (TGF α) [145] and neurotrophic factors [146].

SCF and c-kit have also been shown to regulate the adult testis survival of spermatogonia and spermatocytes [147]. Rat tubules cultures with FSH have been shown to influence germ cell apoptosis, affecting both mitotic and meiotic cell populations [106, 148]. GDNF and CSF1 signaling are important for spermatogonial stem cell renewal in the stem cell niche [149, 150]. Survival and developmental progression of spermatogonia depend upon expression of several genes, including the transcription factor ID4 [151] and the RNA binding protein NANOS2 [152]. Different sub-populations of spermatogonial stem cells express different genes, depending on whether that population is undergoing self-renewal, differentiation and progression, or replenishment of earlier stem cell stages [153, 154].

Spermatogenic stage germ cell development and epigenetics. Many studies have shown that the dynamics of epigenetic modifications and their regulatory networks are essential for normal spermatogenesis. Any perturbations of these epigenetic modifications is likely to cause degrees of infertility and these perturbations could result in phenotypic defects in subsequent generations [3, 155–157]. Two studies have shown that a high fat or low protein in male mice can alter the metabolic gene expression in the offspring mediated by small noncoding RNAs (sncRNAs) derived from transition ncRNAs [158, 159]. Abnormal DNA methylation is associated with altered histone modifications, dysregulation of ncRNA, abnormal protamination, and all of these contribute to male infertility. In the prospermatogonia, prepachytene piRNAs are necessary for silencing mobile elements through guiding the *de novo* DNA methylations of transposable elements in order to guarantee genome stability [160]. In late spermatocytes and round spermatids, the pachytene piRNAs could silence the retrotransposon sequences through degrading the 3'UTR of retrotransposon mRNAs or recruiting the DNMT3L to the retrotransposon locus [161]. Most studies on the different spermatogenesis stages have focused on one type of cells. In the field of infertility, histone modifications are mostly studied in the mature sperm. Our lab investigated the developmental alterations in DNA methylation during gametogenesis from PGCs to sperm. Rat fetal PGCs, prospermatogonia, spermatogonia, meiotic pachytene spermatocytes, haploid round spermatids, caput spermatozoa and mature cauda sperm were isolated and purified. Differential DNA methylation regions between each developmental stage involved were compared. The study identified a dynamic cascade of epigenetic changes during development, the most dramatic happening during the early developmental stages, which suggests complex alterations to regulate genome biology and gene expression during gametogenesis [162].

Epididymal maturation and epigenetics. Although spermatogenesis is complete with the formation of the spermatozoa following spermatogenesis, additional maturation of the sperm occurs in the epididymis [163–166]. The spermatozoa released into the seminiferous

tubules collect in the rete testes and pass through the efferent ducts into the head of the epididymis called the caput epididymis. The spermatozoa in the caput epididymis go through a further maturation as it passes through the caput to the corpus epididymis and finally, to the cauda epididymis. The caput epididymis spermatozoa do not have the capacity to have motility [167, 168]. During the transit through the epididymis, the epididymal epithelial cells produce proteins that are acquired and modify the maturation of the sperm to then in the cauda epididymis gain the capacity to become motile following ejaculation from the vas deferens where sperm are collected following epididymal maturation and stored. Therefore, the caput spermatozoa undergo a final stage of maturation during epididymal transit to the cauda epididymis to mature and gain the capacity to develop motility. The cauda epididymal sperm are then stored in the vas deferens. The molecular level maturation remains to be fully elucidated, but some aspects of epididymal maturation are known [169, 170].

Epigenetic alterations during epididymal maturation of the sperm largely remain to be elucidated [171]. Although the sperm nuclei is transcriptionally silent due to the compaction of DNA with protamines in testicular spermatogenesis, protein and epididymal components like ncRNA can be passed to the sperm and localized in the head of the sperm in the acrosome vesicle [171]. The localization of epigenetic components like ncRNA in the sperm nuclei remains to be established, but has been speculated in previous literature [172, 173]. Therefore, the role of epididymal ncRNA for sperm epididymal maturation requires further research, but is potentially an important epigenetic aspect of sperm maturation to consider [174].

Recent studies have investigated the epigenetic alterations between the caput epididymal spermatozoa and the mature cauda epididymal sperm. Environmentally induced (DDT and vinclozolin) epigenetic alterations in sperm have been shown to alter differential DNA methylation regions (DMRs) between the caput and cauda epididymal stage sperm [175, 176]. In addition, analysis of histone retention sites in the caput spermatozoa versus cauda sperm have shown differential histone retention regions (DHRs) [177]. Therefore, during epididymal maturation of sperm, there are DNA methylation and histone retention alterations that occur and are a further epigenetic developmental aspect of gamete development.

Oogenesis

Oogenesis involves the production of female gametes called eggs, and begins with the differentiation of primordial germ cells into oogonia at the period of sex determination [178]. In mammals, oogonia proliferate mitotically during fetal life to form a pool of primary oocytes that arrest in the prophase stage of the first meiotic division and stay in this state of meiotic arrest until the female reaches adulthood [179]. In the developing embryo, the nests of arrested oogonia are surrounded by somatic pre-granulosa cells [180]. Interaction and communication between oogonia and the surrounding somatic cells are vital for normal follicle and oocyte development [181, 182]. The oogonia nests subsequently break down, many oogonia undergo programmed cell death, and the pre-granulosa cells migrate to surround each remaining arrested oocyte in a process termed primordial follicle assembly [183, 184] (Figure 4). A primordial follicle is composed of a single oocyte surrounded by a single layer of flattened pre-granulosa cells. Oocytes are maintained in primordial follicles until sexual maturity, at which point follicles begin to be recruited out of the pool of primordial follicles and undergo primordial to primary follicle transition [185, 186].




Figure 4. Oogenesis and ovarian follicle stages.

Once a primordial follicle undergoes transition and begins developing, the surrounding flattened pre-granulosa cells become cuboidal and begin proliferating, themselves surrounded by the ovarian stromal cells destined to become theca cells. The follicle is now termed a primary follicle [180]. Subsequent growth and development of the follicle into secondary and pre-antral follicle stages involves continued proliferation of the granulosa cells to form multiple layers, and the initial development of a theca cell layer around the granulosa cells. Follicles with multiple layers of granulosa cells gain sensitivity to follicle stimulating hormone (FSH) secreted by the pituitary, and thus are regulated to grow and develop in cyclic waves in coordination with the estrous cycle [187]. As the granulosa and theca cell layers proliferate a fluid-filled space or antrum forms in the follicle, eventually dividing the granulosa cells into cumulus granulosa surrounding the oocyte and mural granulosa around the inside periphery of the follicle [188]. Extensive cell–cell communication and growth factor signaling between the oocyte and somatic cells occurs at all stages of oogenesis [181, 182]. Most developing follicles do not reach the stage at which ovulation occurs. Rather, follicles at several stages undergo atresia and regress [189–191], (Figure 4).

A luteinizing hormone (LH) surge from the pituitary induces ovulation in late-stage pre-ovulatory follicles, as well as promoting the resumption of meiosis in the oocyte [192]. Meiosis progresses in ovulated oocytes through the production of the first polar body, and then arrests again in metaphase two of the second meiotic division

until the time of fertilization. If fertilization occurs, then meiosis again resumes and progresses to completion with the production of the second polar body and the formation of the female pronucleus. Syngamy is the fusion of the male and female pronuclei in the newly formed zygote [193, 194]. Subsequently, the zygotic genome is activated in a carefully controlled manner to allow expression of needed genes in the newly formed individual, while suppressing expression of undesirable genes such as retrotransposons [195, 196].

Epigenetics during oogenesis and in oocytes

Information about the normal epigenetic changes that occur during oogenesis is limited. This is in part due to the difficulty of evaluating developing oocytes in ovaries, and the relatively small number of oocytes available for isolation and study compared to what can be done with male germ cells. Nonetheless, some knowledge of normal epigenomic development in female germ cells has been determined.

Primordial follicle assembly, primordial to primary follicle transition, and many subsequent stages of oogenesis have been shown to be regulated by small non-coding RNA expression, as detailed later in the *non-coding RNAs* section of *Epigenetic Programming During Gametogenesis* [197–200], (Figure 4). DNA methylation occurs at sites that are differentially imprinted between male and female gametes so that imprinted genes can be mono-allelically expressed in offspring. DNA methylation is gained gradually on imprinted genes in oocytes in developing follicles after the primary follicle stage, continuing through antral follicle stages of development [201,




Figure 5. Epigenetic mechanisms and processes (marks) (modified from [330]).

202]. In the oocytes of antral stage follicles and later, H3K4Me3 histone methylation increases. This is important for normal function in mature oocytes, and is involved with establishing the DNA methylation pattern in mature oocytes [203–206].

Using *in vitro* xenogeneic reconstituted ovaries (xrOvaries) with mouse embryonic ovarian somatic cells, Yamashiro and collaborators studied if human primordial germ cell-like cells (hPGCLC) can undergo further development [207]. They observed around 80% of genome-wide 5mC levels in human-induced pluripotent stem cells (hiPSCs) and incipient mesoderm-like cells (iMeLCs) which decreased progressively to around 20% in hPGCLC-derived cells in culture day 77 cells, then dropped at around 13% in culture day 120 cells. Based on their data, the demethylation occurred throughout the genome. Moreover, the 5mC distribution profiles of the culture days 77 to 120 cells were comparable to those observed in the oogonia and gonocytes at weeks 7 to 10. However, they were different to those seen in the blastocysts [208] and naïve human embryonic stem cells (hESCs) [207, 209]. This study demonstrated that hPGCLC-derived cells demethylate their 5mCs similarly to that of oogonia and gonocytes but not early embryonic cells and their putative *in vitro* counterparts [207].

Epigenetic programming during gametogenesis

Epigenetics

Epigenetics refers to ‘the molecular factors and processes around the DNA that regulate genome activity independent of DNA sequence, and that are mitotically stable’ [3]. These molecular processes include DNA methylation, chromatin structure, histone modifications and retention, non-coding RNAs, and RNA methylation (Figure 5). The epigenome is the complex integration of epigenetic modifications. The first epigenome analysis mapped histone acetylation and methylation in yeast [210]. These epigenetic processes and factors are central for an organism to respond to its environment with changes in gene expression. Moreover, epigenetic mechanisms are required

for a stem cell type to develop into a differentiated cell type, which make them an integral part of normal biology [3, 211, 212].

DNA methylation and histone modifications

De novo DNA methylation in males restarts in prospermatogonia at E14.5 in mice and is fully established at birth [213, 214]. In both female and male germlines, the factors responsible for mediating DNA methylation DNMT3A or 3B and DNMT3L at imprinted loci were identified [215–217]. In the gametes, the sequence identity and the characteristics of imprinted regions are now well characterized; however, the mechanisms targeting the *de novo* methyltransferases to imprinted regions remain to be further investigated [218]. In oocytes, the establishment of DNA methylation at imprinted or retrotransposon-rich sites occurs gradually in growing follicles subsequent to the primary follicle stage (reviewed in [201, 202]).

During meiosis of the gametogenesis process, germ cells stop their progress at the prophase stage to allow parental genomes to exchange genetic information through meiosis recombination [219]. At this stage, chromosomes pair in a homologous manner, and large pieces of the chromosomes can be exchanged through crossover events [219, 220]. These crossover events are necessary to maintain euploidy in gametes. An absence of crossover events has been linked to infertility and aneuploidy in the offspring [221, 222]. These meiotic crossover events happen at genomic hotspots and are enriched in regions outside of promoters that bear histone H3K4me3 peaks and established by Prdm9 [223–228]. Mutations in enzymes involved in histone posttranslational modifications observed in meiosis have been shown to have an impact (decrease or increase) on the DNA double-strand break activity, which suggests a role for histone modifications in the initiation and/or repair activity [229–232]. In oocytes, H3K4Me3 histone methylation increases from the antral follicle stage onward, and is important for meiotic recombination, oocyte maturation, oocyte transcriptional activity, and for the establishment of a normal DNA methylation pattern in mature oocytes [203–206, 233, 234].

Chromatin structure & histones

The chromatin reorganization during meiosis is largely transient. The most extensive modifications in chromatin state, structure, or composition occurs after male and female meiosis [220]. In the sperm, the vast majority of histones are replaced with sperm-specific nuclear proteins called protamines [235]. This process is facilitated by different steps: hyperacetylation of histones in round spermatids believed to weaken the interactions between histones and DNA, this will enable the eviction and replacement of histones by testes-specific histone variants, then by transition proteins to end with protamines [236–238]. Because of their endonuclease-inaccessible toroid structure, protamines manage to package the sperm DNA into a tenfold more compact structure than the heterochromatin found in somatic nuclei [239, 240]. The retained histones in the sperm were believed to be remnants of incomplete histone-to-protamine replacement, but recent studies have demonstrated that these retained histones are present at key developmental gene promoters/enhancers in mature sperm. These retained histones bear both active or repressive histone modification [241, 242].

This programmatic retention and evolutionary conservation of histone localization suggests that epigenetic information can be passed through the paternal lineage. Moreover, alteration in histone levels, or chromatin regulators involved in spermatogenesis leads to developmental defects, which can be passed on to the subsequent generation [243, 244]. Altogether, these studies imply that retained histones serve as molecular carriers of epigenetic memory; however, the mechanisms are yet to be elucidated.

Non-coding RNAs

DNA is not the only means to transmit the information between generations. Non-coding RNAs (ncRNAs) are regulatory elements of gene expression and chromatin structure [245]. The differential susceptibility to these non-coding RNAs contributes to tissue-specific gene expression. Early on, ncRNAs are important in the germline development, but they are also crucial players in posttranscriptional gene control during spermatogenesis and oogenesis. Different classes of ncRNAs exist, but this section will focus on microRNAs (miRNAs), Piwi interacting RNAs (piRNAs), and long non-coding RNAs (lncRNAs) and their role in the PGCs and the gametes.

miRNAs

During the PGC specification, some miRNAs are selectively expressed such as miR-10b, -18a, -93, -106b, -126-3p, -127, -181a, -181b, and -301. All of them have important functions in these cells such as differentiation, migration, and apoptosis in PGCs in mice [246]. For instance, Medeiros and collaborators have shown that in mice a deficiency in miR-290-295 cluster result in an abnormal germ cell with defect in the PGC migration [247]. In the female, an upregulation of miR-29b has been shown to induce PGC development by targeting DNA methyltransferases Dnmt3a and Dnmt3b [248]. In the zebrafish, miR-202-5p has been identified as a potential germ plasma-specific biomarker due to its potential role in the germ cell development [249]. Other microRNAs have been linked to PGC migration in the zebrafish as well, such as miR-430, which regulates *sdfla* and *cxc7* mRNAs key transcripts regulating migration [250].

In the early stages of spermatogenesis, different miRNAs have been described in mammals as being crucial for germ cell self-renewal and differentiation. miR-34c has been identified as promoting mouse spermatogonial stem cell (SSCs) differentiation by

targeting Nanos2 [251]. Moreover, this miRNA has another role in the later stage of spermatogenesis where miR-34c is involved in apoptotic events of spermatocytes and round spermatids [252], and also in the NOTCH signaling, which is important in the control of germ cell differentiation [253]. A list of miRNAs involved in cell cycle regulation have been identified such as miR-293, 291a-5p, 290-5p and 294 [254]. Other miRNAs are involved in later stages of spermatogenesis. The Let-7 miR family is involved in the mouse spermatogonial differentiation, especially in the maturation of undifferentiated spermatogonia to A1 spermatogonia by suppressing Lin28 [255]. In contrast, some miRNAs such as miR-146 play a crucial role in keeping spermatogonia in an undifferentiated state in the mouse [256]. Other miRNAs play a role in the regulation of meiotic and postmeiotic events in the later stages of spermatogenesis such as the miR-449 cluster. During murine spermatogenesis, the upregulation of miR-449 cluster is crucial for the initiation of meiosis [257]. These miRNAs by targeting BCL2 and AFT1 are involved in germ cell apoptosis [258].

piRNAs

Another class of sncRNAs, piRNAs have been discovered in the germline. Their role is to safeguard the germline genome from retrotransposons and protect the genomic stability [259, 260]. These piRNAs are believed to be involved in pathway components of DNA methylation remodeling during early PGC specification in mammals [209]. Moreover, a loss of Piwi function in mice or zebrafish results in a decrease of germ cells by apoptosis, this underlying its role in germ cell maintenance [261].

lncRNAs

The role of long non-coding RNAs in PGC specification has not been described. Some researchers suggest their possible roles in controlling transcription factors such as BLIMP1/PRDM1 or DAZL [262, 263]. In fact, more than 300 binding sites of BLIMP1/PRDM1 in the murine PGCs are associated with non-coding genes whose functions in PGCs specification are still unknown [23, 262]. The lncRNA-Tcam1 and lncRNA-HSVIII have a crucial role in pachytene spermatocytes, which implies their potential participation in the transcriptional regulation of spermatocyte-specific gene expression [264]. lncRNAs have been linked to functions related to post-transcriptional control during spermatogenesis, such as tubulin cofactor A (TBCA), which has the ability to interact with tubulin during the microtubule rearrangement process [265]. However, most studies have been conducted in rats, so not much is known about lncRNAs in the human. In human spermatogenesis, male infertility has been associated with NLC1-C through the control of miRNA expression via RNA-binding proteins [266].

ncRNAs in oocytes

Most of the studies have focused on miRNAs in oogenesis and ovary function using conditional KO mice models to evaluate their involvement in the ovary. By using this approach, a clear role has been outlined for miRNAs in folliculogenesis, oocyte maturation, and ovulation. Other miRNAs are also involved in the assembly of primordial follicles, the transition from primordial to primary follicles, follicular growth, oocyte maturation, ovulation, and the formation of the *corpus luteum* in mammals [198, 199, 267–269]. After a conditional knockout of Dicer1 from follicular granulosa cells in mammals, abnormal oocyte maturation, disrupted follicular

ENVIRONMENTALLY INDUCED EPIGENETIC TRANSGENERATIONAL INHERITANCE

Environmental Toxicants

Vinclozolin (Agricultural Fungicide)

Methoxychlor (Agricultural Pesticide)

Dioxin/TCDD (Industrial Contaminant)

Plastic Compounds (BPA & Phthalates)

Mercury

Permethrin & DEET (Insect Repellants)

DDT (Pesticide)

Tributyltin (Industrial Toxicant & Biocide)

Hydrocarbons (Jet Fuel)

Atrazine

Other Types Exposures

Nutrition (High Fat or Caloric Restriction)

Temperature & Drought (Plant Health & Flowering)

Smoking & Alcohol

Stress (Behavioral)



Plants



Flies



Worms



Fish



Bird



Rodents



Pigs



Humans

Figure 6. Environmentally induced epigenetic transgenerational inheritance. Various exposures and species investigated (modified from [156]).

development and ovulation, increased follicular atresia, and infertility were reported [270–272]. Other studies have demonstrated miRNAs involvement in granulosa cells proliferation, survival, terminal differentiation, steroidogenesis, and cumulus expansion [200, 273–281]. The overexpression of miR-143 in murine 15.5 dpc ovaries has been shown to repress the formation of primordial follicles by stopping the proliferation of pre-granulosa cells. An increased number of primordial follicles were observed in transfected 18.5 dpc ovaries with miR-376a (reviewed by Grossman and Shalgi [197]).

Environmental toxicant exposures resulting in epigenetic changes in gametes

In addition to epigenetic changes being a part of the normal developmental process for gametes, it is also possible that exposure to environmental factors will induce abnormal epigenetic changes to the germ cell epigenome [282, 283]. Such changes may be heritable and affect the phenotype of subsequent generations [156, 284]. Exposure of male mice to the endocrine disruptor bisphenol A (BPA) induces changes to DNA methylation in the fetal germ cells of their developing offspring [285]. Primordial germ cells are also induced to alter DNA methylation in response to hypoglycemic conditions in the uterus [286] or from exposure to the agricultural fungicide vinclozolin [287]. Exposure to a wide variety of environmental factors can lead to DNA methylation changes in spermatozoa and mature sperm [288]. In rodents, direct exposure to arsenic [289], the fungal toxin zearalenone [290], the plastics compounds bisphenol A (i.e. BPA) [291] and phthalates [292], the agricultural fungicide vinclozolin [293], the pesticide dichlorodiphenyltrichloroethane (DDT) [294], and the herbicides glyphosate [295] and atrazine [296], all induce sperm DNA methylation changes. In humans, studies have shown that environmental factors such as exposure to phthalates [297], alcohol [298], flame retardants [299, 300], chemotherapy treatment [301], obesity [302], and exercise [303] are correlated to sperm DNA methylation changes (Figure 6). In one fish species, exposure

to BPA resulted in changes to oocyte DNA methylation in the next generation [304], (Table 1).

Epigenetic changes to histones in germ cells can occur after exposure to environmental factors. For example, zebrafish exposed to BPA showed decreased sperm histone acetylation, as well as impaired primordial germ cell migration, although these findings were not associated with decreased fertility [305]. BPA exposure in a minnow species resulted in changes in oocyte histone methylation in the offspring [304]. Pubertal exposure of mice to the fungicides carbendazim and chlorothalonil caused changes in H3K9me3 levels in sperm [306]. Pubertal exposure to the fungal toxin zearalenone altered mouse histone H3K27 methylation [290]. In mice, exposure to the pesticide chlordecone resulted in altered levels of H3K4Me3 in developing testes [307]. Exposure to chlordecone in mice also resulted in changes in H3K4me3 and H4ac in mature oocytes [308]. Even exposure to chronic restraint stress can alter histone acetylation, methylation and phosphorylation in germinal vesicle-stage oocytes [309].

Another way in which environmental factors can alter histones in sperm is to affect which histones are retained as male germ cells develop. As male germ cells undergo spermiogenesis, most of the histones associated with the DNA are replaced by protamines [236, 310]. Protamines help condense and package DNA into the small sperm head. However, some histones are retained, and they are often located near developmental regulatory genes that are expressed early in embryonic development [311]. Exposure to environmental toxicants has been shown to alter retention of histones in sperm [312]. Men exposed to either cigarette smoke [313] or the smoke of surrounding fires [314] have been shown to have an altered ratio of histones to protamines in sperm. *In utero* exposure to caloric restriction in mice has also been shown to alter histone retention in sperm [315]. In transgenerational studies in rats, it was found that exposure of gestating female F0 generation rats to either DDT or vinclozolin resulted in changes in histone retention in the subsequent transgenerational F3 generation, but interestingly not in

Table 1. Environmental exposures resulting in epigenetic changes in gametes.

| Environmental exposure | Epigenetic change | Cell type | Reference |
|--------------------------------|---|-------------------|---|
| BPA | DNA methylation | Fetal germ cells | Zhang et al. (2012) [285] |
| Uterine hypoglycemia | DNA methylation | PGCs | Ren et al. (2018) [286] |
| Arsenic | DNA methylation | Sperm | Nohara et al. (2019) [289] |
| Zearalenone | DNA methylation | Sperm | Gao et al. (2019) [290] |
| BPA | DNA methylation | Sperm | Rahman et al. (2020) [291] |
| Phthalates | DNA methylation | Sperm | Prados et al. (2015) [292] |
| Vinclozolin | DNA methylation | Sperm | Beck et al. (2017) [293] |
| DDT | DNA methylation | Sperm | Skinner et al. (2018) [294] |
| Glyphosate | DNA methylation | Sperm | Kubsad et al. (2019) [295] |
| Atrazine | DNA methylation | Sperm | McBirney et al. (2017) [296] |
| BPA | DNA methylation | Fish oocyte | Zhu et al. (2020) [304] |
| Phthalates | DNA methylation | Human sperm | Wu et al. (2017) [297] |
| Alcohol | DNA methylation | Human sperm | Ouko et al. (2009) [298] |
| Flame retardants | DNA methylation | Human sperm | Soubry et al. (2017), Greeson et al. (2020) [299, 300] |
| Chemotherapy | DNA methylation | Human sperm | Shnorhavorian et al. (2017) [301] |
| Obesity | DNA methylation | Human sperm | Soubry et al. (2016) [302] |
| Exercise | DNA methylation | Human sperm | Denham et al. (2015) [303] |
| BPA | Histone acetylation | Fish sperm | Lombo et al. (2019) [305] |
| BPA | histone methylation | Fish oocyte | Zhu et al. (2020) [304] |
| Carbendazim and chlorothalonil | Histone H3K9me3 | Sperm | Li et al. (2018) [306] |
| Zearalenone | Histone H3K27 methylation | Sperm | Gao et al. (2019) [290] |
| Chlordecone | Histone H3K4Me3 | Developing testes | Gely-Pernot et al. (2018) [307] |
| Chlordecone | Histone H3K4Me3 | Oocytes | Legoff et al. (2019) [308] |
| Restraint stress | Histone acetylation, methylation, phosphorylation | Oocytes | Wu et al. (2015) [309] |
| Cigarette smoke | Histone retention | Human sperm | Hamad et al. (2014) [313] |
| Smoke | Histone retention | Human sperm | Lettieri et al. (2020) [314] |
| Caloric restriction in utero | Histone retention and DNA methylation | Sperm | Radford et al. (2014) [315] |
| DDT | DNA methylation, non-coding RNA expression, and histone retention | Sperm | Skinner et al. (2018) [294] |
| Vinclozolin | DNA methylation, non-coding RNA expression, and histone retention | Sperm | Ben Maamar et al. (2018) [316] |
| Vinclozolin | miRNA | PGCs | Brieno-Enriquez et al. (2015) [317] |
| Early life trauma | miRNA and lncRNA | Sperm | Dickson et al. (2018), Gapp et al. (2014, 2020) [318–320] |
| Early life stress | miRNA | Human sperm | Dickson et al. (2018) [318] |
| Smoking | miRNA | Human sperm | Marczylo et al. (2012) [321] |
| Obesity | miRNA | Human sperm | Lopez et al. (2018), Donkin et al. (2015) [322, 323] |
| Bariatric surgery | miRNA | Human sperm | Donkin et al. (2015) [323] |

the intervening F1 and F2 generations [294, 312, 316]. However, F1, F2 and F3 exposure-lineage generations all showed changes in DNA methylation and non-coding RNA expression (Table 1).

The expression of non-coding RNA (ncRNA) in germ cells is another epigenetic factor that can be responsive to environmental factors. Mice exposed to vinclozolin *in utero* exhibited changes in micro-RNAs in their primordial germ cells [317]. Rats exposed to either vinclozolin or DDT *in utero* have been shown to have altered levels of piwi-interacting RNAs (piRNAs) and small tRNA fragments in sperm upon reaching adulthood [294, 316]. Traumatic stress can also be an environmental factor that causes epigenetic changes in sperm. In mouse models of early life trauma, changes in miRNAs and long non-coding RNAs were seen in sperm from exposed males [318–320]. Interestingly, the behavioral and metabolic alterations seen in the resulting offspring were

recapitulated by injection of sperm RNAs from traumatized males into fertilized wild-type oocytes [319]. In humans, changes in miRNA expression in sperm have been seen after exposure to early life stress [318], smoking [321], obesity [322, 323], and bariatric surgery [323] (Table 1).

Epigenetic transgenerational inheritance

Exposure to environmental factors, such as toxicants, can induce epigenetic changes in germ cells that can affect the subsequent generations. These epimutation changes are brought to the next generation at fertilization and have the possibility of altering gene expression and phenotype in the developing embryo. Since the germ cell epimutations can affect the earliest stem cells formed in the




Figure 7. Environmentally induced transgenerational epigenetic inheritance. Schematic of multigenerational versus transgenerational environmental exposures (modified from [324]).

embryo, then any subsequent cell type in the embryo and adult animal may have epimutations and changes in gene expression that could affect their phenotype [156].

Epigenetic transgenerational inheritance is defined as germline-mediated inheritance of epigenetic information between generations in the absence of continued direct environmental influences that leads to phenotypic variation [3]. When a male or a non-pregnant female is exposed to an environmental factor that can induce epigenetic change, then epimutations can arise in that individual (the F0 generation), and that individual's germ cells. The germ cells that contribute to forming the next F1 generation were directly exposed to the environmental factor, so epigenetic and phenotypic changes seen in the F1 generation are an example of direct multigenerational exposure, but not transgenerational inheritance [324], (Figure 7). If these F1 generation animals pass on epigenetic and phenotypic changes to the unexposed F2 generation, then this is an example of epigenetic transgenerational inheritance. Similarly, if a pregnant female is exposed to an environmental factor such as a toxicant, the developing F1 generation embryo is directly exposed, and the germ cells in that embryo that go on to form the F2 generation are also directly exposed. Epigenetic and phenotypic changes would have to be seen in subsequent F3 generation or later generations for this to be an example of epigenetic transgenerational inheritance (Figure 7, Table 2) [156, 157, 324].

Epimutations have been reported in transgenerational generation gametes after ancestral exposure to a wide variety of environmental stressors including toxicants [156] and psychological stress [325]. This phenomenon has been observed in several organisms including fish [326] and rodents [157] (reviewed in [156]) (Table 2). Epimutations have been reported in transgenerational F3 generation rat sperm after ancestral exposure to several toxicants [156]. One question that occurs is at what stage of germ cell development do these sperm epimutations arise, or are they continuously present? Two recent studies have addressed this question. The first examined differences in DNA methylation compared to controls after ancestral exposure to DDT [176]. The number of differential DNA methylation regions (DMRs) was determined for the spermatogenic stages of primordial germ cells, prospermatogonia,

spermatogonia, pachytene spermatocytes, round spermatids, caput epididymal spermatozoa, and cauda epididymal sperm. It was found that, of the 265 DMRs present in cauda epididymal sperm, 26% arose in the prospermatogonia stage, 25% in spermatogonia, 18% in pachytene spermatocytes, 5% in round spermatids, 12% in caput epididymal spermatozoa, and 14% only arose in cauda epididymal sperm (Figure 3). Similar observations were made with ancestral vinclozolin exposure [327]. This shows that transgenerational changes in DNA methylation arise throughout gametogenesis.

Similarly, another study examined transgenerational differences in histone retention in sperm from rats ancestrally exposed to DDT or vinclozolin [177]. In mature sperm nuclei, most histone proteins are replaced by protamines to facilitate packaging the DNA into the small sperm head [328]. However, some histones are retained, sometimes near genes that are important to early embryo development [243, 329]. In this study [177], the sites of differential histone retention (DHR) were determined for the spermatogenic stages of round spermatids, caput epididymal spermatozoa, and cauda epididymal sperm. It was found that, of those DHRs that were present in cauda epididymal sperm, about 50% were present in round spermatids, very few arose in caput epididymal spermatozoa, and 40–50% arose at the cauda epididymal sperm stage. Again, this shows that an environmentally induced cascade of histone retention changes occurs during different stages of spermatogenic development.

Recently, two studies investigated the integration of DMR, ncRNA, and DHR by overlapping the different genomic sites between the different epimutations. Rats were ancestrally exposed to DDT or vinclozolin and the F1, F2, and F3 generations sperm epimutations examined. The chromosomal locations of the DMR, ncRNA, and DHR epimutations were distinct with few overlapping, but present in the same regions with similar genomic features such as CpG deserts. Comparing the F1, F2, and F3 generations provided insights into the integration of the different epimutations and the direct exposure versus transgenerational impacts of exposures. Interestingly, the effects observed in the F3 generation were different than those observed in the direct exposure F1 and F2 generations. Direct exposure impacted the ncRNA in the F1 generation, but to a much lesser extent in the F2 and F3 generations.

Table 2. Environmental exposures promoting epigenetic transgenerational inheritance.

| Environmental exposure | Epigenetic change | Cell type | Reference |
|----------------------------|---|--------------------------------------|----------------------------------|
| Several toxicants (review) | DNA methylation | Sperm | Nilsson et al. (2018) [156] |
| DDT | DNA methylation | Sperm | Ben Maamar et al. (2019) [176] |
| DDT | DNA methylation, non-coding RNA, and histone retention | Sperm | Skinner et al. (2018) [294] |
| Vinclozolin | DNA methylation, non-coding RNA expression, and histone retention | Sperm | Ben Maamar et al. (2018) [316] |
| Vinclozolin | DNA methylation | Sperm | Anway et al. (2005) [157] |
| DDT or vinclozolin | Histone retention | Sperm | Ben Maamar et al. (2020) [177] |
| Glyphosate | DNA methylation | Sperm | Kubsad et al. (2019) [295] |
| Atrazine | DNA methylation | Sperm | McBirney et al. (2017) [296] |
| Methoxychlor | DNA methylation | Sperm | Manikkam et al. (2014) [331] |
| Glyphosate | DNA methylation and histone retention | Sperm | Ben Maamar et al. (2020) [332] |
| Dioxin | DNA methylation | Sperm | Manikkam et al. (2012) [333] |
| BPA | DNA methylation | Sperm | Rahman et al. (2020) [291] |
| Phthalates | DNA methylation | Sperm | Prados et al. (2015) [292] |
| Jet fuel | DNA methylation | Sperm | Manikkam et al. (2012) [334] |
| Vinclozolin | tRNA halves | Sperm | Schuster et al. (2016) [335] |
| Methylmercury | DNA methylation | Fish sperm | Carvan et al. (2017) [326] |
| Nutrition change | Gene promoter methylation | Pig liver | Braunschweig et al. (2012) [336] |
| Famine | DNA methylation | Human blood cells | Jiang et al. (2020) [337] |
| Genetic manipulation | Histone modifications | <i>Drosophila</i> embryos | Xia et al. (2016) [338] |
| Genetic manipulation | Histone modifications | <i>Caenorhabditis elegans</i> larvae | Kelly et al. (2014) [339] |
| Vinclozolin | tRNA halves | Sperm | Schuster et al. (2016) [335] |

Table 3. Future research in gametogenesis.

| | |
|----|--|
| 1) | Examine all stages PGCs to gametes for epigenetic and genetic transitions. |
| 2) | Examine and investigate all the epigenetic processes (DNA methylation, histones, ncRNA, and chromatin structure) for the epigenetic regulation of gametogenesis. |
| 3) | Use systems biology and genome-wide analyses to integrate the epigenetics and genetics of the gametogenesis process. |
| 4) | Incorporate environmentally induced epigenetic transgenerational inheritance into our research and understanding of generational impacts of altering gamete epigenetics. |

DMR were predominant in the F1, F2, and F3 generations, whereas the DHRs were observed primarily in the F3 generation. Observations suggest a potential role for ncRNA-directed DNA methylation and DNA methylation-directed histone retention in the environmentally induced epigenetic transgenerational inheritance of altered gametogenesis [294, 316].

Conclusions and future research

Normal fertilization and early embryonic development require the integration of genetic and epigenetic processes in the gametes. Few developmental or physiological processes will not require the integration of these two distinct molecular mechanisms. Understanding the underlying processes and mechanisms *in vivo* will help us better understand the gametogenesis and formation of the sperm and egg, as well as pathologies such as idiopathic infertility. The past few decades have advanced our knowledge of primordial germ cell and gamete development. This has provided insights into the molecular mechanism involved in related disorders, such as infertility and birth defects. These findings are also promising in the field of reproductive therapies.

Although a genetic focus can identify the genes involved in a developmental process such as gametogenesis, an understanding of the epigenetics is required to clarify how the gene expression is controlled and environmental factors regulate the developmental process. The DNA methylation, histone modifications, ncRNA, and chromatin structure are the epigenetic mechanisms required to regulate the gene expression. Therefore, no development or biological process can be understood without the integration of epigenetics and genetics. This is the case for normal developmental processes like spermatogenesis and oogenesis, as well as abnormal processes involved in pathology. The ability of environmental factors such as nutrition, toxicants, or stress to impact developmental processes like gametogenesis involves alterations in the epigenetics to then impact the basic genetics and gene expression. Since the gametogenesis process produces the gametes, sperm and eggs, alterations in gametogenesis impact the future generations physiology, phenotype, and gametes. This non-genetic form of inheritance termed epigenetic transgenerational inheritance requires manipulation during gametogenesis to obtain epigenetically modified sperm and eggs. Therefore, the current review provides the basic information on the gametogenesis process, epigenetic regulation of the developmental process, and impacts on epigenetic transgenerational

inheritance. As the information on epigenetic regulation of gametogenesis is a new area of research, future research is needed to expand our understanding of the integration of epigenetics and genetics in gametogenesis.

The future research suggested includes the following, Table 3. A need to examine all the gametogenesis stages from PGCs to gametes to understand the integration and transition between the different stages of development. A need to integrate all the epigenetic processes and gene expression for a more complete understanding of the epigenetic regulation of gametogenesis. Emphasis on genome-wide analysis and system biology approaches in the study of gametogenesis to fully understand the developmental process and associated pathology. A need for a generational context, to incorporate the environmentally induced epigenetic transgenerational inheritance of gametogenesis alterations into our understanding of gamete development. Without these more complete systems biology approaches, the progress of research and impacts will be diminished.

Supplementary material

Supplementary material is available at *BIOLRE* online.

Acknowledgements

We thank Dr. Jennifer L.M. Thorson for critical review of the manuscript. We thank Dr. Stephanie King for the drawings of the stages of spermatogenesis and oogenesis. We acknowledge Ms. Amanda Quilty for assistance in editing the manuscript, and Ms. Heather Johnson for assistance in preparation of the manuscript.

Conflict of interest: The authors declare no conflict of interest.

Abbreviations

Primordial germ cells (PGCs)
 Bone morphogenetic protein (BMP).
 Knock-out (KO).
 Stromal cell-derived factor 1 (SDF-1).
 Chemokine receptor type 4 (CXCR4).
 5-methylcytosine (5mC).
 Germline reprogramming responsive (GRR).
 Ten eleven translocation (TET).
 Follicle stimulating hormone (FSH).
 Androgen receptors (ARs).
 Ethanedimethane sulfonate treatment (EDS).
 Follistatin (FST).
 Follistatin-like 3 (FSTL3).
 Stem cell factor (SCF).
 Luteinizing hormone (LH).
 Non-coding RNAs (ncRNAs).
 MicroRNAs (miRNAs).
 Piwi interacting RNAs (piRNAs).
 Long non-coding RNAs (lncRNAs).
 Tubulin cofactor A (TBCA).
 Bisphenol A (BPA).
 Differential DNA methylation regions (DMRs).
 Differential histone retention sites (DHRs).

Author Contributions

MBM—Writing and editing manuscript.

EN—Writing and editing manuscript.

MKS—Funding acquisition, writing and editing manuscript.

References

1. Tang WW, Kobayashi T, Irie N, Dietmann S, Surani MA. Specification and epigenetic programming of the human germ line. *Nat Rev Genet* 2016; 17:585–600.
2. Ohinata Y, Ohta H, Shigeta M, Yamanaka K, Wakayama T, Saitou M. A signaling principle for the specification of the germ cell lineage in mice. *Cell* 2009; 137:571–584.
3. Skinner MK. Environmental epigenetic transgenerational inheritance and somatic epigenetic mitotic stability. *Epigenetics* 2011; 6:838–842.
4. Seydoux G, Dunn MA. Transcriptionally repressed germ cells lack a subpopulation of phosphorylated RNA polymerase II in early embryos of *Caenorhabditis elegans* and *Drosophila melanogaster*. *Development* 1997; 124:2191–2201.
5. Boswell RE, Mahowald AP. Tudor, a gene required for assembly of the germ plasm in *Drosophila melanogaster*. *Cell* 1985; 43:97–104.
6. Capowski EE, Martin P, Garvin C, Strome S. Identification of grandchildless loci whose products are required for normal germ-line development in the nematode *Caenorhabditis elegans*. *Genetics* 1991; 129:1061–1072.
7. Christerson LB, McKearin DM. Orb is required for anteroposterior and dorsoventral patterning during *drosophila* oogenesis. *Genes Dev* 1994; 8:614–628.
8. Matova N, Cooley L. Comparative aspects of animal oogenesis. *Dev Biol* 2001; 231:291–320.
9. Simpson AJ, Caballero OL, Jungbluth A, Chen YT, Old LJ. Cancer/testis antigens, gametogenesis and cancer. *Nat Rev Cancer* 2005; 5:615–625.
10. Leitch HG, Smith A. The mammalian germline as a pluripotency cycle. *Development* 2013; 140:2495–2501.
11. Lawson KA, Dunn NR, Roelen BA, Zeinstra LM, Davis AM, Wright CV, Korving JP, Hogan BL. Bmp4 is required for the generation of primordial germ cells in the mouse embryo. *Genes Dev* 1999; 13:424–436.
12. Saitou M, Barton SC, Surani MA. A molecular programme for the specification of germ cell fate in mice. *Nature* 2002; 418:293–300.
13. Saitou M, Yamaji M. Primordial germ cells in mice. *Cold Spring Harb Perspect Biol* 2012; 4:a008375.
14. Tremblay KD, Dunn NR, Robertson EJ. Mouse embryos lacking Smad1 signals display defects in extra-embryonic tissues and germ cell formation. *Development* 2001; 128:3609–3621.
15. Ying Y, Liu XM, Marble A, Lawson KA, Zhao GQ. Requirement of bmp 8b for the generation of primordial germ cells in the mouse. *Mol Endocrinol* 2000; 14:1053–1063.
16. Ying Y, Zhao GQ. Cooperation of endoderm-derived BMP2 and extraembryonic ectoderm-derived BMP4 in primordial germ cell generation in the mouse. *Dev Biol* 2001; 232:484–492.
17. Saffman EE, Lasko P. Germline development in vertebrates and invertebrates. *Cell Mol Life Sci* 1999; 55:1141–1163.
18. Gomperts M, Garcia-Castro M, Wylie C, Heasman J. Interactions between primordial germ cells play a role in their migration in mouse embryos. *Development* 1994; 120:135–141.
19. Ohinata Y, Payer B, O'Carroll D, Ancelin K, Ono Y, Sano M, Barton SC, Obukhanych T, Nussenzweig M, Tarakhovskiy A, Saitou M, Surani MA. Blimp1 is a critical determinant of the germ cell lineage in mice. *Nature* 2005; 436:207–213.
20. Yamaji M, Seki Y, Kurimoto K, Yabuta Y, Yuasa M, Shigeta M, Yamanaka K, Ohinata Y, Saitou M. Critical function of Prdm14 for the establishment of the germ cell lineage in mice. *Nat Genet* 2008; 40:1016–1022.
21. Weber S, Eckert D, Nettersheim D, Gillis AJ, Schafer S, Kuckenbergh P, Ehlermann J, Werling U, Biermann K, Looijenga LH, Schorle H.

- Critical function of AP-2 gamma/TCFAP2C in mouse embryonic germ cell maintenance. *Biol Reprod* 2010; 82:214–223.
22. Magnusdottir E, Dietmann S, Murakami K, Gunesdogan U, Tang F, Bao S, Diamanti E, Lao K, Gottgens B, Azim Surani M. A tripartite transcription factor network regulates primordial germ cell specification in mice. *Nat Cell Biol* 2013; 15:905–915.
 23. Magnusdottir E, Surani MA. How to make a primordial germ cell. *Development* 2014; 141:245–252.
 24. Nakaki F, Hayashi K, Ohta H, Kurimoto K, Yabuta Y, Saitou M. Induction of mouse germ-cell fate by transcription factors in vitro. *Nature* 2013; 501:222–226.
 25. Ancelin K, Lange UC, Hajkova P, Schneider R, Bannister AJ, Kouzarides T, Surani MA. Blimp1 associates with Prmt5 and directs histone arginine methylation in mouse germ cells. *Nat Cell Biol* 2006; 8:623–630.
 26. Hayashi K, de Sousa Lopes SM, Surani MA. Germ cell specification in mice. *Science* 2007; 316:394–396.
 27. Richardson BE, Lehmann R. Mechanisms guiding primordial germ cell migration: Strategies from different organisms. *Nat Rev Mol Cell Biol* 2010; 11:37–49.
 28. Vincent SD, Dunn NR, Sciammas R, Shapiro-Shalef M, Davis MM, Calame K, Bikoff EK, Robertson EJ. The zinc finger transcriptional repressor Blimp1/Prdm1 is dispensable for early axis formation but is required for specification of primordial germ cells in the mouse. *Development* 2005; 132:1315–1325.
 29. Gunesdogan U, Magnusdottir E, Surani MA. Primordial germ cell specification: A context-dependent cellular differentiation event [corrected]. *Philos Trans R Soc Lond B Biol Sci* 2014; 369:20130543.
 30. Kurimoto K, Yabuta Y, Ohinata Y, Shigetani M, Yamanaka K, Saitou M. Complex genome-wide transcription dynamics orchestrated by Blimp1 for the specification of the germ cell lineage in mice. *Genes Dev* 2008; 22:1617–1635.
 31. Fox N, Damjanov I, Martinez-Hernandez A, Knowles BB, Solter D. Immunohistochemical localization of the early embryonic antigen (SSEA-1) in postimplantation mouse embryos and fetal and adult tissues. *Dev Biol* 1981; 83:391–398.
 32. Ginsburg M, Snow MH, McLaren A. Primordial germ cells in the mouse embryo during gastrulation. *Development* 1990; 110:521–528.
 33. Sato M, Kimura T, Kurokawa K, Fujita Y, Abe K, Masuhara M, Yasunaga T, Ryo A, Yamamoto M, Nakano T. Identification of PGC7, a new gene expressed specifically in preimplantation embryos and germ cells. *Mech Dev* 2002; 113:91–94.
 34. Tam PP, Zhou SX. The allocation of epiblast cells to ectodermal and germ-line lineages is influenced by the position of the cells in the gastrulating mouse embryo. *Dev Biol* 1996; 178:124–132.
 35. Chambers I, Silva J, Colby D, Nichols J, Nijmeijer B, Robertson M, Vrana J, Jones K, Grotewold L, Smith A. Nanog safeguards pluripotency and mediates germline development. *Nature* 2007; 450:1230–1234.
 36. Avilion AA, Nicolis SK, Pevny LH, Perez L, Vivian N, Lovell-Badge R. Multipotent cell lineages in early mouse development depend on SOX2 function. *Genes Dev* 2003; 17:126–140.
 37. Scholer HR, Ruppert S, Suzuki N, Chowdhury K, Gruss P. New type of POU domain in germ line-specific protein Oct-4. *Nature* 1990; 344:435–439.
 38. Yamaguchi S, Kimura H, Tada M, Nakatsuji N, Tada T. Nanog expression in mouse germ cell development. *Gene Expr Patterns* 2005; 5:639–646.
 39. Sybirna A, Tang WWC, Pierson Smela M, Dietmann S, Gruhn WH, Brosh R, Surani MA. A critical role of PRDM14 in human primordial germ cell fate revealed by inducible degrons. *Nat Commun* 2020; 11:1282.
 40. Irie N, Weinberger L, Tang WW, Kobayashi T, Viukov S, Manor YS, Dietmann S, Hanna JH, Surani MA. SOX17 is a critical specifier of human primordial germ cell fate. *Cell* 2015; 160:253–268.
 41. Kojima Y, Sasaki K, Yokobayashi S, Sakai Y, Nakamura T, Yabuta Y, Nakaki F, Nagaoka S, Woltjen K, Hotta A, Yamamoto T, Saitou M. Evolutionarily distinctive transcriptional and signaling programs drive human germ cell lineage specification from pluripotent stem cells. *Cell Stem Cell* 2017; 21:517–532 e515.
 42. Anderson R, Copeland TK, Scholer H, Heasman J, Wylie C. The onset of germ cell migration in the mouse embryo. *Mech Dev* 2000; 91:61–68.
 43. Molyneux KA, Stallock J, Schaible K, Wylie C. Time-lapse analysis of living mouse germ cell migration. *Dev Biol* 2001; 240:488–498.
 44. Ewen K, Baker M, Wilhelm D, Aitken RJ, Koopman P. Global survey of protein expression during gonadal sex determination in mice. *Mol Cell Proteomics* 2009; 8:2624–2641.
 45. Gu Y, Runyan C, Shoemaker A, Surani A, Wylie C. Steel factor controls primordial germ cell survival and motility from the time of their specification in the allantois, and provides a continuous niche throughout their migration. *Development* 2009; 136:1295–1303.
 46. Ohta H, Tohda A, Nishimune Y. Proliferation and differentiation of spermatogonial stem cells in the w/wv mutant mouse testis. *Biol Reprod* 2003; 69:1815–1821.
 47. Raz E. Guidance of primordial germ cell migration. *Curr Opin Cell Biol* 2004; 16:169–173.
 48. Runyan C, Schaible K, Molyneux K, Wang Z, Levin L, Wylie C. Steel factor controls midline cell death of primordial germ cells and is essential for their normal proliferation and migration. *Development* 2006; 133:4861–4869.
 49. Blume-Jensen P, Jiang G, Hyman R, Lee KF, O’Gorman S, Hunter T. Kit/stem cell factor receptor-induced activation of phosphatidylinositol 3’-kinase is essential for male fertility. *Nat Genet* 2000; 24:157–162.
 50. Chabot B, Stephenson DA, Chapman VM, Besmer P, Bernstein A. The proto-oncogene c-kit encoding a transmembrane tyrosine kinase receptor maps to the mouse W locus. *Nature* 1988; 335:88–89.
 51. Reith AD, Rottapel R, Giddens E, Brady C, Forrester L, Bernstein A. W mutant mice with mild or severe developmental defects contain distinct point mutations in the kinase domain of the c-kit receptor. *Genes Dev* 1990; 4:390–400.
 52. Moitsidou M, Reichman-Fried M, Stebler J, Kopranner M, Dorries J, Meyer D, Esguerra CV, Leung T, Raz E. Guidance of primordial germ cell migration by the chemokine SDF-1. *Cell* 2002; 111:647–659.
 53. Ara T, Nakamura Y, Egawa T, Sugiyama T, Abe K, Kishimoto T, Matsui Y, Nagasawa T. Impaired colonization of the gonads by primordial germ cells in mice lacking a chemokine, stromal cell-derived factor-1 (SDF-1). *Proc Natl Acad Sci U S A* 2003; 100:5319–5323.
 54. Molyneux KA, Zinszner H, Kunwar PS, Schaible K, Stebler J, Sunshine MJ, O’Brien W, Raz E, Littman D, Wylie C, Lehmann R. The chemokine SDF1/CXCL12 and its receptor CXCR4 regulate mouse germ cell migration and survival. *Development* 2003; 130:4279–4286.
 55. Pepling ME, Spradling AC. Female mouse germ cells form synchronously dividing cysts. *Development* 1998; 125:3323–3328.
 56. Lei L, Spradling AC. Mouse primordial germ cells produce cysts that partially fragment prior to meiosis. *Development* 2013; 140:2075–2081.
 57. Mork L, Maatouk DM, McMahon JA, Guo JJ, Zhang P, McMahon AP, Capel B. Temporal differences in granulosa cell specification in the ovary reflect distinct follicle fates in mice. *Biol Reprod* 2012; 86:37.
 58. Seisenberger S, Andrews S, Krueger F, Arand J, Walter J, Santos F, Popp C, Thienpont B, Dean W, Reik W. The dynamics of genome-wide DNA methylation reprogramming in mouse primordial germ cells. *Mol Cell* 2012; 48:849–862.
 59. Kagiwada S, Kurimoto K, Hirota T, Yamaji M, Saitou M. Replication-coupled passive DNA demethylation for the erasure of genome imprints in mice. *EMBO J* 2013; 32:340–353.
 60. Kurimoto K, Yamaji M, Seki Y, Saitou M. Specification of the germ cell lineage in mice: A process orchestrated by the PR-domain proteins, Blimp1 and Prdm14. *Cell Cycle* 2008; 7:3514–3518.
 61. Seki Y, Yamaji M, Yabuta Y, Sano M, Shigetani M, Matsui Y, Saga Y, Tachibana M, Shinkai Y, Saitou M. Cellular dynamics associated with the genome-wide epigenetic reprogramming in migrating primordial germ cells in mice. *Development* 2007; 134:2627–2638.
 62. Shirane K, Kurimoto K, Yabuta Y, Yamaji M, Satoh J, Ito S, Watanabe A, Hayashi K, Saitou M, Sasaki H. Global landscape and regulatory principles of DNA methylation reprogramming for germ cell specification by mouse pluripotent stem cells. *Dev Cell* 2016; 39:87–103.

63. Hargan-Calvopina J, Taylor S, Cook H, Hu Z, Lee SA, Yen MR, Chiang YS, Chen PY, Clark AT. Stage-specific demethylation in primordial germ cells safeguards against precocious differentiation. *Dev Cell* 2016; 39:75–86.
64. Hackett JA, Sengupta R, Zyliz JJ, Murakami K, Lee C, Down TA, Surani MA. Germline DNA demethylation dynamics and imprint erasure through 5-hydroxymethylcytosine. *Science* 2013; 339:448–452.
65. Yamaguchi S, Shen L, Liu Y, Sandler D, Zhang Y. Role of Tet1 in erasure of genomic imprinting. *Nature* 2013; 504:460–464.
66. Vincent JJ, Huang Y, Chen PY, Feng S, Calvopina JH, Nee K, Lee SA, Le T, Yoon AJ, Faull K, Fan G, Rao A et al. Stage-specific roles for tet1 and tet2 in DNA demethylation in primordial germ cells. *Cell Stem Cell* 2013; 12:470–478.
67. Kobayashi H, Sakurai T, Miura F, Imai M, Mochiduki K, Yanagisawa E, Sakashita A, Wakai T, Suzuki Y, Ito T, Matsui Y, Kono T. High-resolution DNA methylome analysis of primordial germ cells identifies gender-specific reprogramming in mice. *Genome Res* 2013; 23:616–627.
68. Yamaguchi S, Hong K, Liu R, Shen L, Inoue A, Diep D, Zhang K, Zhang Y. Tet1 controls meiosis by regulating meiotic gene expression. *Nature* 2012; 492:443–447.
69. Dawlaty MM, Breiling A, Le T, Raddatz G, Barrasa MI, Cheng AW, Gao Q, Powell BE, Li Z, Xu M, Faull KF, Lyko F et al. Combined deficiency of Tet1 and Tet2 causes epigenetic abnormalities but is compatible with postnatal development. *Dev Cell* 2013; 24:310–323.
70. Popp C, Dean W, Feng S, Cokus SJ, Andrews S, Pellegrini M, Jacobsen SE, Reik W. Genome-wide erasure of DNA methylation in mouse primordial germ cells is affected by AID deficiency. *Nature* 2010; 463:1101–1105.
71. Guibert S, Forne T, Weber M. Global profiling of DNA methylation erasure in mouse primordial germ cells. *Genome Res* 2012; 22:633–641.
72. Tang WW, Dietmann S, Irie N, Leitch HG, Floros VI, Bradshaw CR, Hackett JA, Chinnery PF, Surani MA. A unique gene regulatory network resets the human germline epigenome for development. *Cell* 2015; 161:1453–1467.
73. Guo F, Yan L, Guo H, Li L, Hu B, Zhao Y, Yong J, Hu Y, Wang X, Wei Y, Wang W, Li R et al. The transcriptome and DNA methylome landscapes of human primordial germ cells. *Cell* 2015; 161:1437–1452.
74. Hajkova P, Ancelin K, Waldmann T, Lacoste N, Lange UC, Cesari F, Lee C, Almouzni G, Schneider R, Surani MA. Chromatin dynamics during epigenetic reprogramming in the mouse germ line. *Nature* 2008; 452:877–881.
75. Hajkova P, Erhardt S, Lane N, Haaf T, El-Maarri O, Reik W, Walter J, Surani MA. Epigenetic reprogramming in mouse primordial germ cells. *Mech Dev* 2002; 117:15–23.
76. Hill PW, Amouroux R, Hajkova P. DNA demethylation, Tet proteins and 5-hydroxymethylcytosine in epigenetic reprogramming: An emerging complex story. *Genomics* 2014; 104:324–333.
77. Lee J, Inoue K, Ono R, Ogonuki N, Kohda T, Kaneko-Ishino T, Ogura A, Ishino F. Erasing genomic imprinting memory in mouse clone embryos produced from day 11.5 primordial germ cells. *Development* 2002; 129:1807–1817.
78. Surani MA, Hajkova P. Epigenetic reprogramming of mouse germ cells toward totipotency. *Cold Spring Harb Symp Quant Biol* 2010; 75:211–218.
79. Hill PWS, Leitch HG, Requena CE, Sun Z, Amouroux R, Roman-Trufero M, Borkowska M, Terragni J, Vaisvila R, Linnett S, Bagci H, Dharmalingham G et al. Epigenetic reprogramming enables the transition from primordial germ cell to gonocyte. *Nature* 2018; 555:392–396.
80. Williams K, Christensen J, Pedersen MT, Johansen JV, Cloos PA, Rapp-silber J, Helin K. TET1 and hydroxymethylcytosine in transcription and DNA methylation fidelity. *Nature* 2011; 473:343–348.
81. Vella P, Scelfo A, Jammula S, Chiacchiera F, Williams K, Cuomo A, Roberto A, Christensen J, Bonaldi T, Helin K, Pasini D. Tet proteins connect the O-linked N-acetylglucosamine transferase Ogt to chromatin in embryonic stem cells. *Mol Cell* 2013; 49:645–656.
82. Smith ZD, Meissner A. DNA methylation: Roles in mammalian development. *Nat Rev Genet* 2013; 14:204–220.
83. Seki Y, Hayashi K, Itoh K, Mizugaki M, Saitou M, Matsui Y. Extensive and orderly reprogramming of genome-wide chromatin modifications associated with specification and early development of germ cells in mice. *Dev Biol* 2005; 278:440–458.
84. Gkountela S, Zhang KX, Shafiq TA, Liao WW, Hargan-Calvopina J, Chen PY, Clark AT. DNA demethylation dynamics in the human prenatal germline. *Cell* 2015; 161:1425–1436.
85. Kurimoto K, Yabuta Y, Hayashi K, Ohta H, Kiyonari H, Mitani T, Moritoki Y, Kohri K, Kimura H, Yamamoto T, Katou Y, Shirahige K et al. Quantitative dynamics of chromatin Remodeling during germ cell specification from mouse embryonic stem cells. *Cell Stem Cell* 2015; 16:517–532.
86. Eguizabal C, Herrera L, De Onate L, Montserrat N, Hajkova P, Izpisua Belmonte JC. Characterization of the epigenetic changes during human gonadal primordial germ cells reprogramming. *Stem Cells* 2016; 34:2418–2428.
87. de Kretser DM, Loveland KL, Meinhardt A, Simorangkir D, Wreford N. Spermatogenesis. *Hum Reprod* 1998; 13:1–8.
88. Griswold MD. Action of FSH on mammalian Sertoli cells. In: Russell LD. GMD (ed.) *The Sertoli Cell*. Clearwater, FL: Cache River Press; 1993: 493–508.
89. Zirkin BR, Awoniyi C, Griswold MD, Russell LD, Sharpe R. Is FSH required for adult spermatogenesis? *J Androl* 1994; 15:273–276.
90. Abel MH, Baker PJ, Charlton HM, Monteiro A, Verhoeven G, De Gendt K, Guillouf F, O'Shaughnessy PJ. Spermatogenesis and sertoli cell activity in mice lacking sertoli cell receptors for follicle-stimulating hormone and androgen. *Endocrinology* 2008; 149:3279–3285.
91. Kumar TR, Wang Y, Lu N, Matzuk MM. Follicle stimulating hormone is required for ovarian follicle maturation but not male fertility. *Nat Genet* 1997; 15:201–204.
92. Abel MH, Wootton AN, Wilkins V, Huhtaniemi I, Knight PG, Charlton HM. The effect of a null mutation in the follicle-stimulating hormone receptor gene on mouse reproduction. *Endocrinology* 2000; 141:1795–1803.
93. Dierich A, Sairam MR, Monaco L, Fimia GM, Gansmuller A, LeMeur M, Sassone-Corsi P. Impairing follicle-stimulating hormone (FSH) signaling in vivo: Targeted disruption of the FSH receptor leads to aberrant gametogenesis and hormonal imbalance. *Proc Natl Acad Sci U S A* 1998; 95:13612–13617.
94. Wreford NG, Rajendra Kumar T, Matzuk MM, de Kretser DM. Analysis of the testicular phenotype of the follicle-stimulating hormone beta-subunit knockout and the activin type II receptor knockout mice by stereological analysis. *Endocrinology* 2001; 142:2916–2920.
95. O'Shaughnessy PJ, Monteiro A, Verhoeven G, De Gendt K, Abel MH. Effect of FSH on testicular morphology and spermatogenesis in gonadotrophin-deficient hypogonadal mice lacking androgen receptors. *Reproduction* 2010; 139:177–184.
96. Singh J, Handelsman DJ. The effects of recombinant FSH on testosterone-induced spermatogenesis in gonadotrophin-deficient (hpg) mice. *J Androl* 1996; 17:382–393.
97. Baines H, Nwagwu MO, Hastie GR, Wiles RA, Mayhew TM, Ebling FJ. Effects of estradiol and FSH on maturation of the testis in the hypogonadal (hpg) mouse. *Reprod Biol Endocrinol* 2008; 6:4.
98. Haywood M, Spaliviero J, Jimenez M, King NJ, Handelsman DJ, Allan CM. Sertoli and germ cell development in hypogonadal (hpg) mice expressing transgenic follicle-stimulating hormone alone or in combination with testosterone. *Endocrinology* 2003; 144:509–517.
99. Rodriguez I, Ody C, Araki K, Garcia I, Vassalli P. An early and massive wave of germinal cell apoptosis is required for the development of functional spermatogenesis. *EMBO J* 1997; 16:2262–2270.
100. McLachlan RI, Wreford NG, de Kretser DM, Robertson DM. The effects of recombinant follicle-stimulating hormone on the restoration of spermatogenesis in the gonadotropin-releasing hormone-immunized adult rat. *Endocrinology* 1995; 136:4035–4043.
101. Russell LD, Corbin TJ, Borg KE, De Franca LR, Grasso P, Bartke A. Recombinant human follicle-stimulating hormone is capable of exerting a biological effect in the adult hypophysectomized rat by

- reducing the numbers of degenerating germ cells. *Endocrinology* 1993; 133:2062–2070.
102. Sun YT, Wreford NG, Robertson DM, de Kretser DM. Quantitative cytological studies of spermatogenesis in intact and hypophysectomized rats: Identification of androgen-dependent stages. *Endocrinology* 1990; 127:1215–1223.
 103. Ruwanpura SM, McLachlan RI, Stanton PG, Loveland KL, Meachem SJ. Pathways involved in testicular germ cell apoptosis in immature rats after FSH suppression. *J Endocrinol* 2008; 197:35–43.
 104. Ruwanpura SM, McLachlan RI, Stanton PG, Meachem SJ. Follicle-stimulating hormone affects spermatogonial survival by regulating the intrinsic apoptotic pathway in adult rats. *Biol Reprod* 2008; 78:705–713.
 105. Simorangkir DR, Ramaswamy S, Marshall GR, Pohl CR, Plant TM. A selective monotropic elevation of FSH, but not that of LH, amplifies the proliferation and differentiation of spermatogonia in the adult rhesus monkey (*Macaca mulatta*). *Hum Reprod* 2009; 24:1584–1595.
 106. Henriksen K, Kangasniemi M, Parvinen M, Kaipia A, Hakovirta H. In vitro, follicle-stimulating hormone prevents apoptosis and stimulates deoxyribonucleic acid synthesis in the rat seminiferous epithelium in a stage-specific fashion. *Endocrinology* 1996; 137:2141–2149.
 107. Skinner MK. Cell-cell interactions in the testis. *Endocr Rev* 1991; 12:45–77.
 108. Grover A, Sairam MR, Smith CE, Hermo L. Structural and functional modifications of sertoli cells in the testis of adult follicle-stimulating hormone receptor knockout mice. *Biol Reprod* 2004; 71:117–129.
 109. Hazra R, Corcoran L, Robson M, McTavish KJ, Upton D, Handelsman DJ, Allan CM. Temporal role of Sertoli cell androgen receptor expression in spermatogenic development. *Mol Endocrinol* 2013; 27:12–24.
 110. Pakarainen T, Zhang FP, Makela S, Poutanen M, Huhtaniemi I. Testosterone replacement therapy induces spermatogenesis and partially restores fertility in luteinizing hormone receptor knockout mice. *Endocrinology* 2005; 146:596–606.
 111. Cameron DF, Muffly KE, Nazian SJ. Reduced testosterone during puberty results in a midspemiogenic lesion. *Proc Soc Exp Biol Med* 1993; 202:457–464.
 112. Huang HF, Marshall GR, Rosenberg R, Nieschlag E. Restoration of spermatogenesis by high levels of testosterone in hypophysectomized rats after long-term regression. *Acta Endocrinol* 1987; 116:433–444.
 113. Sun YT, Irby DC, Robertson DM, de Kretser DM. The effects of exogenously administered testosterone on spermatogenesis in intact and hypophysectomized rats. *Endocrinology* 1989; 125:1000–1010.
 114. Awoniyi CA, Sprando RL, Santulli R, Chandrashekar V, Ewing LL, Zirkin BR. Restoration of spermatogenesis by exogenously administered testosterone in rats made azoospermic by hypophysectomy or withdrawal of luteinizing hormone alone. *Endocrinology* 1990; 127:177–184.
 115. Sharpe RM, Fraser HM, Ratnasooriya WD. Assessment of the role of Leydig cell products other than testosterone in spermatogenesis and fertility in adult rats. *Int J Androl* 1988; 11:507–523.
 116. O'Donnell L, Meachem SJ, Stanton PG, RI ML. CHAPTER 21-Endocrine Regulation of Spermatogenesis. In: *Knobil and Neill's Physiology of Reproduction (Third Edition)*, Academic Press, Cambridge, MA, vol. 1. 2006: 1017–1069.
 117. Russell LD, Clermont Y. Degeneration of germ cells in normal, hypophysectomized and hormone treated hypophysectomized rats. *Anat Rec* 1977; 187:347–366.
 118. O'Shaughnessy PJ, Monteiro A, Abel M. Testicular development in mice lacking receptors for follicle stimulating hormone and androgen. *PLoS One* 2012; 7:e35136.
 119. Lyon MF, Hawkes SG. X-linked gene for testicular feminization in the mouse. *Nature* 1970; 227:1217–1219.
 120. Johnston H, Baker PJ, Abel M, Charlton HM, Jackson G, Fleming L, Kumar TR, O'Shaughnessy PJ. Regulation of Sertoli cell number and activity by follicle-stimulating hormone and androgen during postnatal development in the mouse. *Endocrinology* 2004; 145:318–329.
 121. De Gendt K, Swinnen JV, Saunders PT, Schoonjans L, Dewerchin M, Devos A, Tan K, Atanassova N, Claessens F, Lecreuil C, Heyns W, Carmeliet P et al. A Sertoli cell-selective knockout of the androgen receptor causes spermatogenic arrest in meiosis. *Proc Natl Acad Sci U S A* 2004; 101:1327–1332.
 122. Lyon MF, Glenister PH, Lamoreux ML. Normal spermatozoa from androgen-resistant germ cells of chimaeric mice and the role of androgen in spermatogenesis. *Nature* 1975; 258:620–622.
 123. Johnston DS, Russell LD, Friel PJ, Griswold MD. Murine germ cells do not require functional androgen receptors to complete spermatogenesis following spermatogonial stem cell transplantation. *Endocrinology* 2001; 142:2405–2408.
 124. Allan CM, Couse JF, Simanainen U, Spaliviero J, Jimenez M, Rodriguez K, Korach KS, Handelsman DJ. Estradiol induction of spermatogenesis is mediated via an estrogen receptor- α mechanism involving neuroendocrine activation of follicle-stimulating hormone secretion. *Endocrinology* 2010; 151:2800–2810.
 125. Ebling FJ, Brooks AN, Cronin AS, Ford H, Kerr JB. Estrogenic induction of spermatogenesis in the hypogonadal mouse. *Endocrinology* 2000; 141:2861–2869.
 126. Hunt DM, Saksena SK, Chang MC. Effects of estradiol-17 beta on reproduction in adult male rats. *Arch Androl* 1979; 2:129–133.
 127. Zondek B. Mass excretion of (Estrogenic hormone in the urine of the stallion. *Nature* 1934; 133:209–210.
 128. Dorrington JH, Fritz IB, Armstrong DT. Control of testicular estrogen synthesis. *Biol Reprod* 1978; 18:55–64.
 129. Carreau S, Bouraima-Lelong H, Delalande C. Estrogen, a female hormone involved in spermatogenesis. *Adv Med Sci* 2012; 57:31–36.
 130. Chimento A, Sirianni R, Delalande C, Silandre D, Bois C, Ando S, Maggiolini M, Carreau S, Pezzi V. 17 beta-estradiol activates rapid signaling pathways involved in rat pachytene spermatocytes apoptosis through GPR30 and ER α . *Mol Cell Endocrinol* 2010; 320:136–144.
 131. Robertson KM, O'Donnell L, Jones ME, Meachem SJ, Boon WC, Fisher CR, Graves KH, McLachlan RI, Simpson ER. Impairment of spermatogenesis in mice lacking a functional aromatase (*cyp 19*) gene. *Proc Natl Acad Sci U S A* 1999; 96:7986–7991.
 132. Sinkevicius KW, Laine M, Lotan TL, Woloszyn K, Richburg JH, Greene GL. Estrogen-dependent and -independent estrogen receptor- α signaling separately regulate male fertility. *Endocrinology* 2009; 150:2898–2905.
 133. Cacciola G, Chioccarelli T, Fasano S, Pierantoni R, Cobellis G. Estrogens and spermiogenesis: New insights from type 1 cannabinoid receptor knockout mice. *Int J Endocrinol* 2013; 2013:501350.
 134. Shetty G, Weng CC, Bolden-Tiller OU, Huhtaniemi I, Handelsman DJ, Meistrich ML. Effects of medroxyprogesterone and estradiol on the recovery of spermatogenesis in irradiated rats. *Endocrinology* 2004; 145:4461–4469.
 135. Porter KL, Shetty G, Shuttlesworth GA, Weng CC, Huhtaniemi I, Pakarainen P, Meistrich ML. Estrogen enhances recovery from radiation-induced spermatogonial arrest in rat testes. *J Androl* 2009; 30:440–451.
 136. Hedger MP, Winnall WR. Regulation of activin and inhibin in the adult testis and the evidence for functional roles in spermatogenesis and immunoregulation. *Mol Cell Endocrinol* 2012; 359:30–42.
 137. Hakovirta H, Kaipia A, Soder O, Parvinen M. Effects of activin-a, inhibin-a, and transforming growth factor-beta 1 on stage-specific deoxyribonucleic acid synthesis during rat seminiferous epithelial cycle. *Endocrinology* 1993; 133:1664–1668.
 138. Mather JP, Attie KM, Woodruff TK, Rice GC, Phillips DM. Activin stimulates spermatogonial proliferation in germ-Sertoli cell cocultures from immature rat testis. *Endocrinology* 1990; 127:3206–3214.
 139. Nagano M, Ryu BY, Brinster CJ, Avarbock MR, Brinster RL. Maintenance of mouse male germ line stem cells in vitro. *Biol Reprod* 2003; 68:2207–2214.
 140. Guo Q, Kumar TR, Woodruff T, Hadsell LA, DeMayo FJ, Matzuk MM. Overexpression of mouse follistatin causes reproductive defects in transgenic mice. *Mol Endocrinol* 1998; 12:96–106.

141. Oldknow KJ, Seebacher J, Goswami T, Villen J, Pitsillides AA, O'Shaughnessy PJ, Gygi SP, Schneyer AL, Mukherjee A. Follistatin-like 3 (FSTL3) mediated silencing of transforming growth factor beta (TGFbeta) signaling is essential for testicular aging and regulating testis size. *Endocrinology* 2013; **154**:1310–1320.
142. Heller CH, Clermont Y. Kinetics of the germinal epithelium in man. *Recent Prog Horm Res* 1964; **20**:545–575.
143. Reith DA, Bernstein A. Molecular biology of the W and Steel loci. *Genome Anal* 1991; **3**:105–133.
144. Munsie M, Schlatt S, deKretser DM, Loveland KL. Expression of stem cell factor in the postnatal rat testis. *Mol Reprod Dev* 1997; **47**: 19–25.
145. Cupp AS, Skinner MK. Expression, action, and regulation of transforming growth factor alpha and epidermal growth factor receptor during embryonic and perinatal rat testis development. *J Androl* 2001; **22**:1019–1029.
146. Cupp AS, Tessarollo L, Skinner MK. Testis developmental phenotypes in neurotrophin receptor trkA and trkC null mutations: Role in formation of seminiferous cords and germ cell survival. *Biol Reprod* 2002; **66**:1838–1845.
147. Packer AI, Besmer P, Bachvarova RF. Kit ligand mediates survival of type a spermatogonia and dividing spermatocytes in postnatal mouse testes. *Mol Reprod Dev* 1995; **42**:303–310.
148. Shetty J, Marathe GK, Dighe RR. Specific immunoneutralization of FSH leads to apoptotic cell death of the pachytene spermatocytes and spermatogonial cells in the rat. *Endocrinology* 1996; **137**: 2179–2182.
149. Oatley JM, Avarbock MR, Brinster RL. Glial cell line-derived neurotrophic factor regulation of genes essential for self-renewal of mouse spermatogonial stem cells is dependent on Src family kinase signaling. *J Biol Chem* 2007; **282**:25842–25851.
150. Oatley JM, Oatley MJ, Avarbock MR, Tobias JW, Brinster RL. Colony stimulating factor 1 is an extrinsic stimulator of mouse spermatogonial stem cell self-renewal. *Development* 2009; **136**:1191–1199.
151. Oatley MJ, Kaucher AV, Racicot KE, Oatley JM. Inhibitor of DNA binding 4 is expressed selectively by single spermatogonia in the male germline and regulates the self-renewal of spermatogonial stem cells in mice. *Biol Reprod* 2011; **85**:347–356.
152. Ciccarelli M, Giassetti MI, Miao D, Oatley MJ, Robbins C, Lopez-Biladeau B, Waqas MS, Tibary A, Whitelaw B, Lillico S, Park CH, Park KE et al. Donor-derived spermatogenesis following stem cell transplantation in sterile NANOS2 knockout males. *Proc Natl Acad Sci U S A* 2020; **117**:24195–24204.
153. Hermann BP, Cheng K, Singh A, Roa-De La Cruz L, Mutoji KN, Chen IC, Gildersleeve H, Lehle JD, Mayo M, Westernstroer B, Law NC, Oatley MJ et al. The mammalian spermatogenesis single-cell transcriptome, from Spermatogonial stem cells to spermatids. *Cell Rep* 2018; **25**:1650–1667 e1658.
154. Griswold MD, Oatley JM. Concise review: Defining characteristics of mammalian spermatogenic stem cells. *Stem Cells* 2013; **31**:8–11.
155. Leslie M. EPIGENETICS. Sperm RNA fragments modify offspring metabolism. *Science* 2016; **351**:13.
156. Nilsson E, Sadler-Riggleman I, Skinner MK. Environmentally induced epigenetic transgenerational inheritance of disease. *Environmental Epigenetics* 2018; **4**:1, dvy016–13.
157. Anway MD, Cupp AS, Uzumcu M, Skinner MK. Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science* 2005; **308**:1466–1469.
158. Sharma U, Conine CC, Shea JM, Boskovic A, Derr AG, Bing XY, Belleannee C, Kucukural A, Serra RW, Sun F, Song L, Carone BR et al. Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science* 2016; **351**: 391–396.
159. Chen Q, Yan M, Cao Z, Li X, Zhang Y, Shi J, Feng GH, Peng H, Zhang X, Zhang Y, Qian J, Duan E et al. Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* 2016; **351**:397–400.
160. Aravin AA, Sachidanandam R, Bourc'his D, Schaefer C, Pezic D, Toth KF, Bestor T, Hannon GJ. A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol Cell* 2008; **31**:785–799.
161. Stalker L, Russell SJ, Co C, Foster RA, LaMarre J. PIWIL1 is expressed in the canine testis, increases with sexual maturity, and binds small RNAs. *Biol Reprod* 2016; **94**:17.
162. Ben Maamar M, Beck D, Nilsson E, McCarrey JR, Skinner MK. Developmental alterations in DNA methylation during gametogenesis from primordial germ cells to sperm. *Dev Biol* 2021; Second revision submitted and publication pending.
163. Vreeburg JT, Holland MK, Cornwall GA, Rankin TL, Orgebin-Crist MC. Secretion of epididymal proteins and their interactions with spermatozoa. *Bull Assoc Anat (Nancy)* 1991; **75**:171–173.
164. Cornwall GA. Role of posttranslational protein modifications in epididymal sperm maturation and extracellular quality control. *Adv Exp Med Biol* 2014; **759**:159–180.
165. Austin CR. Observations on the penetration of the sperm in the mammalian egg. *Aust J Sci Res B* 1951; **4**:581–596.
166. Chang MC. Fertilizing capacity of spermatozoa deposited into the fallopian tubes. *Nature* 1951; **168**:697–698.
167. Young LG, Hinton BT, Gould KG. Surface changes in chimpanzee sperm during epididymal transit. *Biol Reprod* 1985; **32**:399–412.
168. Han Y, Haines CJ, Feng HL. Role(s) of the serine/threonine protein phosphatase 1 on mammalian sperm motility. *Arch Androl* 2007; **53**:169–177.
169. James ER, Carrell DT, Aston KI, Jenkins TG, Yeste M, Salas-Huetos A. The role of the epididymis and the contribution of Epididymosomes to mammalian reproduction. *Int J Mol Sci* 2020; **21**:5377.
170. Sullivan R, Legare C, Lamontagne-Proulx J, Breton S, Soulet D. Revisiting structure/functions of the human epididymis. *Andrology* 2019; **7**:748–757.
171. Sharma A. Transgenerational epigenetics: Integrating soma to germline communication with gametic inheritance. *Mech Ageing Dev* 2017; **163**:15–22.
172. Conine CC, Sun F, Song L, Rivera-Perez JA, Rando OJ. MicroRNAs absent in caput sperm are required for normal embryonic development. *Dev Cell* 2019; **50**:7–8.
173. Chu C, Zhang YL, Yu L, Sharma S, Fei ZL, Drevet JR. Epididymal small non-coding RNA studies: Progress over the past decade. *Andrology* 2019; **7**:681–689.
174. Sharma U. Paternal contributions to offspring health: Role of sperm small RNAs in intergenerational transmission of epigenetic information. *Front Cell Dev Biol* 2019; **7**:215.
175. Scott IM, Rubinstein GM, Poole FL 2nd, Lipscomb GL, Schut GJ, Williams-Rhaesa AM, Stevenson DM, Amador-Noguez D, Kelly RM, Adams MWW. The thermophilic biomass-degrading bacterium *Caldicellulosiruptor bescii* utilizes two enzymes to oxidize glyceraldehyde 3-phosphate during glycolysis. *J Biol Chem* 2019; **294**:9995–10005.
176. Ben Maamar M, Nilsson E, Sadler-Riggleman I, Beck D, McCarrey JR, Skinner MK. Developmental origins of transgenerational sperm DNA methylation epimutations following ancestral DDT exposure. *Dev Biol* 2019; **445**:280–293.
177. Ben Maamar M, Beck D, Nilsson E, McCarrey JR, Skinner MK. Developmental origins of transgenerational sperm histone retention following ancestral exposures. *Dev Biol* 2020; **465**:31–45.
178. Sanchez F, Smitz J. Molecular control of oogenesis. *Biochim Biophys Acta* 1822; **2012**:1896–1912.
179. Mehlmann LM. Stops and starts in mammalian oocytes: Recent advances in understanding the regulation of meiotic arrest and oocyte maturation. *Reproduction* 2005; **130**:791–799.
180. Rajah R, Glaser EM, Hirshfield AN. The changing architecture of the neonatal rat ovary during histogenesis. *Dev Dyn* 1992; **194**:177–192.
181. El-Hayek S, Clarke HJ. Control of oocyte growth and development by intercellular communication within the follicular niche. *Results Probl Cell Differ* 2016; **58**:191–224.

182. Buccione R, Schroeder AC, Eppig JJ. Interactions between somatic cells and germ cells throughout mammalian oogenesis. *Biol Reprod* 1990; 43:543–547.
183. Kezele P, Nilsson E, Skinner MK. Cell-cell interactions in primordial follicle assembly and development. *Front Biosci* 2002; 7:d1990–d1996.
184. Grive KJ. Pathways coordinating oocyte attrition and abundance during mammalian ovarian reserve establishment. *Mol Reprod Dev* 2020; 87:843–856.
185. Chen Y, Yang W, Shi X, Zhang C, Song G, Huang D. The factors and pathways regulating the activation of mammalian primordial follicles in vivo. *Front Cell Dev Biol* 2020; 8:575706.
186. Skinner MK. Regulation of primordial follicle assembly and development. *Hum Reprod Update* 2005; 11:461–471.
187. Gougeon A. Human ovarian follicular development: From activation of resting follicles to preovulatory maturation. *Ann Endocrinol (Paris)* 2010; 71:132–143.
188. Rodgers RJ, Irving-Rodgers HF. Formation of the ovarian follicular antrum and follicular fluid. *Biol Reprod* 2010; 82:1021–1029.
189. Son WY, Das M, Shalom-Paz E, Holzer H. Mechanisms of follicle selection and development. *Minerva Ginecol* 2011; 63:89–102.
190. Craig J, Orisaka M, Wang H, Orisaka S, Thompson W, Zhu C, Kotsuji F, Tsang BK. Gonadotropin and intra-ovarian signals regulating follicle development and atresia: The delicate balance between life and death. *Front Biosci* 2007; 12:3628–3639.
191. Marcozzi S, Rossi V, Salustri A, De Felici M, Klinger FG. Programmed cell death in the human ovary. *Minerva Ginecol* 2018; 70:549–560.
192. Sen A, Caiazza F. Oocyte maturation: A story of arrest and release. *Front Biosci (Schol Ed)* 2013; 5:451–477.
193. Georgadaki K, Khoury N, Spandidos DA, Zoumpourlis V. The molecular basis of fertilization (review). *Int J Mol Med* 2016; 38:979–986.
194. Bhakta HH, Refai FH, Avella MA. The molecular mechanisms mediating mammalian fertilization. *Development* 2019; 146:dev176966.
195. Svoboda P. Mammalian zygotic genome activation. *Semin Cell Dev Biol* 2018; 84:118–126.
196. Schulz KN, Harrison MM. Mechanisms regulating zygotic genome activation. *Nat Rev Genet* 2019; 20:221–234.
197. Grossman H, Shalgi R. A role of MicroRNAs in cell differentiation during gonad development. *Results Probl Cell Differ* 2016; 58:309–336.
198. Liang Y, Ridzon D, Wong L, Chen C. Characterization of microRNA expression profiles in normal human tissues. *BMC Genomics* 2007; 8:166.
199. Ro S, Song R, Park C, Zheng H, Sanders KM, Yan W. Cloning and expression profiling of small RNAs expressed in the mouse ovary. *RNA* 2007; 13:2366–2380.
200. Yao G, Liang M, Liang N, Yin M, Lu M, Lian J, Wang Y, Sun F. MicroRNA-224 is involved in the regulation of mouse cumulus expansion by targeting Ptx3. *Mol Cell Endocrinol* 2014; 382:244–253.
201. Tomizawa S, Nowacka-Woszek J, Kelsey G. DNA methylation establishment during oocyte growth: Mechanisms and significance. *Int J Dev Biol* 2012; 56:867–875.
202. Sendzikaite G, Kelsey G. The role and mechanisms of DNA methylation in the oocyte. *Essays Biochem* 2019; 63:691–705.
203. Sha QQ, Zhang J, Fan HY. Function and regulation of histone H3 Lysine-4 methylation during oocyte meiosis and maternal-to-zygotic transition. *Front Cell Dev Biol* 2020; 8:597498.
204. Yu C, Fan X, Sha QQ, Wang HH, Li BT, Dai XX, Shen L, Liu J, Wang L, Liu K, Tang F, Fan HY. CFP1 regulates histone H3K4 Trimethylation and developmental potential in mouse oocytes. *Cell Rep* 2017; 20:1161–1172.
205. Sha QQ, Dai XX, Jiang JC, Yu C, Jiang Y, Liu J, Ou XH, Zhang SY, Fan HY. CFP1 coordinates histone H3 lysine-4 trimethylation and meiotic cell cycle progression in mouse oocytes. *Nat Commun* 2018; 9:3477.
206. Andreu-Vieyra CV, Chen R, Agno JE, Glaser S, Anastasiadis K, Stewart AF, Matzuk MM. MLL2 is required in oocytes for bulk histone 3 lysine 4 trimethylation and transcriptional silencing. *PLoS Biol* 2010; 8:e1000453.
207. Yamashiro C, Sasaki K, Yabuta Y, Kojima Y, Nakamura T, Okamoto I, Yokobayashi S, Murase Y, Ishikura Y, Shirane K, Sasaki H, Yamamoto T et al. Generation of human oogonia from induced pluripotent stem cells in vitro. *Science* 2018; 362:356–360.
208. Okae H, Chiba H, Hiura H, Hamada H, Sato A, Utsunomiya T, Kikuchi H, Yoshida H, Tanaka A, Suyama M, Arima T. Genome-wide analysis of DNA methylation dynamics during early human development. *PLoS Genet* 2014; 10:e1004868.
209. von Meyenn F, Berrens RV, Andrews S, Santos F, Collier AJ, Krueger F, Osorno R, Dean W, Rugg-Gunn PJ, Reik W. Comparative principles of DNA methylation reprogramming during human and mouse in vitro primordial germ cell specification. *Dev Cell* 2016; 39:104–115.
210. Pokholok DK, Harbison CT, Levine S, Cole M, Hannett NM, Lee TI, Bell GW, Walker K, Rolfe PA, Herbolsheimer E, Zeitlinger J, Lewitter F et al. Genome-wide map of nucleosome acetylation and methylation in yeast. *Cell* 2005; 122:517–527.
211. Boland MJ, Nazor KL, Loring JF. Epigenetic regulation of pluripotency and differentiation. *Circ Res* 2014; 115:311–324.
212. Avgustinova A, Benitah SA. Epigenetic control of adult stem cell function. *Nat Rev Mol Cell Biol* 2016; 17:643–658.
213. Kato Y, Kaneda M, Hata K, Kumaki K, Hisano M, Kohara Y, Okano M, Li E, Nozaki M, Sasaki H. Role of the Dnmt3 family in de novo methylation of imprinted and repetitive sequences during male germ cell development in the mouse. *Hum Mol Genet* 2007; 16:2272–2280.
214. Li JY, Lees-Murdock DJ, Xu GL, Walsh CP. Timing of establishment of paternal methylation imprints in the mouse. *Genomics* 2004; 84:952–960.
215. Bestor TH, Bourc'his D. Transposon silencing and imprint establishment in mammalian germ cells. *Cold Spring Harb Symp Quant Biol* 2004; 69:381–387.
216. Bourc'his D, Xu GL, Lin CS, Bollman B, Bestor TH. Dnmt3L and the establishment of maternal genomic imprints. *Science* 2001; 294:2536–2539.
217. Kaneda M, Okano M, Hata K, Sado T, Tsujimoto N, Li E, Sasaki H. Essential role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting. *Nature* 2004; 429:900–903.
218. Smallwood SA, Tomizawa S, Krueger F, Ruf N, Carli N, Segonds-Pichon A, Sato S, Hata K, Andrews SR, Kelsey G. Dynamic CpG island methylation landscape in oocytes and preimplantation embryos. *Nat Genet* 2011; 43:811–814.
219. Marston AL, Amon A. Meiosis: Cell-cycle controls shuffle and deal. *Nat Rev Mol Cell Biol* 2004; 5:983–997.
220. Kota SK, Feil R. Epigenetic transitions in germ cell development and meiosis. *Dev Cell* 2010; 19:675–686.
221. Baudat F, Manova K, Yuen JP, Jasin M, Keeney S. Chromosome synapsis defects and sexually dimorphic meiotic progression in mice lacking Spo11. *Mol Cell* 2000; 6:989–998.
222. Zelazowski MJ, Sandoval M, Paniker L, Hamilton HM, Han J, Gribbell MA, Kang R, Cole F. Age-dependent alterations in meiotic recombination cause chromosome segregation errors in spermatocytes. *Cell* 2017; 171:601–614 e613.
223. Baudat F, Buard J, Grey C, Fledel-Alon A, Ober C, Przeworski M, Coop G, de Massy B. PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. *Science* 2010; 327:836–840.
224. Hayashi K, Yoshida K, Matsui Y. A histone H3 methyltransferase controls epigenetic events required for meiotic prophase. *Nature* 2005; 438:374–378.
225. Jeffreys AJ, Kauppi L, Neumann R. Intensely punctate meiotic recombination in the class II region of the major histocompatibility complex. *Nat Genet* 2001; 29:217–222.
226. McVean GA, Myers SR, Hunt S, Deloukas P, Bentley DR, Donnelly P. The fine-scale structure of recombination rate variation in the human genome. *Science* 2004; 304:581–584.
227. Myers S, Bottolo L, Freeman C, McVean G, Donnelly P. A fine-scale map of recombination rates and hotspots across the human genome. *Science* 2005; 310:321–324.

228. Parvanov ED, Petkov PM, Paigen K. Prdm9 controls activation of mammalian recombination hotspots. *Science* 2010; 327:835.
229. Gong F, Miller KM. Mammalian DNA repair: HATs and HDACs make their mark through histone acetylation. *Mutat Res* 2013; 750: 23–30.
230. Hirota K, Mizuno K, Shibata T, Ohta K. Distinct chromatin modulators regulate the formation of accessible and repressive chromatin at the fission yeast recombination hotspot ade6-M26. *Mol Biol Cell* 2008; 19:1162–1173.
231. Sollier J, Lin W, Soustelle C, Suhre K, Nicolas A, Geli V, de La Roche Saint-Andre C. Set1 is required for meiotic S-phase onset, double-strand break formation and middle gene expression. *EMBO J* 2004; 23:1957–1967.
232. Yamashita K, Shinohara M, Shinohara A. Rad6-Bre1-mediated histone H2B ubiquitylation modulates the formation of double-strand breaks during meiosis. *Proc Natl Acad Sci U S A* 2004; 101: 11380–11385.
233. Hanna CW, Taudt A, Huang J, Gahurova L, Kranz A, Andrews S, Dean W, Stewart AF, Colome-Tatche M, Kelsey G. MLL2 conveys transcription-independent H3K4 trimethylation in oocytes. *Nat Struct Mol Biol* 2018; 25:73–82.
234. Stewart KR, Veselovska L, Kim J, Huang J, Saadeh H, Tomizawa S, Smallwood SA, Chen T, Kelsey G. Dynamic changes in histone modifications precede de novo DNA methylation in oocytes. *Genes Dev* 2015; 29:2449–2462.
235. Rando OJ. Intergenerational transfer of epigenetic information in sperm. *Cold Spring Harb Perspect Med* 2016.
236. Bao J, Bedford MT. Epigenetic regulation of the histone-to-protamine transition during spermiogenesis. *Reproduction* 2016; 151: R55–R70.
237. Barral S, Morozumi Y, Tanaka H, Montellier E, Govin J, de Dieuleveult M, Charbonnier G, Coute Y, Puthier D, Buchou T, Boussouar F, Urahama T et al. Histone variant H2A.L.2 guides transition protein-dependent protamine assembly in male germ cells. *Mol Cell* 2017; 66:89–101 e108.
238. Houghoughi N, Barral S, Vargas A, Rousseaux S, Khochbin S. Histone variants: Essential actors in male genome programming. *J Biochem* 2018; 163:97–103.
239. Furrer R. Physical characteristics of mouse sperm nuclei. *Biophys J* 1976; 16:811–825.
240. Hud NV, Allen MJ, Downing KH, Lee J, Balhorn R. Identification of the elemental packing unit of DNA in mammalian sperm cells by atomic force microscopy. *Biochem Biophys Res Commun* 1993; 193:1347–1354.
241. Brykczynska U, Hisano M, Erkek S, Ramos L, Oakeley EJ, Roloff TC, Beisel C, Schubeler D, Stadler MB, Peters AH. Repressive and active histone methylation mark distinct promoters in human and mouse spermatozoa. *Nat Struct Mol Biol* 2010; 17:679–687.
242. Hammoud SS, Nix DA, Zhang H, Purwar J, Carrell DT, Cairns BR. Distinctive chromatin in human sperm packages genes for embryo development. *Nature* 2009; 460:473–478.
243. Ihara M, Meyer-Ficca ML, Leu NA, Rao S, Li F, Gregory BD, Zalenskaya IA, Schultz RM, Meyer RG. Paternal poly (ADP-ribose) metabolism modulates retention of inheritable sperm histones and early embryonic gene expression. *PLoS Genet* 2014; 10:e1004317.
244. Siklenka K, Erkek S, Godmann M, Lambrot R, McGraw S, Lafleur C, Cohen T, Xia J, Suderman M, Hallett M, Trasler J, Peters AH et al. Disruption of histone methylation in developing sperm impairs offspring health transgenerationally. *Science* 2015, 2001–2014; 350:aab2006.
245. Larriba E, del Mazo J. Role of non-coding RNAs in the transgenerational epigenetic transmission of the effects of Reprotoxicants. *Int J Mol Sci* 2016; 17:452.
246. Bhin J, Jeong HS, Kim JS, Shin JO, Hong KS, Jung HS, Kim C, Hwang D, Kim KS. PGC-enriched miRNAs control germ cell development. *Mol Cells* 2015; 38:895–903.
247. Medeiros LA, Dennis LM, Gill ME, Houbaviv H, Markoulaki S, Fu D, White AC, Kirak O, Sharp PA, Page DC, Jaenisch R. Mir-290-295 deficiency in mice results in partially penetrant embryonic lethality and germ cell defects. *Proc Natl Acad Sci U S A* 2011; 108:14163–14168.
248. Takada S, Berezikov E, Choi YL, Yamashita Y, Mano H. Potential role of miR-29b in modulation of Dnmt3a and Dnmt3b expression in primordial germ cells of female mouse embryos. *RNA* 2009; 15:1507–1514.
249. Zhang J, Liu W, Jin Y, Jia P, Jia K, Yi M. MiR-202-5p is a novel germ plasm-specific microRNA in zebrafish. *Sci Rep* 2017; 7:7055.
250. Staton AA, Knaut H, Giraldez AJ. miRNA regulation of Sdf1 chemokine signaling provides genetic robustness to germ cell migration. *Nat Genet* 2011; 43:204–211.
251. Yu M, Mu H, Niu Z, Chu Z, Zhu H, Hua J. miR-34c enhances mouse spermatogonial stem cells differentiation by targeting Nanos2. *J Cell Biochem* 2014; 115:232–242.
252. Liang X, Zhou D, Wei C, Luo H, Liu J, Fu R, Cui S. MicroRNA-34c enhances murine male germ cell apoptosis through targeting ATF1. *PLoS One* 2012; 7:e33861.
253. Bouhallier F, Allioli N, Lavial F, Chalmel F, Perrard MH, Durand P, Samarut J, Pain B, Rouault JP. Role of miR-34c microRNA in the late steps of spermatogenesis. *RNA* 2010; 16:720–731.
254. McIver SC, Stanger SJ, Santarelli DM, Roman SD, Nixon B, McLaughlin EA. A unique combination of male germ cell miRNAs coordinates gonocyte differentiation. *PLoS One* 2012; 7:e35553.
255. Tong MH, Mitchell D, Evanoff R, Griswold MD. Expression of Mirlet7 family microRNAs in response to retinoic acid-induced spermatogonial differentiation in mice. *Biol Reprod* 2011; 85:189–197.
256. Huszar JM, Payne CJ. MicroRNA 146 (Mir146) modulates spermatogonial differentiation by retinoic acid in mice. *Biol Reprod* 2013; 88:15.
257. Bao J, Li D, Wang L, Wu J, Hu Y, Wang Z, Chen Y, Cao X, Jiang C, Yan W, Xu C. MicroRNA-449 and microRNA-34b/c function redundantly in murine testes by targeting E2F transcription factor-retinoblastoma protein (E2F-pRb) pathway. *J Biol Chem* 2012; 287:21686–21698.
258. Chen X, Li X, Guo J, Zhang P, Zeng W. The roles of microRNAs in regulation of mammalian spermatogenesis. *J Anim Sci Biotechnol* 2017; 8:35.
259. Weick EM, Miska EA. piRNAs: From biogenesis to function. *Development* 2014; 141:3458–3471.
260. Ku HY, Lin H. PIWI proteins and their interactors in piRNA biogenesis, germline development and gene expression. *Natl Sci Rev* 2014; 1:205–218.
261. Houwing S, Kamminga LM, Berezikov E, Cronembold D, Girard A, van den Elst H, Filippov DV, Blaser H, Raz E, Moens CB, Plasterk RH, Hannon GJ et al. A role for Piwi and piRNAs in germ cell maintenance and transposon silencing in zebrafish. *Cell* 2007; 129:69–82.
262. Taylor DH, Chu ET, Spektor R, Soloway PD. Long non-coding RNA regulation of reproduction and development. *Mol Reprod Dev* 2015; 82:932–956.
263. Zhang C, Gao L, Xu EY. LncRNA, a new component of expanding RNA-protein regulatory network important for animal sperm development. *Semin Cell Dev Biol* 2016; 59:110–117.
264. Anguera MC, Ma W, Clift D, Namekawa S, Kelleher RJ 3rd, Lee JT. Tsx produces a long noncoding RNA and has general functions in the germline, stem cells, and brain. *PLoS Genet* 2011; 7:e1002248.
265. Nolasco S, Bellido J, Goncalves J, Tavares A, Zabala JC, Soares H. The expression of tubulin cofactor a (TBCA) is regulated by a non-coding antisense TbcA RNA during testis maturation. *PLoS One* 2012; 7:e42536.
266. Lu M, Tian H, Cao YX, He X, Chen L, Song X, Ping P, Huang H, Sun F. Downregulation of miR-320a/383-sponge-like long non-coding RNA NLC1-C (narcolepsy candidate-region 1 genes) is associated with male infertility and promotes testicular embryonal carcinoma cell proliferation. *Cell Death Dis* 2015; 6:e1960.
267. Li M, Liu Y, Wang T, Guan J, Luo Z, Chen H, Wang X, Chen L, Ma J, Mu Z, Jiang AA, Zhu L et al. Repertoire of porcine microRNAs in adult ovary and testis by deep sequencing. *Int J Biol Sci* 2011; 7: 1045–1055.
268. Ling YH, Ren CH, Guo XF, Xu LN, Huang YF, Luo JC, Zhang YH, Zhang XR, Zhang ZJ. Identification and characterization of microRNAs

- in the ovaries of multiple and uniparous goats (*Capra hircus*) during follicular phase. *BMC Genomics* 2014; 15:339.
269. Hossain MM, Ghanem N, Hoelker M, Rings F, Phatsara C, Tholen E, Schellander K, Tesfaye D. Identification and characterization of miRNAs expressed in the bovine ovary. *BMC Genomics* 2009; 10:443.
 270. Hong X, Luense LJ, McGinnis LK, Nothnick WB, Christenson LK. Dicer1 is essential for female fertility and normal development of the female reproductive system. *Endocrinology* 2008; 149:6207–6212.
 271. Nagaraja AK, Andreu-Vieyra C, Franco HL, Ma L, Chen R, Han DY, Zhu H, Agno JE, Gunaratne PH, DeMayo FJ, Matzuk MM. Deletion of dicer in somatic cells of the female reproductive tract causes sterility. *Mol Endocrinol* 2008; 22:2336–2352.
 272. Gonzalez G, Behringer RR. Dicer is required for female reproductive tract development and fertility in the mouse. *Mol Reprod Dev* 2009; 76:678–688.
 273. Cui N, Hao G, Zhao Z, Wang F, Cao J, Yang A. MicroRNA-224 regulates self-renewal of mouse spermatogonial stem cells via targeting DMRT1. *J Cell Mol Med* 2016; 20:1503–1512.
 274. Yan G, Zhang L, Fang T, Zhang Q, Wu S, Jiang Y, Sun H, Hu Y. MicroRNA-145 suppresses mouse granulosa cell proliferation by targeting activin receptor IB. *FEBS Lett* 2012; 586:3263–3270.
 275. Dai A, Sun H, Fang T, Zhang Q, Wu S, Jiang Y, Ding L, Yan G, Hu Y. MicroRNA-133b stimulates ovarian estradiol synthesis by targeting Foxl2. *FEBS Lett* 2013; 587:2474–2482.
 276. Zhang Q, Sun H, Jiang Y, Ding L, Wu S, Fang T, Yan G, Hu Y. MicroRNA-181a suppresses mouse granulosa cell proliferation by targeting activin receptor IIA. *PLoS One* 2013; 8:e59667.
 277. Carletti MZ, Fiedler SD, Christenson LK. MicroRNA 21 blocks apoptosis in mouse periovulatory granulosa cells. *Biol Reprod* 2010; 83:286–295.
 278. Lin F, Li R, Pan ZX, Zhou B, Yu DB, Wang XG, Ma XS, Han J, Shen M, Liu HL. miR-26b promotes granulosa cell apoptosis by targeting ATM during follicular atresia in porcine ovary. *PLoS One* 2012; 7:e38640.
 279. Yang X, Zhou Y, Peng S, Wu L, Lin HY, Wang S, Wang H. Differentially expressed plasma microRNAs in premature ovarian failure patients and the potential regulatory function of mir-23a in granulosa cell apoptosis. *Reproduction* 2012; 144:235–244.
 280. Kitahara Y, Nakamura K, Kogure K, Minegishi T. Role of microRNA-136-3p on the expression of luteinizing hormone-human chorionic gonadotropin receptor mRNA in rat ovaries. *Biol Reprod* 2013; 89:114.
 281. Yin M, Lu M, Yao G, Tian H, Lian J, Liu L, Liang M, Wang Y, Sun F. Transactivation of microRNA-383 by steroidogenic factor-1 promotes estradiol release from mouse ovarian granulosa cells by targeting RBMS1. *Mol Endocrinol* 2012; 26:1129–1143.
 282. Cescon M, Chianese R, Tavares RS. Environmental impact on male (in)fertility via epigenetic route. *J Clin Med* 2020; 9:2520.
 283. Jirtle RL, Skinner MK. Environmental epigenomics and disease susceptibility. *Nat Rev Genet* 2007; 8:253–262.
 284. Xavier MJ, Roman SD, Aitken RJ, Nixon B. Transgenerational inheritance: How impacts to the epigenetic and genetic information of parents affect offspring health. *Hum Reprod Update* 2019; 25:518–540.
 285. Zhang XF, Zhang LJ, Feng YN, Chen B, Feng YM, Liang GJ, Li L, Shen W. Bisphenol A exposure modifies DNA methylation of imprint genes in mouse fetal germ cells. *Mol Biol Rep* 2012; 39:8621–8628.
 286. Ren J, Cheng Y, Ming ZH, Dong XY, Zhou YZ, Ding GL, Pang HY, Rahman TU, Akbar R, Huang HF, Sheng JZ. Intrauterine hyperglycemia exposure results in intergenerational inheritance via DNA methylation reprogramming on F1 PGCs. *Epigenetics Chromatin* 2018; 11:20.
 287. Skinner M, Guerrero-Bosagna C, Haque MM, Nilsson E, Bhandari R, McCarrey J. Environmentally induced transgenerational epigenetic reprogramming of primordial germ cells and subsequent germline. *PLoS One* 2013; 8:1–15 e66318.
 288. Jawaid A, Jehle KL, Mansuy IM. Impact of parental exposure on offspring health in humans. *Trends Genet* 2020; 37:373–388.
 289. Nohara K, Suzuki T, Okamura K. Gestational arsenic exposure and paternal intergenerational epigenetic inheritance. *Toxicol Appl Pharmacol* 2020; 409:115319.
 290. Gao Y, Zhao Y, Zhang H, Zhang P, Liu J, Feng Y, Men Y, Li L, Shen W, Sun Z, Min L. Pubertal exposure to low doses of zearalenone disrupting spermatogenesis through ERalpha related genetic and epigenetic pathways. *Toxicol Lett* 2019; 315:31–38.
 291. Rahman MS, Pang WK, Ryu DY, Park YJ, Pang MG. Multigenerational and transgenerational impact of paternal bisphenol A exposure on male fertility in a mouse model. *Hum Reprod* 2020; 35:1740–1752.
 292. Prados J, Stenz L, Somm E, Strouder C, Dayer A, Paoloni-Giacobino A. Prenatal exposure to DEHP affects spermatogenesis and sperm DNA methylation in a strain-dependent manner. *PLoS One* 2015; 10:e0132136.
 293. Beck D, Sadler-Riggleman I, Skinner MK. Generational comparisons (F1 versus F3) of vinclozolin induced epigenetic transgenerational inheritance of sperm differential DNA methylation regions (epimutations) using MeDIP-Seq. *Environmental Epigenetics* 2017; 3:1, dx016–12.
 294. Skinner MK, Ben Maamar M, Sadler-Riggleman I, Beck D, Nilsson E, McBirney M, Klukovich R, Xie Y, Tang C, Yan W. Alterations in sperm DNA methylation, non-coding RNA and histone retention associate with DDT-induced epigenetic transgenerational inheritance of disease. *Epigenetics Chromatin* 2018; 11:1–24.
 295. Kubsad D, Nilsson EE, King SE, Sadler-Riggleman I, Beck D, Skinner MK. Assessment of glyphosate induced epigenetic transgenerational inheritance of pathologies and sperm Epimutations: Generational toxicology. *Sci Rep* 2019; 9:6372.
 296. McBirney M, King SE, Pappalardo M, Houser E, Unkefer M, Nilsson E, Sadler-Riggleman I, Beck D, Winchester P, Skinner MK. Atrazine induced epigenetic transgenerational inheritance of disease, lean phenotype and sperm Epimutation pathology biomarkers. *PLoS One* 2017; 12:1–37 e0184306.
 297. Wu H, Estill MS, Shershebnv A, Suvorov A, Krawetz SA, Whitcomb BW, Dinnie H, Rahil T, Sites CK, Pilsner JR. Preconception urinary phthalate concentrations and sperm DNA methylation profiles among men undergoing IVF treatment: A cross-sectional study. *Hum Reprod* 2017; 32:2159–2169.
 298. Ouko LA, Shantikumar K, Knezovich J, Haycock P, Schnugh DJ, Ramsay M. Effect of alcohol consumption on CpG methylation in the differentially methylated regions of H19 and IG-DMR in male gametes: Implications for fetal alcohol spectrum disorders. *Alcohol Clin Exp Res* 2009; 33:1615–1627.
 299. Soubry A, Hoyo C, Butt CM, Fieuws S, Price TM, Murphy SK, Stapleton HM. Human exposure to flame-retardants is associated with aberrant DNA methylation at imprinted genes in sperm. *Environ Epigenet* 2017; 3:dx003.
 300. Greeson KW, Fowler KL, Estave PM, Kate Thompson S, Wagner C, Clayton Edenfield R, Symosko KM, Steves AN, Marder EM, Terrell ML, Barton H, Koval M et al. Detrimental effects of flame retardant, PBB153, exposure on sperm and future generations. *Sci Rep* 2020; 10:8567.
 301. Shnorhavorian M, Schwartz SM, Stansfeld B, Sadler-Riggleman I, Beck D, Skinner MK. Differential DNA methylation regions in adult human sperm following adolescent chemotherapy: Potential for epigenetic inheritance. *PLoS One* 2017; 12:1–18 e0170085.
 302. Soubry A, Guo L, Huang Z, Hoyo C, Romanus S, Price T, Murphy SK. Obesity-related DNA methylation at imprinted genes in human sperm: Results from the TIEGER study. *Clin Epigenetics* 2016; 8:51.
 303. Denham J, O'Brien BJ, Harvey JT, Charchar FJ. Genome-wide sperm DNA methylation changes after 3 months of exercise training in humans. *Epigenomics* 2015; 7:717–731.
 304. Zhu L, Liu Y, Xue X, Yuan C, Wang Z. BPA's transgenerational disturbance to transcription of ovarian steroidogenic genes in rare minnow *Gobiocypris rarus* via DNA and histone methylation. *Sci Total Environ* 2020; 762:143055.
 305. Lombo M, Getino-Alvarez L, Depince A, Labbe C, Herraes MP. Embryonic exposure to bisphenol A impairs primordial germ cell migration without jeopardizing male breeding capacity. *Biomolecules* 2019; 9:307.
 306. Li H, Zhang P, Zhao Y, Zhang H. Low doses of carbendazim and chlorothalonil synergized to impair mouse spermatogenesis through epigenetic pathways. *Ecotoxicol Environ Saf* 2020; 188:109908.

307. Gely-Pernot A, Hao C, Legoff L, Multigner L, D'Cruz SC, Kervarrec C, Jegou B, Tevosian S, Smagulova F. Gestational exposure to chlordecone promotes transgenerational changes in the murine reproductive system of males. *Sci Rep* 2018; 8:10274.
308. Legoff L, Dali O, D'Cruz SC, Suglia A, Gely-Pernot A, Hemery C, Kernanec PY, Demmouche A, Kervarrec C, Tevosian S, Multigner L, Smagulova F. Ovarian dysfunction following prenatal exposure to an insecticide, chlordecone, associates with altered epigenetic features. *Epigenetics Chromatin* 2019; 12:29.
309. Wu XF, Yuan HJ, Li H, Gong S, Lin J, Miao YL, Wang TY, Tan JH. Restraint stress on female mice diminishes the developmental potential of oocytes: Roles of chromatin configuration and histone modification in germinal vesicle stage oocytes. *Biol Reprod* 2015; 92: 13.
310. Torres-Flores U, Hernandez-Hernandez A. The interplay between replacement and retention of histones in the sperm genome. *Front Genet* 2020; 11:780.
311. Oikawa M, Simeone A, Hormanseder E, Teperek M, Gaggioli V, O'Doherty A, Falk E, Sporniak M, D'Santos C, Franklin VNR, Kishore K, Bradshaw CR et al. Epigenetic homogeneity in histone methylation underlies sperm programming for embryonic transcription. *Nat Commun* 2020; 11:3491.
312. Ben Maamar M, Sadler-Riggleman I, Beck D, Skinner MK. Epigenetic transgenerational inheritance of altered sperm histone retention Sites. *Sci Rep* 2018; 5308:5301–5310.
313. Hamad MF, Shelko N, Kartarius S, Montenarh M, Hammadeh ME. Impact of cigarette smoking on histone (H2B) to protamine ratio in human spermatozoa and its relation to sperm parameters. *Andrology* 2014; 2:666–677.
314. Lettieri G, D'Agostino G, Mele E, Cardito C, Esposito R, Cimmino A, Giarra A, Trifuoggi M, Raimondo S, Notari T, Febbraio F, Montano L et al. Discovery of the involvement in DNA oxidative damage of human sperm nuclear basic proteins of healthy Young Men living in polluted areas. *Int J Mol Sci* 2020; 21:4198.
315. Radford EJ, Ito M, Shi H, Corish JA, Yamazawa K, Isganaitis E, Seisenberger S, Hore TA, Reik W, Erkek S, Peters A, Patti ME et al. In utero effects. In utero undernourishment perturbs the adult sperm methylome and intergenerational metabolism. *Science* 2014; 345: 1255903.
316. Ben Maamar M, Sadler-Riggleman I, Beck D, McBirney M, Nilsson E, Klukovich R, Xie Y, Tang C, Yan W, Skinner MK. Alterations in sperm DNA methylation, non-coding RNA expression, and histone retention mediate vinclozolin-induced epigenetic transgenerational inheritance of disease. *Environmental Epigenetics* 2018; 4:1, dvy010–19.
317. Brieno-Enriquez MA, Garcia-Lopez J, Cardenas DB, Guibert S, Cleroux E, Ded L, Hourcade Jde D, Peknicova J, Weber M, Del Mazo J. Exposure to endocrine disruptor induces transgenerational epigenetic deregulation of microRNAs in primordial germ cells. *PLoS One* 2015; 10:e0124296.
318. Dickson DA, Paulus JK, Mensah V, Lem J, Saavedra-Rodriguez L, Gentry A, Pagidas K, Feig LA. Reduced levels of miRNAs 449 and 34 in sperm of mice and men exposed to early life stress. *Transl Psychiatry* 2018; 8:101.
319. Gapp K, Jawaaid A, Sarkies P, Bohacek J, Pelczar P, Prados J, Farinelli L, Miska E, Mansuy IM. Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nat Neurosci* 2014; 17:667–669.
320. Gapp K, van Steenwyk G, Germain PL, Matsushima W, Rudolph KLM, Manuella F, Roszkowski M, Vernaz G, Ghosh T, Pelczar P, Mansuy IM, Miska EA. Alterations in sperm long RNA contribute to the epigenetic inheritance of the effects of postnatal trauma. *Mol Psychiatry* 2020; 25:2162–2174.
321. Marczylo EL, Amoako AA, Konje JC, Gant TW, Marczylo TH. Smoking induces differential miRNA expression in human spermatozoa: A potential transgenerational epigenetic concern? *Epigenetics* 2012; 7:432–439.
322. Lopez P, Castro A, Florez M, Miranda K, Aranda P, Sanchez-Gonzalez C, Llopis J, Arredondo M. miR-155 and miR-122 expression of spermatozoa in obese subjects. *Front Genet* 2018; 9:175.
323. Donkin I, Versteyhe S, Ingerslev LR, Qian K, Mechta M, Nordkap L, Mortensen B, Appel EV, Jorgensen N, Kristiansen VB, Hansen T, Workman CT et al. Obesity and bariatric surgery drive epigenetic variation of spermatozoa in humans. *Cell Metab* 2015; 23:369–78.
324. Skinner MK. What is an epigenetic transgenerational phenotype? F3 or F2. *Reprod Toxicol* 2008; 25:2–6.
325. Rodgers AB, Morgan CP, Leu NA, Bale TL. Transgenerational epigenetic programming via sperm microRNA recapitulates effects of paternal stress. *Proc Natl Acad Sci U S A* 2015; 112:13699–13704.
326. Carvan MJr, Kalluvila TA, Klingler RH, Larson JK, Pickens M, Mora-Zamorano FX, Connaughton VP, Sadler-Riggleman I, Beck D, Skinner MK. Mercury-induced epigenetic transgenerational inheritance of abnormal neurobehavior is correlated with sperm epimutations in zebrafish. *PLoS One* 2017; 12:1–26 e0176155.
327. Skinner MK, Nilsson E, Sadler-Riggleman I, Beck D, JR MC. Transgenerational sperm DNA methylation Epimutation developmental origins following ancestral Vinclozolin exposure. *Epigenetics* 2019; 14:721–739.
328. Oliva R, Dixon GH. Vertebrate protamine genes and the histone-to-protamine replacement reaction. *Prog Nucleic Acid Res Mol Biol* 1991; 40:25–94.
329. Hammoud SS, Nix DA, Hammoud AO, Gibson M, Cairns BR, Carrell DT. Genome-wide analysis identifies changes in histone retention and epigenetic modifications at developmental and imprinted gene loci in the sperm of infertile men. *Hum Reprod* 2011; 26:2558–2569.
330. Skinner MK. A new kind of inheritance. *Sci Am* 2014; 311:44–51.

Challenging dogmas: How transgenerational epigenetics reshapes our views on life

Harrison D. Wang¹ | Patrick Allard^{1,2} 

¹The Institute for Society and Genetics, University of California, Los Angeles, California, USA

²Molecular Biology Institute, University of Los Angeles, Los Angeles, California, USA

Correspondence

Patrick Allard, 621 Charles E. Young Dr S., Life Science Bldg 3360, University of California, Los Angeles, CA 90095.
Email: pallard@ucla.edu

Funding information

Burroughs Wellcome Fund; John Templeton Foundation, Grant/Award Number: 60742; National Institute of Environmental Health Sciences, Grant/Award Number: ES027487; NIEHS

Abstract

The emergence of the field of transgenerational epigenetics inheritance (TEI) has profoundly reshaped our understanding of the relationships between environment, soma, and germ cells as well as of heredity. TEI refers to the changes in chromatin state, gene expression, and/or phenotypes that are transmitted across several generations without involving changes to the DNA sequences. TEI has direct connections with, and feeds from, the fields of molecular biology, genetics, developmental biology, and reproductive biology, among others. However, the expansion of TEI-related research, has profoundly reshaped boundaries within each field and often led to the erosion of theories and concepts considered as tenets of biology. We first explore how the molecularization of biology has shifted the definition of epigenetics to include the notion of heredity and how epigenetics has refined our understanding of the central dogma of biology. The demonstrated transfer of environmental information from soma to germ cell through extracellular vesicles and subsequent alteration of health outcomes in offspring has put a definite end to the long-held principle of the Weismann barrier. TEI has also simultaneously led to the revival of the inheritance of acquired characteristics while further eroding the concept of an epigenetic “blank slate” in mammals. Using an historical framework, and via the exploration of central studies in the field, in this perspective article, we will draw a compelling argument for the revolutionary aspect of TEI in biology.

KEYWORDS

transgenerational inheritance, TEI, epigenetics

1 | INTRODUCTION

Transgenerational epigenetics inheritance (TEI) can be defined as changes in chromatin state, gene expression, and/or phenotypes that are transmitted across several generations without involving changes to the DNA sequences. This subfield of epigenetics has grown tremendously in the last decade due to major findings in a variety of model systems that have helped reshape the molecular understanding of heredity. As we will argue, TEI has also been

able to find its place as a field of study because of the erosion of long-held views and dogmas in biology. Together TEI and epigenetics have contributed to the overwhelming amount of evidence for the Weismann barrier to be laid to rest, and have encouraged the revisitation and refinement of concepts, such as the central dogma of molecular biology. Here, we will start by framing the discourse through the lens of Waddington's understanding of the relationship between genotype and phenotype and explore how progress in the field of TEI and research on lived experiences,

such as stress and environmental exposure have brought renewed attention to the concept of soft inheritance.

2 | WADDINGTON AND THE DEFINITION OF THE EPIGENOME

In his seminal 1942 publication, "The Epigenotype," Conrad H. Waddington described the study of mechanisms that exist between genotypes and the phenotypic effects they bring about. Specifically, Waddington provided a prototype for a definition of epigenetics by stating that it refers to "the causal mechanisms at work [...] linked together in a network [...] by which the genes of the genotype bring about phenotypic effects." (Waddington, 2012). Thus, Waddington theorized, in a way akin to modern systems biology, that the developmental processes existing in this space form a network such that early perturbations could funnel to downstream consequences on phenotype. Waddington mentions the example of the "gray-lethal" that prohibits the absorption of bone which has repercussions on other body systems (Grunberg, 1938). For the teeth in particular, impaired coordination and delayed calcification prevents the proper mastication of food. The lack of bone absorption also creates added pressure on nerves in the lower jaw and prevents the animal from taking liquids, leading to starvation and death. Thus, the spontaneous "gray-lethal" mutation of the gene disrupts bone absorption which in turn leads to neuralgic pain and death. From our current perspective, this example has very little to do with our present use of the term epigenetics.

Waddington's concept of the epigenotype was deeply influenced by the current of thoughts in embryology and the debates around the nature and position of the components responsible for carrying out an organism's developmental plan (Felsenfeld, 2014). Obviously, since Waddington's original framework, and with the molecularization of biology, the definition of epigenetics has shifted from its focus on genotype-phenotype interactions towards the stability of expression states and cellular inheritance. It was verified that DNA methylation sites were palindromic, and that DNA methyltransferases were responsible for methylation of unmodified or hemi-methylated DNA (reviewed in Goll & Bestor, 2005). Importantly, these DNA methylation marks are copied on daughter strands after replication, which results in the transmission of the methylated state to future cellular generations (Goll & Bestor, 2005). This finding has led to the insertion of heritability into the definition of epigenetics and, in 1996, Riggs reinterpreted epigenetics as "the study of mitotically or meiotically heritable changes in gene function that cannot be explained by changes in DNA sequence" which has become the near standard definition of epigenetics (Riggs & Porter, 1996). Thus, the shift in definition of epigenetics opened a space for TEI to grow as an area of investigation: if the epigenetic marks are heritable then what molecular mechanisms exist that allow them to be transmitted? If some epigenetic marks are stable over several or many generations, and considering that DNA mutations are reversible, then what truly separates DNA-based heredity from TEI?

3 | REVISITING THE CENTRAL DOGMA OF MOLECULAR BIOLOGY

As the definition of epigenetics shifted, so did the central dogma of biology. First defined by Francis Crick in 1957, the central dogma explains the direction of flow for genetic information in our cells. In its purest form, it translates to "DNA makes RNA, and RNA makes protein." (Crick, 1958, 1970). Crick also specified that "the transfer of information from nucleic acid to nucleic acid, or from nucleic acid to protein may be possible, but transfer from protein to protein, or from protein to nucleic acid is impossible." However, epigenetic pathways conflict with this rigidity and have transformed our interpretation of the genome at large. By altering the patterns of DNA methylation and that of histone modifications and noncoding RNAs (ncRNAs), the epigenetic machinery of writers, erasers, and readers directs in what cellular context, and when, specific genes should be expressed (Gillette & Hill, 2015). Similarly, messenger RNAs (mRNAs) are also subject to posttranslational modification to both modulate gene expression and control their metabolism (Frye et al., 2018). For example, established by a multiprotein writer complex, N⁶-methyladenosine (m⁶A) is the most abundant modification of eukaryotic mRNAs and directly and indirectly affects the binding of various reader proteins, which either target mRNAs for degradation or translation (Yue et al., 2015). Recent studies have focused on the role of m⁶A during development for embryonic and adult stem cell differentiation (Roundtree et al., 2017). Specifically, m⁶A marks transcripts that contain vital developmental regulators to ensure proper transcriptome switching during cell fate transitions. Other work indicates that m⁶A could interact with chromatin regulatory complexes and long noncoding RNAs to influence transcription as well (Patil et al., 2016).

Finally, ncRNAs also play a central role in the epigenetic regulation of chromatin states. In the fission yeast *S. pombe*, small interfering RNA targets and silences the repetitive pericentromeric region (Reinhart & Bartel, 2002). To accomplish this, three interacting protein complexes are involved. The RITS complex first determines the genomic location of the heterochromatin (Li et al., 2009). Then, the RNA-directed RNA polymerase complex (RDRC) amplifies small RNAs from the selected locus (Motamedi et al., 2004). Finally, the Clr4-containing complex establishes the heterochromatic mark (Zhang et al., 2008). Counterintuitively, some level of transcription is necessary to maintain silencing of the heterochromatic locus: the pericentromeric transcripts initially recognized by the RITS complex are synthesized into double-stranded RNA and processed into new small interfering RNAs that load additional RITS complexes, creating a reinforcing loop (Colmenares et al., 2007). The RITS complex also recruits other writers to promote histone deacetylation and H3K9 methylation (Moazed, 2009). In another well-understood example, in female mammals, the long noncoding RNA Xist plays a vital role in X chromosome inactivation. Xist-mediated silencing requires the presence of Xist A-repeats, which are structurally conserved (Wutz et al., 2002). As Xist spreads, it

directly recruits polycomb repressive complex 2 that mediates the tri-methylation of H3K27 (Engreitz et al., 2013).

Thus, while sequence-based information may follow the unidirectional nature of Crick's central dogma, epigenetic mechanisms significantly add complexity to, and transform, the absolute linear schematic of the central dogma to resemble a molecularized version of Waddington's network.

4 | THE END OF THE WEISMANN BARRIER

While other concepts and dogmas have been refined by epigenetics and TEI over time, none has been affected as much as the Weismann barrier (Bline et al., 2020). The Weismann Barrier refers to the unidirectional and irreversible flow of developmental potential from germline to soma and has informed discussion about evolutionary genetic inheritance for much of the past century. However, recent transgenerational epigenetic inheritance research has refuted core tenets of the theory and has indicated a larger role for environmental influence than previously held. The concept of a barrier originates from Weismann's germ plasm theory, which he used to justify how organisms remain relatively constant through generations but adapt to their environment over evolutionary time (Weismann, 1893). He argued that after fertilization, a zygote had two paths for replication. The first choice was "embryonic" cell division, where daughter cells received some parts of the nuclear content. The resultant "idioplasm" gave rise to bodily tissues of the organism. The second choice was "ordinary" cell division, where the daughter cell received all nuclear contents of the parent. Here, a "reserve" germ plasm was produced, to be further transmitted to new generations. Importantly, Weismann postulated that idioplasm could under no circumstances be reconstructed into germ plasm. Therefore, any bodily changes brought about by external stressors during an organism's lifespan would not be passed on to their children. This inability of idioplasm to revert to germ plasm and be inherited by offspring necessitated the existence of a barrier that prevented interaction between the two, which was later named the Weismann Barrier.

While the Weismann Barrier hinges on the separation of germ cells and somatic cells, recent studies have demonstrated that the two interact and exchange epigenetic information to create an additional source of heredity shaped by the parental environment. Multiple groups have now demonstrated and characterized the transfer of small RNAs between somatic and germline cells in mammals (Bohacek & Mansuy, 2015; Jawaid et al., 2018). For instance, in mice, metabolic tracing successfully verified that caput epididymosomes transport small RNA cargos initially synthesized in the epididymal epithelium to spermatozoa (Sharma et al., 2018). The content of these cargos is altered based on the various stress conditions or dietary perturbations encountered by the father pre-conception. With regard to models of stress, zygotes injected with sperm RNAs from males subjected to the MSUS paradigm – maternal separation coupled with unpredictable maternal stress – give rise to

offspring that exhibit depression-like symptoms. These offspring display an increased tendency to remain afloat during forced swim tests, which has been established as an accurate measure of passive coping (Sharma & Rando, 2014). Remarkably, in other studies, the same increase in floating time can be elicited transgenerationally (until the F3) when MSUS is applied during gestation (van Steenwyk et al., 2018). The ability for sperm to carry experiential information to the offspring is not limited to stress-related cues. Offspring generated from mice raised on high-fat diets show impaired glucose metabolism and a small 30–40 nt sperm RNA (tsRNAs) fractions was shown to be the primary mediator of such effects (Chen et al., 2016). Interestingly, the tsRNA cargo requires itself epigenetic modification as demonstrated by the dependence of the transmission of impaired glucose metabolism on active tRNA methyltransferase DNMT2 (Zhang et al., 2018), again highlighting the complex interplay and layering of epigenetic information.

In a recent elegant set of experiments, van Steenwyk and colleagues showed that early life trauma in mice and humans raises the levels of specific serum lipid metabolite species that have the ability to activate PPAR γ in sperm. Interestingly, injection in male mice of serum from mice subjected to MSUS or of a PPAR γ agonist is sufficient to recreate the MSUS-induced glycemic deregulation in offspring (van Steenwyk et al., 2020). Together, these tantalizing developments in the field of TEI go exactly against the concept of the Weismann barrier: external influences on the soma can be physically transferred to the germ cells and be inherited by the offspring.

5 | THE INHERITANCE OF ACQUIRED CHARACTERISTICS AND THE RETURN OF SOFT INHERITANCE

While new advances in the field of TEI are putting some old concepts to rest, they are concomitantly reviving others that had largely been abandoned. In interesting ways, TEI resembles the concepts underlying the theory of inheritance of acquired characteristics, a common belief among eighteenth and nineteenth-century naturalists until it was mostly discarded for Darwin's views of evolution and the Mendel, Boveri-Sutton, Morgan principles of heredity. The theory of inheritance of acquired characteristics, or the ability for acquired characteristics during an organism's lifetime to be passed to their offspring, has long been associated with Jean Baptiste-Lamarck, even though most of the underlying principles of the theory were not his own (Burkhardt, 2013). Instead, Lamarck simply delineated specific conditions necessary for it to occur. Unlike his contemporaries, he argued that organisms took on new forms because of environmentally-shaped habits they acquired and not vice versa. To support this, Lamarck described how the webbed feet of the wading bird resulted from its continual exposure to sinking in mire when searching for prey. The bird will get in the habit of contracting its legs to elongate itself, and after successive generations this habit will be biology encoded in the species.

Conversely, he postulated that constant disuse of an organ or part would cause it to deteriorate until it finally disappears.

While these ideas were eventually dismissed in favor of Darwinian views of evolution, the growth of TEI has signaled the return of a modernized version of Lamarckism: soft inheritance. Soft inheritance, bolstered by a growing body of evidence that variation in epigenetic states is not random but rather initiated and guided by the environment, posits that environment cues can influence hereditary information. While it remains a challenge to comprehensively identify the mechanisms of epigenetic inheritance from other environmental cues, the aforementioned studies indicate that parental lived experience, such as stress can impact the health of descendants spanning several generations.

Interestingly, for soft inheritance and TEI to align at the molecular level, yet another concept needed refining: the ability of early mammalian zygotes and early germ cells, termed primordial germ cells, to profoundly reprogram their epigenome, thereby producing an epigenetic “blank slate” (Messerschmidt et al., 2014). However, this “blank slate” specifically refers to the waves of DNA demethylation during early development while other epigenetic marks do not see reprogramming to a similar extent. Furthermore, even DNA methylation is not completely lost, it is reduced to around 10%, instead being actively maintained at critical genomic regions, such as repetitive elements (Tang et al., 2016).

The ability of environmental information to cause detectable and heritable alterations to the epigenome has now been described in a multitude of model organisms and involve all three types of epigenetic modifications. For example, in nematodes, the plastic component Bisphenol A causes transgenerational reproductive defects that are dependent on two histone modifications, H3K9me3 and H3K27me3 (Camacho et al., 2018; Weinhouse et al., 2018). Endocrine disruptors also lead to detectable transgenerational changes in DNA methylation in Zebrafish (Akemann et al., 2020) as well as in rats (Gillette et al., 2018). In the latter study, the authors examined the number of differentially methylated regions (DMRs) occurring in sperm and specific brain nuclei involved in stress response and social behavior caused by in utero exposure to two endocrine disruptors Vinclozolin and the polychlorinated biphenyl mixture A1221. They observed a high number of DMRs in the sperm, and to a lesser extent also in the brain regions, of male F1 and F3 progeny (Gillette et al., 2018). Remarkably, there was substantial overlap in the number and direction (hyper vs hypo-methylation) of the DMRs in sperm between F1 and F3 generations. This study opens exciting avenues to investigate whether in mammals, DMRs are maintained “as is” across generation or whether a relay-mechanism exist that would involve cross-talks with other epigenetic marks or other mechanisms.

6 | CONCLUSION

While the examples we have examined in this paper have been confined to stress, diet, and environmental chemicals, the field of TEI encompasses many other types of environmental information. However, the field still suffers from a relative fragmentation of the research that forms it:

fragmentation by model organism, type of epigenetic modification, environmental cue, and sex. Questions of the applicability of the model organisms to humans are also not to be ignored, and continued examination of human cohorts, such as the often-cited example of the Dutch Hunger Winter study (Stein et al., 1975), while difficult, will provide critical proof of the relevance of findings in other model systems. Nonetheless, advances in the fields of epigenetics and TEI have already reshaped, or put to rest, numerous views previously considered as rules and/or central to biology.

ACKNOWLEDGMENTS

Patrick Allard is supported by the John Templeton Foundation, NIEHS R01 ES027487, and the Burroughs Wellcome Fund.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

ORCID

Patrick Allard  <http://orcid.org/0000-0001-7765-1547>

REFERENCES

- Akemann, C., Meyer, D. N., Gurdziel, K., & Baker, T. R. (2020). TCDD-induced multi- and transgenerational changes in the methylome of male zebrafish gonads. *Environmental Epigenetics*, 6(1), dvaa010. <https://doi.org/10.1093/eep/dvaa010>
- Bline, A. P., Le Goff, A., & Allard, P. (2020). What is lost in the Weismann barrier? *Journal of Developmental Biology*, 8(4), 35. <https://doi.org/10.3390/jdb8040035>
- Bohacek, J., & Mansuy, I. M. (2015). Molecular insights into transgenerational non-genetic inheritance of acquired behaviours. *Nature Reviews Genetics*, 16(11), 641–652. <https://doi.org/10.1038/nrg3964>
- Burkhardt, R. W., Jr (2013). Lamarck, evolution, and the inheritance of acquired characters. *Genetics*, 194(4), 793–805. <https://doi.org/10.1534/genetics.113.151852>
- Camacho, J., Truong, L., Kurt, Z., Chen, Y. W., Morselli, M., Gutierrez, G., Pellegrini, M., Yang, X., & Allard, P. (2018). The memory of environmental chemical exposure in *C. elegans* is dependent on the Jumonji demethylases *jmjd-2* and *jmjd-3/utx-1*. *Cell Reports*, 23(8), 2392–2404. <https://doi.org/10.1016/j.celrep.2018.04.078>
- Chen, Q., Yan, M., Cao, Z., Li, X., Zhang, Y., Shi, J., Feng, G. H., Peng, H., Zhang, X., Zhang, Y., Qian, J., Duan, E., Zhai, Q., & Zhou, Q. (2016). Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science*, 351(6271), 397–400. <https://doi.org/10.1126/science.aad7977>
- Colmenares, S. U., Buker, S. M., Buhler, M., Dlakić, M., & Moazed, D. (2007). Coupling of double-stranded RNA synthesis and siRNA generation in fission yeast RNAi. *Molecular Cell*, 27(3), 449–461. <https://doi.org/10.1016/j.molcel.2007.07.007>
- Crick, F. H. (1958). On protein synthesis. *Symposia of the Society for Experimental Biology*, 12(No. 138-63), 8.
- Crick, F. H. (1970). Central dogma of molecular biology. *Nature*, 227(5258), 561–563. <https://doi.org/10.1038/227561a0>
- Engreitz, J. M., Pandya-Jones, A., McDonel, P., Shishkin, A., Sirokman, K., Surka, C., Kadri, S., Xing, J., Goren, A., Lander, E. S., Plath, K., & Guttman, M. (2013). The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. *Science*, 341(6147), 1237973. <https://doi.org/10.1126/science.1237973>
- Felsenfeld, G. (2014). A brief history of epigenetics. *Cold Spring Harbor Perspectives in Biology*, 6(1), a018200. <https://doi.org/10.1101/cshperspect.a018200>

- Frye, M., Harada, B. T., Behm, M., & He, C. (2018). RNA modifications modulate gene expression during development. *Science*, 361(6409), 1346–1349. <https://doi.org/10.1126/science.aau1646>
- Gillette, R., Son, M. J., Ton, L., Gore, A. C., & Crews, D. (2018). Passing experiences on to future generations: Endocrine disruptors and transgenerational inheritance of epimutations in brain and sperm. *Epigenetics*, 2018 13(10-11), 1106–1126. <https://doi.org/10.1080/15592294.2018.1543506>
- Gillette, T. G., & Hill, J. A. (2015). Readers, writers, and erasers: Chromatin as the whiteboard of heart disease. *Circulation Research*, 116(7), 1245–1253. <https://doi.org/10.1161/CIRCRESAHA.116.303630>
- Goll, M. G., & Bestor, T. H. (2005). Eukaryotic cytosine methyltransferases. *Annual Review of Biochemistry*, 74, 481–514. <https://doi.org/10.1146/annurev.biochem.74.010904.153721>
- Grünberg, H. (1938). Some new data on the grey-lethal mouse. *Journal of Genetics*, 36(1), 153–170.
- Jawaid, A., Roszkowski, M., & Mansuy, I. M. (2018). Transgenerational epigenetics of traumatic stress. *Progress in Molecular Biology and Translational Science*, 158, 273–298. <https://doi.org/10.1016/bs.pmbts.2018.03.003>
- Li, H., Motamedi, M. R., Yip, C. K., Wang, Z., Walz, T., Patel, D. J., & Moazed, D. (2009). An alpha motif at Tas3 C terminus mediates RITS cis spreading and promotes heterochromatic gene silencing. *Molecular Cell*, 34(2), 155–167. <https://doi.org/10.1016/j.molcel.2009.02.032>
- Messerschmidt, D. M., Knowles, D. B., & Solter, D. (2014). DNA methylation dynamics during epigenetic reprogramming in the germline and preimplantation embryos. *Genes and Development*, 28(8), 812–828. <https://doi.org/10.1101/gad.234294.113>
- Moazed, D. (2009). Small RNAs in transcriptional gene silencing and genome defence. *Nature*, 457(7228), 413–420. <https://doi.org/10.1038/nature07756>
- Motamedi, M. R., Verdel, A., Colmenares, S. U., Gerber, S. A., Gygi, S. P., & Moazed, D. (2004). Two RNAi complexes, RITS and RDRC, physically interact and localize to noncoding centromeric RNAs. *Cell*, 119(6), 789–802. <https://doi.org/10.1016/j.cell.2004.11.034>
- Patil, D. P., Chen, C. K., Pickering, B. F., Chow, A., Jackson, C., Guttman, M., & Jaffrey, S. R. (2016). m(6)A RNA methylation promotes XIST-mediated transcriptional repression. *Nature*, 537(7620), 369–373. <https://doi.org/10.1038/nature19342>
- Reinhart, B. J., & Bartel, D. P. (2002). Small RNAs correspond to centromere heterochromatic repeats. *Science*, 297, 1831.
- Riggs, A. D., & Porter, T. N. (1996). Overview of epigenetic mechanisms. *Cold Spring Harbor Monograph Series*, 32, 29–46.
- Roundtree, I. A., Evans, M. E., Pan, T., & He, C. (2017). Dynamic RNA modifications in gene expression regulation. *Cell*, 169(7), 1187–1200. <https://doi.org/10.1016/j.cell.2017.05.045>
- Sharma, U., & Rando, O. J. (2014). Father-son chats: Inheriting stress through sperm RNA. *Cell Metabolism*, 19(6), 894–895. <https://doi.org/10.1016/j.cmet.2014.05.015>
- Sharma, U., Sun, F., Conine, C. C., Reichholz, B., Kukreja, S., Herzog, V. A., Ameres, S. L., & Rando, O. J. (2018). Small RNAs are trafficked from the epididymis to developing mammalian sperm. *Developmental Cell*, 46(4), 481–494.e6. <https://doi.org/10.1016/j.devcel.2018.06.023>
- van Steenwyk, G., Roszkowski, M., Manuella, F., Franklin, T. B., & Mansuy, I. M. (2018). Transgenerational inheritance of behavioral and metabolic effects of paternal exposure to traumatic stress in early postnatal life: evidence in the 4th generation. *Environmental Epigenetics*, 4(2), dv023. <https://doi.org/10.1093/eep/dvy023>
- van Steenwyk, G., Gapp, K., Jawaid, A., Germain, P. L., Manuella, F., Tanwar, D. K., Zamboni, N., Gaur, N., Efimova, A., Thumfart, K. M., Miska, E. A., & Mansuy, I. M. (2020). Involvement of circulating factors in the transmission of paternal experiences through the germline. *The EMBO Journal*, 39(23), e104579. <https://doi.org/10.15252/embj.2020104579>
- Stein, Z., Susser, M., Saenger, G., & Marolla, F. (1975). *Famine and human development: The Dutch hunger winter of 1944–1945*. Oxford University Press.
- Tang, W. W. C., Kobayashi, T., Irie, N., Dietmann, S., & Surani, M. A. (2016). Specification and epigenetic programming of the human germ line. *Nature Reviews Genetics*, 2016 Oct 17(10), 585–600. <https://doi.org/10.1038/nrg.2016.88>
- Waddington, C. H. (2012). The epigenotype. 1942. *International Journal of Epidemiology*, 41(1), 10–13. <https://doi.org/10.1093/ije/dyr184>
- Weinhouse, C., Truong, L., Meyer, J. N., & Allard, P. *Caenorhabditis elegans* as an emerging model system in environmental epigenetics. *Environmental and Molecular Mutagenesis*, 59(7), 560–575.
- Weismann, A. (1893). *The germplasm: A theory of heredity*. Charles Scribner's Sons.
- Wutz, A., Rasmussen, T. P., & Jaenisch, R. (2002). Chromosomal silencing and localization are mediated by different domains of Xist RNA. *Nature Genetics*, 30(2), 167–174. <https://doi.org/10.1038/ng820>
- Yue, Y., Liu, J., & He, C. (2015). RNA N6-methyladenosine methylation in post-transcriptional gene expression regulation. *Genes and Development*, 29(13), 1343–1355. <https://doi.org/10.1101/gad.262766.115>
- Zhang, K., Mosch, K., Fischle, W., & Grewal, S. I. S. (2008). Roles of the Ctr4 methyltransferase complex in nucleation, spreading and maintenance of heterochromatin. *Nature Structural & Molecular Biology*, 15(4), 381–388. <https://doi.org/10.1038/nsmb.1406>
- Zhang, Y., Zhang, X., Shi, J., Tuorto, F., Li, X., Liu, Y., Liebers, R., Zhang, L., Qu, Y., Qian, J., Pahima, M., Liu, Y., Yan, M., Cao, Z., Lei, X., Cao, Y., Peng, H., Liu, S., Wang, Y., ... Chen, Q. (2018). Dnmt2 mediates intergenerational transmission of paternally acquired metabolic disorders through sperm small non-coding RNAs. *Nature Cell Biology*, 20(5), 535–540. <https://doi.org/10.1038/s41556-018-0087-2>

How to cite this article: Wang, H. D., & Allard, P. (2022). Challenging dogmas: how transgenerational epigenetics reshapes our views on life. *J. Exp. Zool*, 337, 70–74. <https://doi.org/10.1002/jez.2465>

"Systems Biology of Reproduction"

Spring 2024 (Even Years) – Course Syllabus
 Biol 475/575 Undergraduate/Graduate (3 Credit)
 SLN: (475) – 06763, (575) – 06764
 Time - Tuesday and Thursday 10:35 am-11:50 am
 Course Lectures in person and recorded on Canvas/Panopto and Discussion Sessions live in person and on WSU Zoom for all campuses (Hybrid Course)
 Room – CUE 418
 Course Director – Michael Skinner, Abelson Hall 507, 335-1524, skinner@wsu.edu
 Co-Instructor – Eric Nilsson, Abelson Hall 507, 225-1835, nilsson@wsu.edu
Learning Objective -
 Current literature based course on the Systems Biology of Reproduction. Learning Systems approaches to the biology of reproduction from a molecular to physiological level of understanding.

Schedule/Lecture Outline –

| | | | |
|----------|---------|---------|---|
| January | 9 & 11 | Week 1 | Systems Biology Introduction |
| | 16 & 18 | Week 2 | Molecular/ Cellular/ Reproduction Systems |
| | 23 & 25 | Week 3 | Sex Determination Systems |
| Jan /Feb | 30 & 1 | Week 4 | Male Reproductive Tract Development & Function |
| February | 6 & 8 | Week 5 | Female Reproductive Tract Development & Function |
| | 13 & 15 | Week 6 | Gonadal Developmental Systems Biology |
| | 20 & 22 | Week 7 | Testis Systems Biology |
| | 27 & 29 | Week 8 | Ovary Systems Biology |
| March | 5 & 7 | Week 9 | Epigenetics and Transgenerational Gonadal Disease |
| | 11 – 15 | Week 10 | Spring Break |
| | 19 & 21 | Week 11 | Gametogenesis/ Stem Cells/ Cloning |
| | 26 & 28 | Week 12 | Hypothalamus- Pituitary Development & Function |
| April | 2 & 4 | Week 13 | Reproductive Endocrinology Systems |
| | 9 & 11 | Week 14 | Fertilization & Implantation Systems |
| | 16 & 18 | Week 15 | Fetal Development & Birth Systems |
| | 23 & 25 | Week 16 | Assisted Reproduction/Contraception |
| Apr/May | 30 & 2 | Week 17 | Exam or Grant Review |

Spring 2024 – Systems Biology of Reproduction Lecture Outline – Epigenetics and Transgenerational Reproductive Disease

Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday

March 5, 2024

Week 9

Epigenetics and Transgenerational Reproductive Disease

- Environmental Induced Pathology
- Transgenerational Disease Phenotype
- Compound Specificity
- Epigenetic Mechanism
- Epimutations and Exposure Specificity
- Transgenerational Transcriptome
- Transgenerational Testis Disease
- Transgenerational Ovary Disease
- Broader Impact

REQUIRED READING

Ben Maamar M, Nilsson EE, Skinner MK. Epigenetic transgenerational inheritance, gametogenesis and germline development. *Biol Reprod.* 2021 Sep 14;105(3):570-592.

Wang HD, Allard P. Challenging dogmas: How transgenerational epigenetics reshapes our views on life. *J Exp Zool A Ecol Integr Physiol.* 2022 Jan;337(1):70-74.

Spring 2024 – Systems Biology of Reproduction
 Discussion Outline – Epigenetics and Transgenerational Reproductive Disease
 Michael K. Skinner – Biol 475/575
 CUE 418, 10:35-11:50 am, Tuesday & Thursday
 March 7, 2024
 Week 9

Epigenetics and Transgenerational Reproductive Disease

Primary Papers:

1. Kubsad, et al. (2019) *Scientific Reports* 9 :6372
2. Sadler-Riggelman, et al. (2019) *Environmental Epigenetics* 5(3) dvz013
3. Beck, et al. (2021) *Epigenetics & Chromatin* 14:6

Discussion

Student 11: Reference 1 above

- What transgenerational reproductive phenotypes were observed?
- What is glyphosate and effects on sperm?
- What is generational toxicology?


Student 1: Reference 2 above

- What environmental exposures were used?
- What was the Sertoli cell effects observed?
- What basic information on testis disease was obtained?

Student 2: Reference 3 above

- What technology and epigenetic analysis was used?
- What differential epigenetic regions were observed?
- What transgenerational integration of epigenetic processes was observed?

GENETIC DETERMINISM




ENVIRONMENTAL IMPACT ON BIOLOGY

- Regional Disease Frequencies
- Low Frequency of Genetic Component of Disease
- Increases In Disease Frequencies
- Identical Twins and Variable Disease Frequency
- Environmental Exposures and Disease
- Evolution and Rapid Induction

GROWING SICKNESS

Although people are living longer, they are also living with more chronic conditions, as seen here in data for the developed world.


Number of conditions ■ 0 ■ 1-2 ■ 3-4 ■ 5-6 ■ 7-8 ■ ≥9




ENVIRONMENTAL IMPACT ON BIOLOGY

- Regional Disease Frequencies
- Low Frequency of Genetic Component of Disease
- Increases In Disease Frequencies
- Identical Twins and Variable Disease Frequency
- Environmental Exposures and Disease
- Evolution and Rapid Induction

GENETIC DETERMINISM




EPIGENETIC SOLUTIONS TO GENETIC DETERMINISM FAILURES



EPIGENETICS

Molecular factors/processes around the DNA that regulate genome activity, independent of DNA sequence, and are mitotically stable


Epigenetics




EPIGENETIC MECHANISMS AND MARKS

- DNA Methylation
- Histone Modifications
- Chromatin Structure
- Non-coding RNA
- RNA methylation

Transgenerational Inheritance of Disease




Model Endocrine Disruptor: Vinclozolin




- Vinclozolin is a systemic fungicide (e.g. Wine Industry)
- Vinclozolin and its metabolites are **anti-androgenic**
- Late embryonic/early postnatal exposure causes abnormal reproductive tract development and gonadal function

Vinclozolin Effects on Spermatogenic Cell Apoptosis



Germ Cell Apoptosis



* P<0.05


Vinclozolin

Complete Male Infertility (10%)


F3 Generation Males

Control

Vinclozolin




Sperm Concentration P60-P150



* P<0.05

Vinclozolin


Sperm Motility P60-P150




* P<0.05

Vinclozolin


Transgenerational Disease in Female F1-F4







F3 Female Puberty



F3 Male Puberty





TRANSGENERATIONAL DISEASE ETIOLOGY

| MALE | FEMALE |
|---|---|
| <ul style="list-style-type: none"> Spermatogenic Defect (>90%) Male infertility (complete ~10%, severe 20%) Prostate disease (~50%) | <ul style="list-style-type: none"> Kidney disease (~30-40%) Increase in mammary tumor formation (~10-20%) Behavior (Mate Preference, Anxiety, & Stress) (>90%) Pubertal Abnormalities (~25%) Obesity (~10-50%) Lean (~10-20%) Immune (~20%) |
| | <ul style="list-style-type: none"> Pre-eclampsia-like during late pregnancy (~10%) Premature Ovarian Failure POF (>90%) Ovarian Polycystic Ovarian Disease (>90%) Female Premature Pubertal Onset (>90%) Parturition (birth) defects |

Compound Specificity


Other Endocrine Disruptors And Environmental Compounds

Environmental Compound Specificity


| (Exposure Groups) | (Direct) F1 | F3 (Transgenerational) |
|---|-------------|------------------------|
| A. Vinclozolin [agricultural fungicide] | Yes | Yes |
| B. Flutamide [anti-androgenic pharmaceutical] | Yes | No |
| C. TCDD/Dioxin (Industrial pollutant) | Yes | Yes |
| D. Plastics Compounds [Bisphenol-A BPA, Phthalate-DEHP & DBP] | Yes | Yes |
| E. Jet Fuel [JP8] (Hydrocarbon Mixture) | Yes | Yes |
| F. Pesticide & Insect Repellent [Permethrin & DEET] | No* | Yes |
| G. DDT (pesticide) | Yes | Yes |
| H. Methoxychlor (pesticide, replace DDT) | Yes | Yes |
| I. Mercury (Industrial pollutant) | Yes | Yes |
| J. Atrazine (agricultural herbicide) | No* | Yes |
| K. Glyphosate (pesticide herbicide) | No* | Yes |

Assessment of Glyphosate Induced Epigenetic Transgenerational Inheritance of Pathologies and Sperm Epimutations: Generational Toxicology.


Kubsad D, Nilsson EE, King SE, Sadler-Riggelman I, Beck D, Skinner MK. Sci Rep. 2019 Apr 23;9(1):6372.




Effects of glyphosate-based herbicides and glyphosate exposure on sex hormones and the reproductive system: From epidemiological evidence to mechanistic insights.
Tajai P, Pruksakorn D, Chattipakorn SC, Chattipakorn N, Shinlapawittayatorn K.
Environ Toxicol Pharmacol. 2023 Sep;102:104252




Maternal genetic intergenerational and transgenerational effects on hormone synthesis in ovarian granulosa cells of offspring exposed to cadmium during pregnancy.
Luo L, Li J, Sun Y, Lv Y, Liu J, Li Y, Zhang C, Zhang W.
Ecotoxicol Environ Saf. 2023 Sep 15;263:115278.





Di(2-ethylhexyl) phthalate induces reproductive toxicity and transgenerational reproductive aging in *Caenorhabditis elegans*.
Zhang Y, Li J, Shi W, Lu L, Zhou Q, Zhang H, Liu R, Pu Y, Yin L.
Environ Pollut. 2023 Nov 1;336:122259.




The brominated flame retardant, 1,2,5,6-tetrabromocyclooctane (TBCO), causes multigenerational effects on reproductive capacity of Japanese medaka (*Oryzias latipes*).
Devoy C, Raza Y, Kleiner M, Jones PD, Doering JA, Wiseman S.
Chemosphere. 2023 Feb;313:137561.






Transgenerational Disease Etiology

- Spermatogenic Defect (>90%)
- Male infertility (complete ~10%, severe 20%)
- Kidney disease (~30-40%)
- Prostate disease (~50%)
- Increase in mammary tumor formation (~10-20%)
- Behavior (Mate Preference, Anxiety & Stress) (>90%)
- Pre-eclampsia-like during late pregnancy (~10%)
- Premature Ovarian Failure POF (>90%)
- Ovarian Polycystic Ovarian Disease (>90%)
- Female Premature Pubertal Onset (>90%)
- Obesity (~10-50%)





ENVIRONMENTALLY INDUCED EPIGENETIC TRANSGENERATIONAL INHERITANCE

| | |
|--|---|
| Environmental Toxicants | |
| Vinclozolin (Agricultural Fungicide) | Permethrin & DEET (Insect Repellants) |
| Methoxychlor (Agricultural Pesticide) | DDT (Pesticide) |
| Dioxin/TCDD (Industrial Contaminant) | Tributyltin (Industrial Toxicant & Biocide) |
| Plastic Compounds (BPA & Phthalates) | Hydrocarbons (Jet Fuel) |
| Atrazine (Herbicide) | Glyphosate (Pesticide / Herbicide) |
| Other Types Exposures | |
| Nutrition (High Fat or Caloric Restriction) | Smoking & Alcohol |
| Temperature & Drought (Plant Health & Flowering) | Stress Trauma (Behavioral) |

Plants

Flies

Worms

Fish

Birds


Rodents





Pigs

Humans


Summary

- Transient Embryonic Exposure Effects Adult
 - Sex Determination Period
 - Fetal Basis of Disease
- Spermatogenic Fertility Defect & Other Diseases
- Transgenerational Phenotype
 - WHAT MECHANISM?







Chromatin Structure – Composition and Function during Spermiogenesis
 Khochbin, S. and Ward, W.S




Environmentally Induced New Transgenerational Histone Retention





Developmental Origins of Transgenerational Sperm Histone Retention Following Ancestral Exposures
 Ben Maamar M, Beck D, Nilsson E, McCarrey JR, and Skinner MK
 Developmental Biology (submitted)





Ancestral vinclozolin exposure alters the epigenetic transgenerational inheritance of sperm small noncoding RNAs (CC)
 Andrew Schuster, Michael K. Skinner, Wei Yan
 Environmental Epigenetics (2016) 2 (1): 1-15






Small RNAs in the Transgenerational Inheritance of Epigenetic Information.
 Duempelmann L, Skribbe M, Bühler M.
 Trends Genet. 2020 Jan 14. doi: 10.1016/j.tig.2019.12.001. [Epub ahead of print]



Integration of sperm ncRNA-directed DNA methylation and DNA methylation-directed histone retention in epigenetic transgenerational inheritance.
 Beck D, Ben Maamar M, Skinner MK.
 Epigenetics Chromatin. 2021 Jan 12;14(1):6.





Epigenetic Mechanisms of Gene Regulation

Spermatogenesis



- DNA Methylation
- Histone Modification
- Chromatin Structure
- DNA Organization into Domains (eg Loops)
- Nuclear Compartmentalization (eg nuclear matrix)
- Noncoding functional RNAs

Other Mechanisms:

- Replaced Protamines
- Condensation
- Condensation
- Condensation
- Silencing (?)


Epigenetics and male reproduction: the consequences of paternal lifestyle on fertility, embryo development, and children lifetime health.

Stuppia L, Franzago M, Ballerini P, Gatta V, Antonucci I.
Clin Epigenetics. 2015 Nov 11;7:120.




Developmental origins of transgenerational sperm DNA methylation epimutations following ancestral DDT exposure.





Ben Maamar M, Nilsson E, Sadler-Riggleman I, Beck D, McCarey JR, Skinner MK.
Dev Biol. 2019 Jan 15;445(2):280-293.



Transgenerational sperm DNA methylation epimutation developmental origins following ancestral vinclozolin exposure.

Skinner MK, Nilsson E, Sadler-Riggleman I, Beck D, Ben Maamar M, McCarey JR.
Epigenetics. 2019 Jul;14(7):721-739.






F3 Generation Sperm Epigenome Mapping
(Epimutations)
(Epigenetic Biomarkers for Ancestral Exposures)

Transgenerational Sperm Epigenome Alterations
(>200 differential DNA methylation sites)


Potential DNA Sequence Motif Identified (EDM1) & CpG%
(Susceptibility epigenetic transgenerational mark)

Genome Activity Alterations?
(transcriptome)


E16 Testis Transcriptome




Tissue Specific Transgenerational Transcriptomes (F3)




Clustering Tissues Male Female



Epigenetic Control Regions



Epigenetic Control Region





Transgenerational Transcriptomes (F3 Generation)


- Variety Somatic Tissues and Cell Types
- Cell and Tissue Specific Altered Transcriptomes (>10X more genes than epimutations)
- Gene Clusters associate with Epimutations (Epigenetic Control Regions)
- Epimutation transmission to disease?**

Ovarian Granulosa Cell Vinclozolin Lineage F3 Generation Transgenerational Transcriptome (~500 genes)


(A) Ovarian Disease Associated Genes




(B) Polycystic Ovarian Disease Associated Genes




Environmental toxicant induced epigenetic transgenerational inheritance of ovarian pathology and granulosa cell epigenome and transcriptome alterations: ancestral origins of polycystic ovarian syndrome and primary ovarian insufficiency.
 Nilsson E, Klukovich R, Sadler-Riggleman I, Beck D, Xie Y, Yan W, Skinner MK.
 Epigenetics. 2018;13(8):875-895.




Epigenetic Transgenerational Inheritance of Sertoli Cell Abnormalities




Epigenetic transgenerational inheritance of testis pathology and Sertoli cell epimutations: generational origins of male infertility.
 Sadler-Riggleman I, Klukovich R, Nilsson E, Beck D, Xie Y, Yan W, Skinner MK.
 Environ Epigenet. 2019 Aug 29;5(3):dvz013.




Environmental Toxicant Induced Epigenetic Transgenerational Inheritance of Prostate Pathology and Stromal-Epithelial Cell Epigenome and Transcriptome Alterations: Ancestral Origins of Prostate Disease.
 Klukovich R, Nilsson E, Sadler-Riggleman I, Beck D, Xie Y, Yan W, Skinner MK.
 Sci Rep. 2019 Feb 18;9(1):2209.




Adipocyte Epigenetic Alterations and Potential Therapeutic Targets in Transgenerationally Inherited Lean and Obese Phenotypes Following Ancestral Exposures
 King SE, Nilsson E, Beck D, Skinner MK
 Adipocyte 2019 8(1) 362-378




EPIGENETIC TRANSGENERATIONAL INHERITANCE




Epigenetic transgenerational inheritance of parent-of-origin allelic transmission of outcross pathology and sperm epimutations.
 Ben Maamar M, King SE, Nilsson E, Beck D, Skinner MK.
 Dev Biol. 2020 Feb 1;458(1):106-119.



ROLE OF GERM LINE IN EPIGENETIC TRANSGENERATIONAL INHERITANCE



EPIGENETIC AND GENETIC CASCADE OF EVENTS INVOLVED IN DEVELOPMENT



Summary

- Epigenetic Transgenerational Inheritance
 - Non-Genetic Form Inheritance
 - Toxicology / Environmental
 - Molecular Mechanism for Disease Etiology and DOHAD
 - All Epigenetic Processes Involved and Integrated in Germline
 - Generational Toxicology

Environmental Epigenetics, Disease and Evolution

- Integration Epigenetics and Genetics Essential Biology
- Evolution and Disease Etiology Requires Inclusion Epigenetics
- Doom and Gloom/ Biomarkers & Preventative Medicine

Epigenetic Alterations Promote Genetic Instability

| Genetic Mutation | Epigenetic Alteration | DNA Sequence Alteration |
|-----------------------------|---|--|
| Point Mutation (SNP) | DNA Methylation (CpG) | Susceptibility C → T Conversion |
| Copy Number Variation (CNV) | Hypomethylation (Repeats) | Susceptibility Repeat Element Alteration (CNV) |
| Transposon Migration | Hypomethylation DNA | Susceptibility Transposon Migration |
| Translocation | DNA Methylation and Histone Alterations | Susceptibility Translocation at Break Point |
| Telomere Length | DNA Methylation Alteration | Alteration in Telomere Length |

Environmentally Induced Epigenetic Transgenerational Inheritance of Sperm Epimutations Promote Genetic Mutations

Skinner MK, Guerrero-Bosagna C, Haque M. Epigenetics 2015; 10:8, 762-771





Table 1. (A) Vinclozolin F3 Generation Sperm Genome-wide CNV and Epimutations

| Parameters | F1 Generation Sperm CNV | F3 Generation Sperm CNV | F3 Generation Epimutation Sperm |
|-------------------------------|--------------------------|----------------------------|---------------------------------|
| Number (Single Probes) | 540(294 Gain / 246 Loss) | 491(24648 Gain / 264 Loss) | 9932 |
| Number (≥3 Probes) | 39(21 Gain / 18 Loss) | 506(461 Gain / 45 Loss) | 191 |
| Mean Size (base) | 11,633 | 12,637 | 2,131 |
| Mean CpG Density (CpG/100 bp) | 1.1 | 1.0 | 0.9 |


Tertiary Epimutations - A Novel Aspect of Epigenetic Transgenerational Inheritance Promoting Genome Instability
 PLoS One. 2016 Dec 19;11(12):e0168038.
 McCarrey JR, Lehto JD, Raju SS, Wang Y, Nilsson EE, Skinner MK.



Transgenerational Phenotype

| | DNA Mutation | Epigenetic Mutation |
|---------------|--|---------------------|
| Frequency - | <0.01% (Hot Spot 1-5%) | High (30-100%) |
| Reproducible- | Random Event | Highly Reproducible |
| Genetics- | Mendelian (decline frequency generationally) | Non-Mendelian |

Sperm Epimutations Promotes Epigenetic Transgenerational Inheritance of Genetic Mutations



Epigenetics 2015 10:762
 PlosONE 2016 e0168038

Summary

- Epigenetic Transgenerational Inheritance
 - Non-Genetic Form Inheritance
 - Toxicology / Environmental
 - Molecular Mechanism for Disease Etiology and DOHAD
 - All Epigenetic Processes Involved and Integrated in Germline
 - Generational Toxicology




Environmental Epigenetics, Disease and Evolution

- Integration Epigenetics and Genetics Essential Biology
- Evolution and Disease Etiology Requires Inclusion Epigenetics
- Doom and Gloom/ Biomarkers & Preventative Medicine

Evolutionary Biology

- Mechanism-Adaptation (ie DNA mutations)
- Problem- Timing
- Environment- Influence (?How)
- Determinant- Sexual Selection

Collaboration- David Crews, UT Austin



Role of Epigenetics in the Speciation of Darwin Finches and Evolutionary Biology

Goal-


Demonstrate the role of environment and epigenetics as critical molecular factors in evolutionary biology.

Objectives-


-Use Darwin Finches to determine the genetic and epigenetic differences between variety species (ie For, Ful, Spa, Par, Cra).

-Use Chromosomal Genomic Hybridization (CGH) to associate copy number variation between species (genetic differences).





-Use Methylated DNA Immunoprecipitation followed by tiling array (MeDIP-Chip) to associate epimutations between species (epigenetic differences).





Speciation of Darwin Finch- Role Epigenetics and Genetics




Collaboration- Dale Clayton, Utah






| Exposure | Pathology | Reference |
|--|---|--|
| Toxicants | | |
| Vinclozolin | Testis, Prostate, Kidney Disease, Tumors, Immune Gender Specific Changes in Anxiety-like Behavior Immune and Reproductive | Anway, et al., 2005 [3]; 2006 [32] Skinner, et al., 2008 [33] Nilsson, et al., 2008 [34] |
| Methoxychlor | Testis, Kidney, Ovary, Obesity | Anway, et al., 2005 [3], Manikkam, 2014 [35] |
| Permethrin/DEET | Prostate, Kidney Disease | Manikkam, et al., 2012 [36] |
| Dioxin | Prostate, Kidney, Fertility, Pregnancy | Manikkam, et al., 2012 [37] Bruner-Tran, 2011 [38] |
| BPA/Phthalates | Prostate, Kidney, Obesity | Manikkam et al., 2013 [39] |
| Hydrocarbon Mixture (Jet Fuel) | Prostate, Kidney, Obesity, Immune and Reproduction | Tracey et al., 2013 [40] |
| Vinclozolin, Permethrin/DEET, Plastics, Dioxin, Jet Fuel | Polycystic Ovaries, Reduced Primordial Follicle Pool | Nilsson et al., 2012 [41] |
| DDT | Obesity, Kidney, Testis | Skinner, et al., 2013 [5] |
| Phthalate | Testis and Spermatogonial Stem Cell | Doyle, et al., 2013 [42] |
| Tributyltin | Obesity and Adipose Cell | Chamorro-Garcia, et al., 2013 [43] |
| BPA | Social Behavior, Implantation, Litter Size, Sperm | Wolstenholme, et al., 2012 [44]; Sallan, et al., 2009 [45] |
| Others | | |
| Caloric Restriction | Cardiovascular Mortality | Bygren, et al., 2014 [46] |
| High Fat Diet | Growth and Insulin Sensitivity | Dunn and Bale, 2011 [6] |
| Folate | Congenital Malformations | Padmanabhan, et al., 2013 [47] |
| Drought | DNA Methylation Changes | Zheng, et al., 2013 [7] |
| Heat/Salt | Flowering and Salt Tolerance | Suter and Widmer, 2013 [48] |
| Prediabetes | Glucose Tolerance and Insulin Sensitivity | Wei, et al., 2014 [49] |
| Smoking | Abnormal Pulmonary Function | Rehan, 2013 [50] |
| Alcohol | Endocrine and Neuronal Function | Govorko, 2012 [51] |
| Heat Stress | Increased Hsp70 Production and Tolerance to Heat Stress | Norouzitallab, et al., 2014 [8] |




Male germline transmits fetal alcohol adverse effect on hypothalamic proopiomelanocortin gene across generations.

Govorko D, et al. (2012) Biol Psychiatry. 1;72(5):378-88.



Progressive, transgenerational changes in offspring phenotype and epigenotype following nutritional transition.

Burdge GC, et al. (2011) PLoS One.6(11):e28282.




Offspring phenotype and mRNA expression of genes involved in hepatic gluconeogenesis and ketogenesis.

Change in offspring body weight on day 70 compared to weaning, offspring energy intake on day 70, fasting glucose and β -hydroxybutyrate concentrations on postnatal day 70, Hepatic PPAR α , carnitine palmitoyltransferase-1 (CPT-1), glucocorticoid receptor (GR), phosphoenolpyruvate carboxykinase (PEPCK), (I) glucose-6-phosphatase (G-6-Pase) mRNA expression. Values are mean \pm SD for n = 5-7 rats per group. Values with different letters are significantly different ($P < 0.05$).

Transgenerational epigenetic inheritance of longevity in Caenorhabditis elegans.

Greer EL, et al. (2011) Nature. 19;479(7373):365-71.



Transgenerational epigenetic inheritance in plants.



Hauser MT, et al. (2011) Biochim Biophys Acta. 2011 Aug;1809(8):459-68.

Table 1
Impact of variations in the epigenetic status on plant growth and development.

| Plant species | Induction | Epigenetic status | Effect | Reference |
|--|------------------------------|---|--|-----------|
| Dryas octopetala DWARF1 | | DNA hypomethylation | Growth | [209,210] |
| Wheat Tritica aestivum | Glutenin gene | DNA methylation | Induced growth | [211] |
| Tobacco Nicotiana glauca | Transposon | Hypomethylation | Transgene silencing | [212] |
| Tobacco Nicotiana glauca | Transposon | Loss of expression of transgene, hypermethylation of 5'-UTR | Transgene silencing | [213] |
| Arabidopsis | Transposon | Loss of expression of <i>lpa</i> transgene | Hygromycin sensitivity | [214] |
| Peruvia | Transposon | Loss of expression of transgene, hypermethylation of 5'-UTR | Hygromycin sensitivity | [215] |
| Arabidopsis AGAMOUS | <i>met1, ddm1</i> background | DNA hypomethylation | Fiscal organ development, patterning | [216,217] |
| Silene latifolia | 5-azacytidine | DNA hypomethylation | Sex change to androecium-agamistosis | [218] |
| Arabidopsis SUPERMAN/dark kent | <i>met1, ddm1</i> background | DNA hypomethylation | Fiscal organ development, patterning | [219,220] |
| Arabidopsis FRI | Natural | Epigenetic with hypermethylated multiple FRI genes | FRI2 silencing results in late flowering plants | [221-224] |
| Utricularia vulgaris LLOYDIEA | Natural | Cytosine DNA methylation of glutelin genes | Fiscal development | [36] |
| Arabidopsis FLOWERING WAGENINGEN (FWA) | Natural | Loss of methylation, ectopic expression of FWA | Late flowering | [225] |
| Arabidopsis BAUCHE1 | <i>ddm1</i> background | Loss of methylation | Twisted leaves, dwarfing, and reduced fertility | [226] |
| Arabidopsis Flowering Receptor Locus (RLC) | Natural | H3K9 and H3K27 dimethylation at RLC locus after vernalization | Flower induction after vernalization | [227,228] |
| Rice Oryza sativa | 5-azacytidine | DNA hypomethylation, chromatin remodeling | Number of leaves, flowering age, stem height, biomass, growth | [229,230] |
| Rice Oryza sativa | Nutrient conditions, heat | DNA methylation; presence of insertion element | Biomass, growth | [136] |
| Tomato | Natural | SRP-box gene | Fruit ripening | [231] |
| Arabidopsis | Natural | Non-LTR retrotransposon <i>Sadha</i> is differentially expressed in natural accessions as G1 (active), C1 and L1 (inactive) | Fruit ripening | [232] |
| Arabidopsis BONSAI | <i>ddm1</i> background | DNA hypomethylation and silencing of APC13:RNS gene by nearby a long interspersed element | Growth | [233] |
| Melon Cucumis melon | Natural transposon insertion | Spreading of DNA methylation to WPI1 promoter | Sex determination, expression of WPI1 leads to carpel abortion | [234] |
| Wild potato Solanum nasutum | Species of hybrid origin | Phenotypic instability of flower development | Flower development | [235] |

Environmental heat and salt stress induce transgenerational phenotypic changes in Arabidopsis thaliana.

Suter L, Widmer A. (2013) PLoS One. 9;8(4):e60364.



Exposure to the environmental endocrine disruptor TCDD and human reproductive dysfunction: Translating lessons from murine models

Reprod Toxicol. 2017 Mar;68:59-71.
Bruner-Tran KL, Gnecco J, Ding T, Glore DR, Pensabene V, Osteen KG.

Pregnant Dam (F0 generation)



F1 fetus (contains the F2 germ cells)

Cumulative overview of reproductive characteristics in control and toxicant exposed female mice.^a


| Mouse History | Pregnancy Rate | Term Delivery ^b | Adenomyosis | Methylation of PR ^c |
|---------------|----------------|----------------------------|-------------|--------------------------------|
| Control | 95-100% | 100% | 0% | 0% |
| F1 | 46 | 37 | 73 | 86 |
| F3 | 50 | 60 | 56 | 33 |

^a Data compiled from references [10,101,102].
^b Delivery was considered term when occurring E19-E21 (plug date is considered E0.5).

^c Whole uteri collected from non-pregnant mice at estrus.

Transgenerational Effects of Di-(2-Ethylhexyl) Phthalate (DEHP) on Stress Hormones and Behavior.

Quinnies KM, Doyle TJ, Kim KH, Rissman EF.
Endocrinology. 2015 Sep;156(9):3077-83.



Mean (\pm SEM) corticosterone levels (ng/mL) measured in plasma from 150-mg/kg DEHP-female and control F3 mice.


(A) Under baseline conditions DEHP-lineage mice tended to have lower corticosterone levels than control females ($\#P = .10$). (B) After 15 minutes of restraint stress, DEHP-lineage females had lower corticosterone than control females; *, $P < .05$ (control-lineage females stressed, $n = 10$; control-lineage females baseline, $n = 9$; DEHP-lineage females stressed, $n = 8$; DEHP-lineage females baseline, $n = 10$; control-lineage males stressed, $n = 6$; control-lineage males baseline, $n = 6$; DEHP-lineage males stressed, $n = 8$; DEHP-lineage males baseline, $n = 7$).

The history of Distilbene® (Diethylstilbestrol) told to grandchildren--the transgenerational effect.

Fénichel P, Brucker-Davis F, Chevalier N.
Ann Endocrinol (Paris). 2015 Jul;76(3):253-9.

Table 1
In utero exposure to Distilbene®: historical.

| | |
|------|---|
| 1936 | DES Synthesis. Dodd and Larson. [22] |
| 1946 | Distilbene® advocated for spontaneous miscarriage and premature labour: 5 to 125 mg/day from 6WA to 35WA |
| 1953 | Randomised double-blind study $n = 840$ not superior to placebo. Dieckman. [3] |
| 1970 | 7 cases of clear cell adenocarcinoma (CCA) of the vagina and cervix in very young women exposed to DES in utero. Herbst [4] |
| 1972 | US ban by the FDA |
| 1975 | First case of CCA described in France |
| 1976 | Disappears from medical references as treatment for miscarriage in France |
| 1977 | Excluded for pregnancy |
| 1977 | Uterine defects related to DES. Kauffmann. [23] |
| 1981 | Pregnancy complications. Papiernik. [24] |
| 1990 | Genital abnormalities described in men |
| 2006 | RR $\times 3$ breast cancer after 50 years of age in second generation. Palmer. [11] |
| 2002 | Third generation hypospadias. Klip. [2] |
| 2009 | Epigenetic mechanisms of foetal programming after exposure to DES hypermethylation of HOX 10 gene in uterine cancer. Bruner. [25] |




Genital tract and reproductive characteristics in daughters of women and men prenatally exposed to diethylstilbestrol (DES)
 Wautier A, Tournaire M, Devouche E, et al.
 Therapie. Sep-Oct 2020;75(5):439-448.

Table 2 Uterine defects.

| Uterine defects N % | Third generation women | | | | | General population | | Second generation = prenatally exposed women |
|-------------------------------|------------------------|--------------------|-------------------------------|--------------------|--------------------|--------------------|---------------|--|
| | Our population | | | | | Fertile (f) | Infertile (g) | |
| | Total (a) | Exposed father (b) | Exposed father and mother (c) | Exposed mother (d) | Exposed mother (e) | | | Prenatally exposed women (h) |
| Population (aged 18 or older) | 759 (380) | 76 (34) | 11 (7) | 672 (339) | 28 | — | — | 60 |
| MRKH syndrome | 3 0.4% (0.8%) | 11.3% (2.9%) | 0 | 2 0.3% (0.6%) | 0 | — | 1/4900 | — |
| Doubling uterus | 3 0.4% (0.8%) | 0 | 0 | 3 0.4% (0.9%) | 0 | — | 1.2% | — |
| Bicornuate uterus | 6 0.8% (1.6%) | 0 | 0 | 6 0.9% (1.8%) | 0 | — | 1.7% | — |
| All uterine defects | 121.6% (3.2%) | 11.3% (2.9%) | 0 | 111.6% (3.2%) | 0 | 0.17% | 3.5% | 40 66% |


This table shows uterine defects in: third generation women = daughters of parents prenatally exposed to DES in our population (a,b,c,d), daughters of exposed mother in Kaufman et al., 2002 (e), general population (f, g), second generation = women prenatally exposed to DES (h), MRKH: Mayer-Rokitanski-Kuster-Hauser Syndrome.

Prenatal maternal infection promotes tissue-specific immunity and inflammation in offspring
 Lim AI, McFadden T, Link VM, et al.
 Science. 2021 Aug 27;373(6558):eabf3002.




Maternal infection promotes offspring intestine-specific immunity and inflammation. The direct response of fetal intestinal epithelial cells to IL-6 during maternal infection confers an enduring epigenetic memory to adult intestinal epithelial stem cells. As such, offspring epithelial cells exhibit enhanced reactivity toward the microbiota and heightened ability to control oral infection. However, these responses come at the cost of greater predisposition to gut inflammation.


Broad phenotypic impact of the effects of transgenerational heat stress in dairy cattle: a study of four consecutive generations.
 Weller JI, Ezra E, Gershoni M.
 Genet Sel Evol. 2021 Sep 6;53(1):69.




Transmission of chromatin states across generations in C. elegans.
 Özdemir I, Steiner FA.
 Semin Cell Dev Biol. 2021 Nov 22;S1084-9521(21)00284-6.



The Transgenerational Transmission of the Paternal Type 2 Diabetes-Induced Subfertility Phenotype
 Zatecka E, Bohuslavova R, Valaskova F, et al.
 Front Endocrinol (Lausanne). 2021 Nov 5;12:763863.



Polycystic ovary syndrome is transmitted via a transgenerational epigenetic process.
 Mimouni NEH, Paiva I, Barbotin A-L, et al.
 Cell Metab. 2021 Mar 2;33(3):513-530.e8.




Impact of Parental Exposure on Offspring Health in Humans.
 Jawaid A, Jehle K-L, Mansuy IM.
 Trends Genet. 2021 Apr;37(4):373-388.

Table 6. *In Utero* Exposure


| Type of disorder | Environmental exposure | Transmitted intergenerational/epigenetic/transgenerational phenotype | Associated epigenetic changes | Number of affected offspring/generations | Number of tested persons (n) | Refs |
|---------------------------------|---|---|-------------------------------|--|--|------|
| Nutritional/metabolic disorders | Changes in food availability in women | Increased risk of sudden death and cardiovascular mortality | Not assessed | 2 | 277 families | [63] |
| | Maternal LBW ^a and IUGR ^b | Increased risk of LBW and IUGR in offspring | Not assessed | 2 | 280 030 maternal observations 125 078 paternal observations | [64] |
| Toxicants | Maternal and grandmaternal smoking during pregnancy | Asthma | Not assessed | 2 | 4082 offspring | [65] |
| | Grandmaternal smoking during pregnancy | Asthma (independent of mother's smoking status) | Not assessed | 2 | 53 169 offspring | [66] |
| | Grandmaternal smoking during pregnancy | Asthma within the first 6 years of life (persistent childhood asthma phenotype) | Not assessed | 2 | 44 593 grandmothers 46 107 mothers 65 956 offspring | [67] |
| | Grandmaternal smoking during pregnancy | Increased birth weight, birth length, and body mass index of grandsons | Not assessed | 2 | 12 707 parents 9677 children | [68] |

^aLBW, low birth weight.
^bIUGR, intrauterine growth retardation.



Fetal Basis of Adult Onset Disease

| | |
|------------|----------------|
| Embryo | Most Sensitive |
| Postnatal | Sensitive |
| Pubertal | Less Sensitive |
| Adult | Insensitive |
| Aged Adult | Most Disease |



Summary


- Epigenetic Transgenerational Inheritance
 - Non-Genetic Form Inheritance
 - Toxicology / Environmental
 - Molecular Mechanism for Disease Etiology and DOHAD
 - All Epigenetic Processes Involved and Integrated in Germline
 - Generational Toxicology

Environmental Epigenetics, Disease and Evolution

- Integration Epigenetics and Genetics Essential Biology
- Evolution and Disease Etiology Requires Inclusion Epigenetics
- Doom and Gloom/ Biomarkers & Preventative Medicine

Glyphosate Induced Transgenerational DNA Methylation and Histone Retention Sperm Epigenetic Biomarkers for Disease


Ben Maamar M, Beck D, Thorson JLM, Nilsson E, Kubsad D, Skinner MK (2020) Epigenetics 9: 1-18




Environmental Induced Epigenetic Transgenerational Inheritance of Pathology: Systems Epigenetics in Disease Etiology and Generational Toxicology
 Daniel Beck, Eric E. Nilsson, Millissia Ben Maamar, Michael K. Skinner (Sci Reports 2022, 12:5452)

Exposure Specific DMRs
Exposure Induced F3 Generation Sperm DMRs at edger p-value <1e-06


| Jet fuel | Pesticides | Plastics | Dioxin | Methoxy | Glyphosate | Atrazine |
|----------|------------|----------|---------|---------|------------|----------|
| p<1e-06 | p<1e-06 | p<1e-06 | p<1e-06 | p<1e-06 | p<1e-04 | p<1e-06 |
| 31,517 | 18,791 | 168,543 | 37,475 | 6188 | 1395 | 3166 |




Transgenerational Disease Specific DMRs are Distinct



Pathway Associated Gene Networks for Kidney Disease




Novel Systems Epigenetics Disease Etiology Paradigm




Sperm DNA Methylation Epimutation Biomarkers for Male Infertility and FSH Therapeutic Responsiveness
Luján S, Caroppo E, Niederberger C, Arce J-C, Sadler-Riggleman I, Beck D, Nilsson E, Skinner MK
Scientific Reports (2019) 9:16786

Infertility Sperm DMR Chromosomal Locations



PCA Infertility DMR Signature Analysis




Sperm DNA Methylation Epimutation Biomarker for Paternal Offspring Autism Susceptibility
Garrido N, et al., and Skinner MK
Clinical Epigenetics 2021 (13:6)

DMR Identification


| p-value | All Window | Multiple Window |
|---------|------------|-----------------|
| 0.001 | 13274 | 255 |
| 1e-04 | 3214 | 31 |
| 1e-05 | 805 | 6 |
| 1e-06 | 223 | 1 |
| 1e-07 | 60 | 0 |

Significant windows


| | 1 | 2 | 3 |
|---------------|-----|---|---|
| Number of DMR | 799 | 3 | 3 |



Autism versus Control DMR PCA



DMR Associated Genes and Autism




Preterm Birth Buccal Cell Epigenetic Biomarkers to Facilitate Preventative Medicine
Paul Winchester, Eric Nilsson, Daniel Beck, and Michael K. Skinner
(In Review)

Mother DMFs

| p-value | All Window | Multiple Window |
|---------|------------|-----------------|
| 0.001 | 601 | 53 |
| 1e-04 | 165 | 28 |
| 1e-05 | 56 | 17 |
| 1e-06 | 32 | 12 |
| 1e-07 | 20 | 9 |

Significant windows


| | 1 | 2 | 3 | 4 | 5 |
|-------------------------|-----|----|---|---|---|
| Number of DMR (p<1e-04) | 137 | 18 | 5 | 2 | 3 |





Extended Overlaps

| | p<1e-04 | p<0.05 | Mother | Father | Female Child | Male Child |
|--------------|------------|-----------|------------|-----------|--------------|------------|
| Mother | 165 (100%) | 31 (49%) | 36 (56%) | 50 (30%) | | |
| Father | 54 (74%) | 70 (100%) | 37 (64%) | 34 (47%) | | |
| Female Child | 79 (58%) | 62 (46%) | 136 (100%) | 45 (34%) | | |
| Male Child | 17 (28%) | 15 (25%) | 11 (18%) | 81 (100%) | | |

Mother DMR Associated Gene Correlations



Environmental Factors → Gestating Mother (F0)



Abnormalities
Transcriptome
Gene Networks
Epimutations

Environment →

“Systems Biology of Reproduction”

Spring 2022 (Even Years) – Course Syllabus

BIOL 475/575 Level Undergraduate/Graduate (3 Credit)

SLN: (475) – 05504, (575) – 05505

Time - Tuesday and Thursday 10:35 am-11:50 am

Course Lectures in person and on Canvas/Panopto and Discussion Sessions in person and on WSU Zoom for all campuses

Room – CUE 418

Course Director – Michael Skinner, Abelson Hall 507, 335-1524, skinner@wsu.edu

Co-Instructor – Eric Nilsson, Abelson Hall 507, 225-1835, nilsson@wsu.edu

Learning Objective -

Current literature based course on the Systems Biology of Reproduction. Learning Systems approaches to the biology of reproduction from a molecular to physiological level of understanding.

Schedule/Lecture Outline –

| | | | |
|----------|---------|---------|---|
| January | 11 & 13 | Week 1 | Systems Biology Introduction |
| | 18 & 20 | Week 2 | Molecular/ Cellular/ Reproduction Systems |
| | 25 & 27 | Week 3 | Sex Determination Systems |
| February | 1 & 3 | Week 4 | Male Reproductive Tract Development & Function |
| | 8 & 10 | Week 5 | Female Reproductive Tract Development & Function |
| | 15 & 17 | Week 6 | Gonadal Developmental Systems Biology |
| | 22 & 24 | Week 7 | Testis Systems Biology |
| March | 1 & 3 | Week 8 | Ovary Systems Biology |
| | 8 & 10 | Week 9 | Epigenetics and Transgenerational Gonadal Disease |
| | 14 – 18 | Week 10 | Spring Break |
| | 22 & 24 | Week 11 | Gametogenesis/ Stem Cells/ Cloning |
| | 29 & 31 | Week 12 | Hypothalamus-Pituitary Development & Function |
| April | 5 & 7 | Week 13 | Reproductive Endocrinology Systems |
| | 12 & 14 | Week 14 | Fertilization & Implantation Systems |
| | 19 & 21 | Week 15 | Fetal Development & Birth Systems |
| | 26 & 28 | Week 16 | Assisted Reproduction/Contraception |
| May | 3 & 5 | Week 17 | Exam or Grant Review |